

ANTICANCER RESEARCH

International Journal of Cancer Research and Treatment

Editorial Office: International Institute
of Anticancer Research,
1st km Kapandritiou - Kalamou Rd.,
Kapandriti, P.O.Box 22, Attiki 19014,
Greece
Fax: 0030-22950-53389
Tel: 0030-22950-52945
e-mail: journals@iiar-anticancer.org

Dear Sir/Madam,

Please find the enclosed galley proofs of your article. Below are instructions on how to help facilitate the timely publication of your article in ANTICANCER RESEARCH.

1. Please read, correct and return the proofs to the Editorial Office within 2-3 working days.
2. Proofs may be Faxed, e-mailed or returned by the fastest mail available. Delays in the return of your proofs may necessitate transferring your paper to a later issue of the journal.
3. While reviewing your manuscript, please be on the lookout for unintended changes in the meaning of the text through editorial corrections or language improvements.
4. With the exception of the above, corrections should be limited to typographical errors.
5. To promote our rapid publication policy, figures are not sent to authors unless alterations have been made through the editing process.
6. **Should you require extra reprints of your article, please order them now.** Reprints ordered at this time will reach you considerably faster than if ordered after the publication of your article.
7. For further information on your article (publication date, volume, page numbers *etc.*), please contact the Editorial Office by Fax/Tel: 0030-22950-53389, e-mail: journals@iiar-anticancer.org, or by sending a letter to the Editorial Office: International Institute of Anticancer Research, 1st km Kapandritiou - Kalamou Rd., Kapandriti, P.O.Box 22, Attiki 19014, Greece.
8. In all correspondence with us please provide your manuscript number, your complete postal address, telephone/fax numbers and e-mail address.

Thank you for taking the time to study these guidelines.

Yours,



J.G. Delinassios
Managing Editor

Enclosures

Analysis of EGFR Overexpression, *EGFR* Gene Amplification and the EGFRvIII Mutation in Portuguese High-grade Gliomas

MARTA VIANA-PEREIRA¹, JOSÉ MANUEL LOPES^{2,3}, SUZIE LITTLE⁴, FERNANDA MILANEZI^{1,2},
DIANA BASTO¹, FERNANDO PARDAL⁵, CHRIS JONES⁴ and RUI MANUEL REIS¹

¹Life and Health Sciences Research Institute (ICVS), School of Health Sciences, University of Minho, Braga;

²IPATIMUP, Porto;

³Department of Pathology, S. Jovo Hospital, Faculty of Medicine, University of Porto, Porto;

⁵Department of Pathology, S. Marcos Hospital, Braga, Portugal;

⁴Paediatric Oncology, Institute of Cancer Research, Sutton, U.K.

Abstract. *Background:* Patients with malignant gliomas do not respond to any current therapy. Epidermal growth factor receptor (EGFR) controls several oncogenic processes, being frequently up-regulated in gliomas due to overexpression, gene amplification and gene mutation. EGFR inhibitors are being tried in gliomas, yet the molecular determinants of therapeutic response are unclear. *Materials and Methods:* EGFR overexpression, EGFRvIII mutation and EGFR amplification were determined by immunohistochemistry and chromogenic in situ hybridization (CISH) in 27 primary glioblastomas (GBM), 24 anaplastic oligodendrogliomas (AO) and four anaplastic oligoastrocytomas (AOA). *Results:* EGFR overexpression was associated with EGFR amplification, being found in 48% and 53% GBM, 33% and 40% AO and 75% and 67% AOA, respectively. EGFRvIII was found in 22% GBM, 8% AO and was absent in AOA. No association was observed between EGFR alterations and patient survival. *Conclusion:* We characterized, for the first time, EGFR molecular alterations in Portuguese patients with malignant glioma and identified a subpopulation of patients presenting putative biomarkers for EGFR-based therapies.

Gliomas are the most frequent primary central nervous system (CNS) tumors (1). According to the World Health Organization (WHO), gliomas are histologically divided into astrocytic, oligodendroglial and mixed oligoastrocytic

Correspondence to: Rui Manuel Reis, Life and Health Sciences Research Institute (ICVS), School of Health Sciences, University of Minho, Campus de Gualtar, 4710-057 Braga, Portugal. Tel: +351 253604825, Fax: +351 253604820, e-mail: rreis@ecsau.de.uminho.pt

Key Words: Amplification, EGFR, EGFRvIII, glioblastoma, glioma, oligodendrogloma, overexpression.

tumors, and are classified into four grades of malignancy (1, 2). Oligodendrogliomas and oligoastrocytomas are stratified into grade II and grade III (anaplastic) tumors; on the other hand astrocytomas can be subdivided into grade II, grade III and grade IV (2). The most malignant form (WHO grade IV), glioblastoma (GBM), is also the most frequent glioma subtype (1, 2). GBMs can be divided into primary glioblastomas, which arise *de novo* and are molecularly characterized by epidermal growth factor receptor (EGFR) overexpression/amplification, and secondary glioblastomas, which are derived from lower-grade astrocytomas and are characterized by TP53 mutations (3). The prognosis of patients with GBM is very poor, with survival usually less than twelve months (1, 4). Recently, the introduction of temozolomide-based chemotherapy in concomitancy with radiotherapy, led to increased GBM patient survival (15 months) (4). However, these results are far from being satisfactory and there are still patients that do not respond favorably to any described therapy. Therefore, it is necessary to understand the molecular features of gliomas in order to identify novel and effective therapeutic targets.

Receptor protein tyrosine kinases (RTKs) are major regulators of cell growth signaling and their importance in tumorigenesis and progression has been extensively investigated (5). Epidermal growth factor receptor (EGFR)/HER1 is a member of the class I epidermal growth factor family, which also includes HER2, HER3 and HER4. EGFR is a transmembrane glycoprotein that is stimulated by growth factors, namely the transforming growth factor- α (TGF- α) and EGF ligands, which bind to the extracellular domain of the receptor (6). Ligand binding to EGFR activates the receptor through dimerization leading to signal transduction and activation of downstream intracellular pathways, mainly RAS/RAF/MAPK, PI3K/AKT and STATs, that regulate cellular proliferation, differentiation, migration and survival (5). Several mechanisms of aberrant

EGFR activation have been reported in cancer cells, namely overexpression of the ligand EGF and of the receptor, *EGFR* gene amplification and activating mutations. Moreover, dysregulated EGFR activation is known to act oncogenically, stimulating the growth and spread of cancer cells (5). Therefore, due to the paramount role of EGFR in tumorigenesis, several therapeutic strategies are being developed to target and inhibit signalling *via* the receptor (7-9). Monoclonal antibodies (mAbs) bind to the extracellular domain of EGFR inhibiting this region and blocking ligand binding. Cetuximab (IMC-C225, Erbitux) and Panitumumab (ABX-EGF, Vectibix) are EGFR-binding mAbs currently approved for the treatment of colorectal cancer (10). Alternatively, EGFR tyrosine kinase inhibitors (TKIs) bind to the intracellular tyrosine kinase domain, blocking kinase activity as well as the downstream signaling. Gefitinib (ZD1839, Iressa) and erlotinib (OSI-774, Tarceva), are EGFR TKIs already approved for non-small cell lung cancer (NSCLC) patients (11).

In high-grade gliomas, *EGFR* is the most frequently amplified oncogene, being present in 40% of primary GBM and poorly characterized in anaplastic astrocytomas and anaplastic oligodendrogliomas (AO) (12-14). Importantly, *EGFR* gene amplification is known to be associated with protein overexpression in gliomas (2). Additionally, in about half of GBM cases with *EGFR* amplification, the event is coupled with *EGFR* gene mutations, with EGFRvIII (also known as Δ EGFR and del2-7EGFR) the most common in GBM (2, 15, 16). This EGFR mutant oncoprotein lacks a portion of the extracellular ligand-binding domain as a result of a genomic deletion involving exons 2 to 7, resulting in a mutant protein unable to bind EGFR ligands, yet constitutively activated in a ligand-independent manner leading to overproliferation of cancer cells (6,17). EGFR represents an attractive therapeutic target in malignant gliomas and even though there are no anti-EGFR agents approved for glioma treatment at present, there are several ongoing clinical trials evaluating the efficacy of mAbs and TKIs in gliomas (7, 18, 19).

In the present study, we aimed to characterize the most common mechanisms (*EGFR* overexpression, gene amplification and EGFRvIII mutation) involved in EGFR activation in a Portuguese cohort of high-grade (astrocytic, oligodendroglial and mixed) gliomas and to identify the subset of patients that would potentially benefit from EGFR-targeted therapies.

Materials and Methods

Patients and tumor samples. Sixty-two formalin-fixed, paraffin-embedded samples of sporadic gliomas from 55 patients were retrieved from the Department of Pathology of the S. João Hospital, Porto and S. Marcos Hospital, Braga, Portugal, as well

as the available patients' clinical data, as described elsewhere (20). Tumor samples were classified according to the WHO classification of CNS tumors (2): as 31 primary GBM (WHO grade IV), 26 AO (WHO grade III), one oligodendroglioma (O) (WHO grade II) and four anaplastic oligoastrocytomas (AOA) (WHO grade III) (Tables I and II). Twenty-seven (49.1%) patients were male and 28 (50.9%) were female; the mean age was 56.1 years (range 27-79 years) (Tables I and II).

EGFR and EGFRvIII immunohistochemistry. Formalin-fixed, paraffin-embedded tissue sections were used in the immunohistochemical analysis. Previously documented mouse monoclonal anti-EGFR (clone 31G7, 1:100; Zymed[®] Laboratories Inc., South San Francisco, CA, USA) and mouse monoclonal anti-EGFRvIII antibody (clone G100, 1:100; Zymed[®] Laboratories Inc.) were used as primary antibodies (18, 21). Anti-EGFRvIII antibody specifically recognizes EGFRvIII and does not cross-react with the wild-type form (18).

Immunohistochemistry for EGFR was performed as described elsewhere (22). Briefly, after deparaffinization, sections were rehydrated and washed. Antigen retrieval was achieved by 20 min incubation at 37°C with a bacterial protease extracted from *Streptomyces griseus* (Sigma-Aldrich Co., St. Louis, MO, USA). Endogenous peroxidase activity was blocked using 3% H₂O₂ in methanol for 10 min and samples were then incubated with Ultra V block (Lab Vision Corporation, Fremont, CA, USA) for 10 min at room temperature (RT). After incubation with primary antibody, overnight at RT, the secondary biotinylated goat anti-polyvalent antibody (Lab Vision Corporation) was applied for 10 min followed by 10 min incubation with Streptavidine Peroxidase (Lab Vision Corporation) at RT.

For the EGFRvIII reaction, sections were deparaffinized, incubated in 0.3% H₂O₂ in methanol for 30 min to block endogenous peroxidase activity and rehydrated. Antigen retrieval was achieved by microwaving sections in 10 mM citrate buffer (pH 6.0) three times for 5 min at 700 W. Sections were then incubated with Ultra V block (Lab Vision Corporation) for 20 min at RT. After incubation with primary antibody overnight at 4°C, the biotinylated "universal" secondary antibody (Vector Laboratories, Burlingame, CA, USA) was applied for 30 min followed by R.T.U. Vectastain[®]Elite ABC reagent (Vector Laboratories) incubation for 45 min at 37°C. Both EGFR and EGFRvIII sections were incubated with the chromogen 3,3'-diaminobenzidine (DAB) (Ultravision Detection System Anti-polyvalent, HRP/DAB; DAKO Corporation, Carpinteria, CA, USA), for 10 min at RT. Haematoxylin counterstaining was performed in a Leica Auto Stainer XL (Meyer Instruments Inc., Houston, TX, USA).

A specimen of human skin and a human glioblastoma with documented expression of EGFRvIII were used as positive controls for EGFR and EGFRvIII, respectively (22,23). Neoplastic cells with membranous and/or cytoplasmic intense immunoreactivity were considered positively stained. Both the distribution and intensity of the immunoreactivity were semi-quantitatively scored as follows: - (0%), + (<10%), ++ (10-50%), and +++ (>50%). Samples with scores (-) and (+) were considered negative, and those with scores (++) and (+++) were considered positive as described elsewhere (22,23).

Chromogenic in situ hybridization (CISH). The presence of EGFR gene amplification was assessed with CISH using Spot-Light

Table I. Results of EGFR analysis in glioblastomas.

Case No.	Age (years)/gender	IHC - EGFR	CISH - EGFR Amplification	IHC - EGFRvIII
12	27/F	+	Not Ampl	-
13	54/F	+++	Ampl	-
14	41/M	-	Not Ampl	-
19	61/F	-	Not Ampl	-
20	77/M	+++	Ampl	-
21	67/F	-	Not Ampl	-
22	50/F	+	Ampl	++
23	73/M	+	Not Ampl	-
24	57/M	+++	Ampl	-
24†	57/M	+++	Ampl	++
26	28/M	+++	NC	-
29	51/F	+++	Ampl	-
30	69/M	-	NC	-
30†	69/M	-	ND	-
32	53/M	+++	NC	++
32†	53/M	++	ND	+
33	59/F	+++	Ampl	+
36	56/F	+++	Ampl	+++
37	66/M	-	NC	-
38	68/F	-	ND	-
40	60/F	-	NC	-
41	58/F	-	Not Ampl	-
42	60/M	+++	Ampl	-
43	73/M	+++	Ampl	-
45	66/F	++	Not Ampl	+++
48	54/M	+++	Not Ampl	-
48†	54/M	+	Not Ampl	-
94	66/F	+	NC	-
96	79/M	+	Ampl	-
97	72/M	-	Not Ampl	-
231	62/F	++	NC	+++

ND, not determined; NC, not conclusive; Ampl, amplified; Not Ampl, not amplified; †, recurrence.

amplification probes for EGFR (Zymed®Laboratories Inc.). CISH was performed using Spot-Light CISH Polymer Detection Kit (Zymed®Laboratories Inc.) in accordance with the manufacturer’s protocol, and as described elsewhere (22). Amplification was defined as more than 5 signals per nucleus in more than 50% of cancer cells, or when large gene copy clusters were seen. Signals were evaluated at x400 and x600 and at least 60 morphologically unequivocal neoplastic cells were counted for the presence of the gene probe signals. CISH hybridizations were evaluated in a blinded manner, by two independent observers, on a multi-headed microscope.

Statistical analysis. Statistical analysis was performed using SPSS for Windows (version 14.0; SPSS Inc., Chicago, IL, USA). The correlation between categorical variables was calculated for statistical significance using Pearson’s chi-square test and the threshold for significance was $p \leq 0.05$.

Follow-up information was available for 40 out of 55 patients, with follow-up periods ranging from 1 to 93 months (median 11

Table II. Results of EGFR analysis in anaplastic oligodendrogliomas and oligoastrocytomas.

Case No.	Age (years)/gender	IHC - EGFR	CISH - EGFR Amplification	IHC - EGFRvIII
Anaplastic oligodendrogliomas				
60	65/M	++	Not Ampl	-
62	47/F	+	Not Ampl	-
64	65/F	+	NC	-
66	50/M	+	Not Ampl	-
68	36/F	-	Not Ampl	-
70	60/F	-	NC	-
71	53/M	++	Not Ampl	-
72	65/M	+++	Ampl	++
73	47/M	-	Not Ampl	-
76	70/F	-	Not Ampl	-
81	64/F	+	NC	-
83	54/M	+	Not Ampl	-
110	44/M	-	Not Ampl	-
110†	44/M	-	Not Ampl	-
112	42/F	+	Ampl	-
115	40/M	++	Not Ampl	-
116	51/M	-	Ampl	+++
118	64/F	++	Ampl	-
119	68/F	+	Ampl	-
163	51/M	+++	Ampl	-
168‡	36/M	+	ND	ND
168†	36/M	-	Not Ampl	+
170	73/M	-	Not Ampl	-
172	45/M	+++	ND	ND
172†	45/M	+++	Ampl	-
202	65/F	+++	Ampl	-
259	52/F	-	NC	-
Anaplastic oligoastrocytomas				
63	46/F	+	Not Ampl	-
203	36/F	++	NC	-
205	33/M	++	Ampl	-
206	55/F	+++	Ampl	+

ND, not determined; NC, not conclusive; Ampl, amplified; Not Ampl, not amplified; †, recurrence; ‡, oligodendroglioma grade II.

months, mean 16 months). Survival duration was defined as the time between diagnosis and death. Associations among EGFR expression, amplification, EGFRvIII expression and patients’ age with patients’ survival were assessed using Cox proportional hazards regression analyses.

Results

EGFR protein overexpression. EGFR immunohistochemical analysis was performed on 62 glioma samples from 55 patients. Results are summarized on Table I and Table II. The EGFR neoplastic staining was membranous and/or cytoplasmic, without immunoreactivity of endothelial tumor cells (Figure 1 A




Figure 1. EGFR cytoplasmic/membranous immunohistochemistry expression in GBM (A, x200) and AO (B, x400). EGFR amplification revealed by high nuclear signaling in GBM (C, x100) and AO (D, x100; inset, x600). EGFRvIII cytoplasmic/membranous immunohistochemistry expression in GBM (E, x200) and AO (F, x400).

and B). EGFR overexpression (2+/3+) was detected in 24 of 55 tumors (44%). A high percentage of positive cases was found in all histological types of gliomas analyzed,

namely, 48% (13/27) of GBM (Table I, Figure 1A), 33% (8/24) of AO and 75% (3/4) of AOA (Table II, Figure 1B).

EGFR amplification. CISH analysis was possible in 54 tumors and was conclusive in 42 (78%) of the cases (Tables I and II). The pattern of EGFR signals observed using CISH was compatible with double minute amplification (Figure 1C and D). Overall, *EGFR* gene amplification was detected in 20 (48%) high-grade gliomas. Specifically, *EGFR* gene amplification was observed in 53% of GBM (10/19) (Table I, Figure 1C), 40% of AO (8/20) (Table II, Figure 1D) and 67% of AOA (2/3) (Table II). Of the 20 gliomas with *EGFR* amplification, 15 (75%) exhibited EGFR overexpression ($p=0.001$). All recurrent cases with available CISH analysis for both primary and recurrent samples, namely two GBM (cases 24 and 48) and one AO (case 110), showed the same *EGFR* amplification status.

EGFRvIII protein overexpression. EGFRvIII expression analysis was performed in 60 samples of 55 gliomas and results are summarized in Table I and Table II. EGFRvIII staining was predominantly cytoplasmic and was observed only in neoplastic cells, not in tumor endothelial tumor cells (Figure 1E and F).

EGFRvIII overexpression was observed in 22% of GBM (6/27) (Table I, Figure 1E), 8% of AO (2/24) (Table II, Figure 1F) and was absent in AOA (0/4) (Table II). In GBM, all but one sample with EGFRvIII overexpression also exhibited EGFR overexpression (Table I, $p=0.114$). Similarly, with the exception of one case, all informative samples with EGFRvIII overexpression also showed *EGFR* gene amplification (Table I, $p=0.596$). Both AO cases with EGFRvIII overexpression depicted *EGFR* gene amplification, (Table II, $p=0.068$) one of them also with EGFR overexpression (Table II, $p=0.602$).

Clinical significance of EGFR overexpression, EGFR amplification and EGFRvIII overexpression. Association between patient age and EGFR overexpression was assessed for glioma, GBM and AO patients (data not shown). EGFR overexpression in GBM patients tended ($p=0.082$) to occur in younger patients. Multivariate analysis (Cox proportional hazards) was used to determine the association between patient age, EGFR overexpression, *EGFR* amplification or EGFRvIII overexpression and overall survival for GBM and AO. The median age of patients was calculated for GBM (60 years) and AO (51 years) and these values were used to split patients into younger and older groups. Survival time was calculated after one year for GBM patients and after five years for AO patients. No correlation was found between EGFR overexpression, *EGFR* amplification, or EGFRvIII overexpression and overall survival of GBM and AO patients (Table III). EGFR overexpression showed a tendency ($p=0.054$, Table III) for being associated with shorter survival in AO patients. There was a significant

Table III. Multivariate analysis for the effect of EGFR overexpression, EGFR amplification, EGFRvIII overexpression and age on survival of GBM and AO patients.

Variable	Hazard ratio (95% confidence interval)	p-value
<i>GBM (1-year survival)</i>		
EGFR overexpression	3.655 (0.731-18.277)	0.114
EGFR amplification	1.073 (0.167-6.885)	0.941
EGFRvIII overexpression	0.732 (0.131-4.095)	0.723
Median age (<60 or ≥60 years)	1.222 (0.204-7.333)	0.826
<i>AO (5-year survival)</i>		
EGFR overexpression	36.927 (0.942-1447.576)	0.054
EGFR amplification	0.112 (0.007-1.710)	0.116
EGFRvIII overexpression	NS	
Median age (<51 or ≥51 years)	0.065 (0.007-0.570)	0.014

NS: Insufficient data for statistical evaluation.

correlation ($p=0.014$, Table III) between being older and better survival in AO patients.

Discussion

In this study, we characterized, for the first time, the presence of EGFR alterations in a series of high grade gliomas from Portuguese patients. We studied a total of 31 GBM and observed that the frequency of GBM presenting EGFR overexpression (48%), *EGFR* amplification (53%) and EGFRvIII overexpression (22%) were in line with the published literature (24-26). Similarly to several studies a positive correlation ($p=0.012$) was found between EGFR overexpression and gene amplification in GBM (14, 18, 24, 27). Additionally, all but one GBM with EGFRvIII overexpression presented gene amplification, which is also in agreement with previous studies (24, 25). It was previously reported that older patients with GBM have higher rates of EGFR overexpression and amplification (24, 28). We found no statistical differences between patient age and EGFR overexpression, *EGFR* gene amplification, or EGFRvIII overexpression in the same cases.

EGFR activation status in anaplastic oligodendrogliomas has been less frequently reported (13, 27, 29, 30). We analyzed 24 cases and observed EGFR overexpression in 33%, *EGFR* amplification in 40% and EGFRvIII mutation in 8% of anaplastic oligodendrogliomas. EGFR overexpression tended to correlate with gene amplification ($p=0.094$). Other authors described *EGFR* amplification in anaplastic oligodendrogliomas and frequencies varied from 0-42.5% (13, 27, 31). A possible reason for this discrepancy is the distinct methodologies used. Regarding EGFRvIII expression, results are few and contradictory. Wikstrand and colleagues analyzed 5 anaplastic oligodendrogliomas and described a frequency of

20% of EGFRvIII expression (30); however, in another study from the same group EGFRvIII was not found to be expressed in 25 anaplastic oligodendrogliomas (29).

The influence of EGFR overexpression, *EGFR* gene amplification and EGFRvIII overexpression in patient prognosis has been highly controversial for gliomas (24, 25, 28, 32-37). To clarify this issue, we performed a multivariate analysis and found no association between these EGFR alterations and patients' overall survival in our cohort.

EGFR is becoming an important therapeutic target, with some anti-EGFR drugs already being used in clinical practice and several novel EGFR inhibitors under development (38). Among the two major approaches, using EGFR-TKIs and mAbs, the former seems to be more suitable for gliomas due to their low molecular weight potentially being better at overcoming the blood-brain barrier (39). In NSCLC, the presence of activating mutations in the EGFR kinase domain was associated with selective EGFR-TKI sensitivity, allowing the selection of patients with a higher probability of clinical response to gefitinib and erlotinib (11). However, these mutations have never been found in glioma cell lines (11), or in glioma patients (11, 40, 41). Recently, Lee and colleagues reported EGFR activation in GBM due to missense mutations in the EGFR extracellular domain (42). They reported that transformed cells with the EGFR ectodomain mutations had increased sensitivity to erlotinib, however studying DNA samples from a previous clinical trial, these authors were unable to associate EGFR ectodomain mutations with clinical responses to EGFR TKI inhibitors (42).

The few clinical trials with gefitinib and erlotinib in gliomas included a small number of patients and variable molecular markers, insufficient for conclusive results regarding patients' response and *EGFR* molecular status. Rich and colleagues, in a phase II trial, described a monotherapy study with gefitinib in GBM patients and found neither objective tumor response nor association between EGFR expression, *EGFR* amplification, or EGFRvIII expression and gefitinib response (43). Franceschi and colleagues' recent phase II trial investigated the role of gefitinib in patients with high-grade gliomas and reported an 18% stable disease rate without any correlation of EGFR expression or gene status and tumor response (44). Vogelbaum *et al.* described a monotherapy with erlotinib in patients with recurrent GBM and found 25% with stable disease and 25% with partial response rates; although *EGFR* amplification was observed in about 50% of tumors, it was not associated with erlotinib response (45). In a phase II study with erlotinib in recurrent GBM, Cloughesy *et al.* described a 33% stable disease rate and EGFR expression associated with a slight tendency for better patient outcome (46). Haas-Kogan *et al.* reported that the presence of EGFR overexpression and gene

amplification, associated with low levels of activated Akt, were associated with response to erlotinib, suggesting that these molecular alterations could be predictive markers for EGFR-TKI sensitivity in gliomas (18). Mellinghoff *et al.* described that coexpression of EGFRvIII and phosphatase and tensin homolog (PTEN) (RTK downstream negative regulator) was associated with a better response to gefitinib and erlotinib in recurrent GBM (19). In general, results with erlotinib seem to be more promising than with gefitinib, possibly due to its targeting of both EGFRvIII mutant and wild-type EGFR (39).

There have also been clinical trials involving EGFR-targeted mAbs in gliomas (9, 47). A phase I/II clinical trial using the EGFR mAb h-R3 in malignant glioma enrolled 29 patients with an objective response in 37.9% and stable disease in 41.4% of the cases (47). Another phase I/II trial using cetuximab in GBM is ongoing (9). There are several *in vitro* and *in vivo* studies reporting EGFR-targeted mAbs effects in glioma. Recently, Johns *et al.* studied the efficacy of two EGFR-specific mAbs (mAbs 806 and 528) against glioma cell line-derived xenografts expressing EGFR and EGFRvIII and reported that their efficacy was dependent on EGFR overexpression and receptor activation status (48). Currently, considerable efforts are being made to design anti-EGFRvIII strategies, such as mAbs and vaccines, since this mutated form constitutes a tumor-specific target that is not present in normal cells.

In conclusion, in the present study, we found that a high percentage of GBM and AO exhibited EGFR overexpression and amplification, as did a significant proportion of GBM and a small proportion of AO expressing EGFRvIII. Our results represent the first step for the identification of Portuguese glioma patients who could respond to specific therapies targeting EGFR.

References

- Behin A, Hoang-Xuan K, Carpentier AF and Delattre JY: Primary brain tumours in adults. *Lancet* 361: 323-331, 2003.
- Louis DN, Ohgaki H, Wiestler OD, Cavenee WK, Burger PC, Jouvet A, Scheithauer BW and Kleihues P: The 2007 WHO classification of tumours of the central nervous system. *Acta Neuropathol* 114: 97-109, 2007.
- Benjamin R, Capparella J and Brown A: Classification of glioblastoma multiforme in adults by molecular genetics. *Cancer J* 9: 82-90, 2003.
- Stupp R, Mason WP, van den Bent MJ, Weller M, Fisher B, Taphoorn MJ, Belanger K, Brandes AA, Marosi C, Bogdahn U, Curschmann J, Janzer RC, Ludwin SK, Gorlia T, Allgeier A, Lacombe D, Cairncross JG, Eisenhauer E and Mirimanoff RO: Radiotherapy plus concomitant and adjuvant temozolomide for glioblastoma. *N Engl J Med* 352: 987-996, 2005.
- Scaltriti M and Baselga J: The epidermal growth factor receptor pathway: A model for targeted therapy. *Clin Cancer Res* 12: 5268-5272, 2006.

- 6 Kuan CT, Wikstrand CJ and Bigner DD: EGF mutant receptor vIII as a molecular target in cancer therapy. *Endocr Relat Cancer* 8: 83-96, 2001.
- 7 Nicholas MK, Lukas RV, Jafri NF, Faoro L and Salgia R: Epidermal growth factor receptor-mediated signal transduction in the development and therapy of gliomas. *Clin Cancer Res* 12: 7261-7270, 2006.
- 8 Mendelsohn J and Baselga J: Epidermal growth factor receptor targeting in cancer. *Semin Oncol* 33: 369-385, 2006.
- 9 Combs SE, Heeger S, Haselmann R, Edler L, Debus J and Schulz-Ertner D: Treatment of primary glioblastoma multiforme with cetuximab, radiotherapy and temozolomide (GERT)- phase I/II trial: study protocol. *BMC Cancer* 6: 133, 2006.
- 10 Diasio RB and Fourie J: Targeting the epidermal growth factor receptor in the treatment of colorectal cancer - State of the art. *Drugs* 66: 1441-1463, 2006.
- 11 Lynch TJ, Bell DW, Sordella R, Gurubhagavatula S, Okimoto RA, Brannigan BW, Harris PL, Haserlat SM, Supko JG, Haluska FG, Louis DN, Christiani DC, Settleman J and Haber DA: Activating mutations in the epidermal growth factor receptor underlying responsiveness of non-small-cell lung cancer to gefitinib. *N Engl J Med* 350: 2129-2139, 2004.
- 12 Nieder C, Schlegel J, Andratschke N, Thamm R, Grosu AL and Molls M: The role of growth factors in central nervous system tumours. *Anticancer Res* 23: 1681-1686, 2003.
- 13 Reifenberger J, Reifenberger G, Ichimura K, Schmidt EE, Wechsler W and Collins VP: Epidermal growth factor receptor expression in oligodendroglial tumors. *Am J Pathol* 149: 29-35, 1996.
- 14 Schober R, Bilzer T, Waha A, Reifenberger G, Wechsler W, von Deimling A, Wiestler OD, Westphal M, Kemshead JT and Vega F: The epidermal growth factor receptor in glioblastoma: genomic amplification, protein expression, and patient survival data in a therapeutic trial. *Clin Neuropathol* 14: 169-174, 1995.
- 15 Schwachheimer K, Huang S and Cavenee WK: EGFR gene amplification-rearrangement in human glioblastomas. *Int J Cancer* 62: 145-148, 1995.
- 16 Rao RD, Uhm JH, Krishnan S and James CD: Genetic and signaling pathway alterations in glioblastoma: relevance to novel targeted therapies. *Front Biosci* 8: e270-e280, 2003.
- 17 Wikstrand CJ, Reist CJ, Archer GE, Zalutsky MR and Bigner DD: The class III variant of the epidermal growth factor receptor (EGFRvIII): characterization and utilization as an immunotherapeutic target. *J Neurovirol* 4: 148-158, 1998.
- 18 Haas-Kogan DA, Prados MD, Tihan T, Eberhard DA, Jelluma N, Arvold ND, Baumber R, Lamborn KR, Kapadia A, Malec M, Berger MS and Stokoe D: Epidermal growth factor receptor, protein kinase B/Akt, and glioma response to erlotinib. *J Natl Cancer Inst* 97: 880-887, 2005.
- 19 Mellingerhoff IK, Wang MY, Vivanco I, Haas-Kogan DA, Zhu S, Dia EQ, Lu KV, Yoshimoto K, Huang JH, Chute DJ, Riggs BL, Horvath S, Liau LM, Cavenee WK, Rao PN, Beroukhir R, Peck TC, Lee JC, Sellers WR, Stokoe D, Prados M, Cloughesy TF, Sawyers CL and Mischel PS: Molecular determinants of the response of glioblastomas to EGFR kinase inhibitors. *N Engl J Med* 353: 2012-2024, 2005.
- 20 Costa BM, Ferreira P, Costa S, Canedo P, Oliveira P, Silva A, Pardal F, Suriano G, Machado JC, Lopes JM and Reis RM: Association between functional EGF+61 polymorphism and glioma risk. *Clin Cancer Res* 13: 2621-2626, 2007.
- 21 Penault-Llorca F, Cayre A, Arnould L, Bibeau F, Bralet MP, Rochaix P, Savary J and Sabourin JC: Is there an immunohistochemical technique definitively valid in epidermal growth factor receptor assessment? *Oncol Rep* 16: 1173-1179, 2006.
- 22 Reis-Filho JS, Pinheiro C, Lambros MB, Milanezi F, Carvalho S, Savage K, Simpson PT, Jones C, Swift S, Mackay A, Reis RM, Hornick JL, Pereira EM, Baltazar F, Fletcher CD, Ashworth A, Lakhani SR and Schmitt FC: EGFR amplification and lack of activating mutations in metaplastic breast carcinomas. *J Pathol* 209: 445-453, 2006.
- 23 Biernat W, Huang H, Yokoo H, Kleihues P and Ohgaki H: Predominant expression of mutant EGFR (EGFRvIII) is rare in primary glioblastomas. *Brain Pathol* 14: 131-136, 2004.
- 24 Shinojima N, Tada K, Shiraiishi S, Kamiryō T, Kochi M, Nakamura H, Makino K, Saya H, Hirano H, Kuratsu J, Oka K, Ishimaru Y and Ushio Y: Prognostic value of epidermal growth factor receptor in patients with glioblastoma multiforme. *Cancer Res* 63: 6962-6970, 2003.
- 25 Aldape KD, Ballman K, Furth A, Buckner JC, Giannini C, Burger PC, Scheithauer BW, Jenkins RB and James CD: Immunohistochemical detection of EGFRvIII in high malignancy grade astrocytomas and evaluation of prognostic significance. *J Neuropathol Exp Neurol* 63: 700-707, 2004.
- 26 Marquez A, Wu R, Zhao J, Tao J and Shi Z: Evaluation of epidermal growth factor receptor (EGFR) by chromogenic *in situ* hybridization (CISH) and immunohistochemistry (IHC) in archival gliomas using bright-field microscopy. *Diagn Mol Pathol* 13: 1-8, 2004.
- 27 Ekstrand AJ, James CD, Cavenee WK, Seliger B, Pettersson RF and Collins VP: Genes for epidermal growth factor receptor, transforming growth factor alpha, and epidermal growth factor and their expression in human gliomas *in vivo*. *Cancer Res* 51: 2164-2172, 1991.
- 28 Simmons ML, Lamborn KR, Takahashi M, Chen P, Israel MA, Berger MS, Godfrey T, Nigro J, Prados M, Chang S, Barker FG and Aldape K: Analysis of complex relationships between age, p53, epidermal growth factor receptor, and survival in glioblastoma patients. *Cancer Res* 61: 1122-1128, 2001.
- 29 McLendon RE, Wikstrand CJ, Matthews MR, Al-Baradei R, Bigner SH and Bigner DD: Glioma-associated antigen expression in oligodendroglial neoplasms. Tenascin and epidermal growth factor receptor. *J Histochem Cytochem* 48: 1103-1110, 2000.
- 30 Wikstrand CJ, McLendon RE, Friedman AH and Bigner DD: Cell surface localization and density of the tumor-associated variant of the epidermal growth factor receptor, EGFRvIII. *Cancer Res* 57: 4130-4140, 1997.
- 31 Franco-Hernandez C, Martinez-Glez V, Alonso ME, de Campos JM, Isla A, Vaquero J, Gutierrez M and Rey JA: Gene dosage and mutational analyses of EGFR in oligodendrogliomas. *Int J Oncol* 30: 209-215, 2007.
- 32 Bigner SH, Burger PC, Wong AJ, Werner MH, Hamilton SR, Muhlbaier LH, Vogelstein B and Bigner DD: Gene amplification in malignant human gliomas: clinical and histopathologic aspects. *J Neuropathol Exp Neurol* 47: 191-205, 1988.
- 33 Ohgaki H and Kleihues P: Population-based studies on incidence, survival rates, and genetic alterations in astrocytic and oligodendroglial gliomas. *J Neuropathol Exp Neurol* 64: 479-489, 2005.

- 34 Mizoguchi M, Betensky RA, Batchelor TT, Bernay DC, Louis DN and Nutt CL: Activation of STAT3, MAPK, and AKT in malignant astrocytic gliomas: correlation with EGFR status, tumor grade, and survival. *J Neuropathol Exp Neurol* 65: 1181-1188, 2006.
- 35 Dirks PB and Rutka JT: Current concepts in neuro-oncology: the cell cycle – a review. *Neurosurgery* 40: 1000-1013, 1997.
- 36 Heimberger AB, Hlatky R, Suki D, Yang D, Weinberg J, Gilbert M, Sawaya R and Aldape K: Prognostic effect of epidermal growth factor receptor and EGFRvIII in glioblastoma multiforme patients. *Clin Cancer Res* 11: 1462-1466, 2005.
- 37 Pelloski CE, Ballman KV, Furth AF, Zhang L, Lin E, Sulman EP, Bhat K, McDonald JM, Yung WK, Colman H, Woo SY, Heimberger AB, Suki D, Prados MD, Chang SM, Barker FG 2nd, Buckner JC, James CD and Aldape K: Epidermal growth factor receptor variant III status defines clinically distinct subtypes of glioblastoma. *J Clin Oncol* 25: 2288-2294, 2007.
- 38 Roberts PJ and Der CJ: Targeting the Raf-MEK-ERK mitogen-activated protein kinase cascade for the treatment of cancer. *Oncogene* 26: 3291-3310, 2007.
- 39 Halatsch ME, Schmidt U, Behnke-Mursch J, Unterberg A and Wirtz CR: Epidermal growth factor receptor inhibition for the treatment of glioblastoma multiforme and other malignant brain tumours. *Cancer Treat Rev* 32: 74-89, 2006.
- 40 Barber TD, Vogelstein B, Kinzler KW and Velculescu VE: Somatic mutations of EGFR in colorectal cancers and glioblastomas. *N Engl J Med* 351: 2883, 2004.
- 41 Marie Y, Carpentier AF, Omuro AM, Sanson M, Thillet J, Hoang-Xuan K and Delattre JY: EGFR tyrosine kinase domain mutations in human gliomas. *Neurology* 64: 1444-1445, 2005.
- 42 Lee JC, Vivanco I, Beroukhi R, Huang JHY, Feng WL, DeBiasi RM, Yoshimoto K, King JC, Nghiemphu P, Yuza Y, Xu Q, Greulich H, Thomas RK, Paez JG, Peck TC, Linhart DJ, Glatt KA, Getz G, Onofrio R, Ziaugra L, Levine RL, Gabriel S, Kawaguchi T, O'Neill K, Khan H, Liao LM, Nelson SF, Rao PN, Mischel P, Pieper RO, Cloughesy T, Leahy DJ, Sellers WR, Sawyers CL, Meyerson M and Mellinghoff IK: Epidermal growth factor receptor activation in glioblastoma through novel missense mutations in the extracellular domain. *Plos Medicine* 3: 2264-2273, 2006.
- 43 Rich JN, Reardon DA, Peery T, Dowell JM, Quinn JA, Penne KL, Wikstrand CJ, Van Duyn LB, Dancey JE, McLendon RE, Kao JC, Stenzel TT, Rasheed BKA, Tourt-Uhlig SE, Herndon JE, Vredenburgh JJ, Sampson JH, Friedman AH, Bigner DD and Friedman HS: Phase II trial of gefitinib in recurrent glioblastoma. *J Clin Oncol* 22: 133-142, 2004.
- 44 Franceschi E, Cavallo G, Lonardi S, Magrini E, Tosoni A, Grosso D, Scopece L, Blatt V, Urbini B, Pession A, Tallini G, Crino L and Brandes AA: Gefitinib in patients with progressive high-grade gliomas: a multicentre phase II study by Gruppo Italiano Cooperativo di Neuro-Oncologia (GICNO). *Br J Cancer* 96: 1047-1051, 2007.
- 45 Vogelbaum MA, Peereboom D, Stevens G, Barnett G and Brewer C: Phase II trial of the EGFR tyrosine kinase inhibitor erlotinib for single agent therapy of recurrent glioblastoma multiforme: Interim results. *J Clin Oncol* 22: 121 (suppl.) (abstract), 2004.
- 46 Cloughesy T, Yung A, Vredenburgh J, Aldape K, Eberhard D, Prados M, Vandenberg S, Klencke B and Mischel P: Phase II study of erlotinib in recurrent GBM: Molecular predictors of outcome. *J Clin Oncol* 23: 115 (suppl.) (abstract), 2005.
- 47 Ramos TC, Figueredo J, Catala M, Gonzalez S, Selva JC, Cruz TM, Toledo C, Silva S, Pestano Y, Ramos M, Leonard L, Torres O, Marinello P, Perez R and Lage A: Treatment of high-grade glioma patients with the humanized anti-epidermal growth factor receptor (EGFR) antibody h-R3 - Report from a phase I/II trial. *Cancer Biol Ther* 5: 375-379, 2006.
- 48 Johns TG, Perera RM, Vernes SC, Vitali AA, Cao DX, Cavenee WK, Scott AM and Furnari FB: The efficacy of epidermal growth factor receptor-specific antibodies against glioma xenografts is influenced by receptor levels, activation status, and heterodimerization. *Clin Cancer Res* 13: 1911-1925, 2007.

Received November 9, 2007

Revised December 21, 2007

Accepted January 8, 2008

ANTICANCER RESEARCH

International Journal of Cancer Research and Treatment

ISSN: 0250-7005

January 8, 2008

Dr. Rui Manuel Reis

Re: Your manuscript No. **8478-V** entitled « **Analysis of EGFR Overexpression...**»

Dear Dr

Referring to your above manuscript for publication in AR, please allow us to use this form letter in reply:

1. *Referee' s recommendations:*

- Urgent to be published immediately.
- Accepted in the presented form.
- Accepted with minor changes.
- Accepted with grammatical or language corrections.
- Remarks:

2. *Excess page charges.*

- Your article has approx. **8** printed pages and is in excess of the allotted number by approx. **4** printed pages. The charges are EURO € **180** per excess page, totalling EURO € **504 (-30% Discount)**
We ask you to confirm acceptance of these charges.
- Your article includes **1** pages with color figures. The charges are EURO € **650** per color page, totalling EURO € **650**
- Our invoice is sent by air mail to the corresponding author.

3. Your article will appear in Volume **28**, Issue No. **2**, **2008**

4. Please order your reprints now. This will facilitate our prompt planning of future issues and rapid publication of your article. Reprints will be delivered by air mail within one month from publication.

We would appreciate your prompt reply.

With many thanks,

Yours sincerely,



J.G. Delinassios
Managing Editor

EDITORIAL OFFICE: INTERNATIONAL INSTITUTE OF ANTICANCER RESEARCH
1st km Kapandritiou - Kalamou Rd., Kapandriti, P.O.B. 22, Attiki 19014, Greece. Tel.: 0030-22950-52945;
Tel & Fax:0030-22950-53389; e-mail: journals@iiar-anticancer.org

ANTICANCER RESEARCH

International Journal of Cancer Research and Treatment

ISSN: 0250-7005

Editorial Office: International Institute of Anticancer Research,
1st km Kapandritiou-Kalamou Rd., Kapandriti, P.O.B. 22, Attiki 19014, Greece
Fax: 0030-22950-53389; Tel: 0030-22950-52945; e-mail: journals@iiar-anticancer.org

Please type or print the requested information on the reprint order form and return it to the Editorial Office by fax or e-mail.

Reprints must be paid for in advance.

If your paper is subject to charges for excess pages or color plates, please add these charges to the payment for reprints.

The reprints are not to be sold.

PRICE LIST FOR REPRINTS WITHOUT COVER

Page length	PDF file	Number of copies requested (prices are in Euro)									
		100	200	300	400	500	1000	1500	2000	3000	5000
1-4pp	167	335	387	438	503	554	851	1135	1470	2038	3225
5-8	219	438	503	580	645	722	1083	1445	1832	2554	4012
9-12	277	554	619	709	787	877	1341	1780	2219	3096	4824
13-16	354	709	787	890	993	1096	1625	2141	2657	3676	5715
17-20	419	838	929	1032	1148	1277	1883	2451	3044	4244	6527

For reprints with cover: Please add EURO 140.00 per 100 copies.

Postage: Please add 4% on the above prices.

Reprint Order Form

Of my paper No. **8478-V** comprising **8** printed pages, entitled « **Analysis of EGFR Overexpression...** » accepted for publication in ANTICANCER RESEARCH Vol. **28** No. **2**

- I require a total of _____ copies at EURO _____
- I do not require reprints.
- Please send me a PDF file of the article at EURO _____.
- Please send me a copy of this issue containing my paper at EURO 45.00.
- Please enter my personal subscription to ANTICANCER RESEARCH at the special Author' s price of EURO 405.00 (Year: 2008).
- A check for the above amounts payable to J.G. Delinassios, Executive Publisher of Anticancer Research Journal, is enclosed.
- Please send an invoice to:

For EC countries: Please give your VAT number:

City and Date:

Exact postal address:

Tel:

Fax:

Signature: