

INTERNATIONAL PUBLICATIONS USA

PanAmerican Mathematical Journal
Volume 24(2014), Number 1, 75–92

The Stability of a Generalized Radical Reciprocal Quadratic Functional Equation in Felbin's Space

K. Ravi

Sacred Heart College
Department of Mathematics
Tirupattur-635 601, TamilNadu, India
shckravi@yahoo.co.in

J.M.Rassias

National and Capodistrian University of Athens
Pedagogical Department E.E
Section of Mathematics and Informatics
4, Agamemnonos Str, Aghia Paraskevi
Athens, Attikis 15342, Greece
jrassias@primedu.uoa.gr

Sandra Pinelas

Academia Militar
Departamento de Ciências Exactas e Naturais
Av.Conde Castro Guimarães, 2720-113 Amadora, Portugal
sandra.pinelas@gmail.com

P.Narasimman

Jeppiaar Institute of Technology
Department of Mathematics
Kunnam, Sriperumbudur(TK), Chennai- 631 604, India
simma_gtec@yahoo.co.in

Communicated by Allan Peterson

(Received December 1, 2013; Revised Version Accepted February 17, 2014)

Abstract

In this paper, the authors introduce a new generalized radical reciprocal quadratic functional equation of the form

$$f\left(\frac{\sqrt{x^2+y^2}}{n}\right) \pm mf\left(\sqrt{x^2+y^2}\right) = (n^2 \pm m) \frac{f(x)f(y)}{f(x)+f(y)}$$

and investigate its generalized Hyers-Ulam stability in Felbin's type fuzzy normed linear spaces. Also we give counter example for the Hyers-Ulam-Rassias stability of the radical reciprocal quadratic functional equation for some cases.

AMS (MOS) Subject Classification: 39B52, 39B55, 39B62, 39B82

Key words: Reciprocal functional equations, Felbin's Space, Generalized Hyers-Ulam stability.

1 Introduction

The stability problem of functional equations originates from the fundamental question: When is it true that a mathematical object satisfying a certain property approximately must be close to an object satisfying the property exactly?

In connection with the above question, in 1940, S.M. Ulam [36] raised a question concerning the stability of homomorphisms. Let G be a group and let G' be a metric group with $d(.,.)$. Given $\epsilon > 0$ does there exist a $\delta > 0$ such that if a function $f : G \rightarrow G'$ satisfies the inequality $d(f(xy), f(x)f(y)) < \delta$ for all $x, y \in G$, then there is a homomorphism $H : G \rightarrow G'$ with $d(f(x), H(x)) < \epsilon$ for all $x \in G$?

The first partial solution to Ulam's question was given by D.H. Hyers [9]. He considered the case of approximately additive mappings $f : E \rightarrow E'$ where E and E' are Banach spaces and f satisfies Hyers inequality

$$\|f(x+y) - f(x) - f(y)\| \leq \epsilon$$

for all $x, y \in E$, it was shown that the limit $a(x) = \lim_{n \rightarrow \infty} \frac{f(2^n x)}{2^n}$ exists for all $x \in E$ and that $a : E \rightarrow E'$ is the unique additive mapping satisfying

$$\|f(x) - a(x)\| \leq \epsilon.$$

Moreover, it was proved that if $f(tx)$ is continuous in t for each fixed $x \in E$, then a is linear. In this case, the Cauchy additive functional equation $f(x+y) = f(x) + f(y)$ is said to satisfy the Hyers-Ulam stability.

In 1978, Th.M. Rassias [35] provided a generalized version of the theorem of Hyers which permitted the Cauchy difference to become unbounded. He proved the following theorem.

Theorem 1. [Th.M. Rassias] *If a function $f : E \rightarrow E'$ between Banach spaces satisfies the inequality*

$$\|f(x+y) - f(x) - f(y)\| \leq \theta (\|x\|^p + \|y\|^p) \quad (1)$$

for some $\theta \geq 0$, $0 \leq p < 1$ and for all $x, y \in E$, then there exists a unique additive function $a : E \rightarrow E'$ such that

$$\|f(x) - a(x)\| \leq \frac{2\theta}{2-2^p} \|x\|^p \quad (2)$$

for all $x \in E$. Moreover, if $f(tx)$ is continuous in t for each fixed $x \in E$, then a is linear.