

Claúdia Torres, Edgar Leitão¹, Joana Leal¹, Sara Camelo¹, João Paulo Sousa^{1,2,3} and Ângela Maria Pereira^{1,4,5}

¹Physiotherapy Department, Escola Superior de Saúde Egas Moniz; ² Departamento de Desporto e Saúde, Escola de Ciência e Tecnologia, Universidade de Évora; ³Comprehensive Health Research Center; ⁴Centro de investigação interdisciplinar Egas Moniz; ⁵Hospital Garcia de Orta, Almada, Portugal

E-mail contact: amcfap@gmail.com

Introduction:

The shoulder of handball players suffers from the ongoing repetition of movement that may lead to the development of joint instability and modification of proprioceptive conditions(1). Articular components of the shoulder joint are considered to have extreme importance on static and dynamic stabilization and quality of proprioceptive information so athlete's performance can be compromised whenever they are affected (2).

Objective:

Therefore, the objective of our study was to verify which factors may contribute to the development of shoulder instability in handball players. The focus was placed on shoulder strength parameters (SSP) and joint position sense (JPS) evaluation of internal and external rotation.

Materials and Methods:

Our study followed a cross-sectional design. The sample was composed by eleven handball players of both genders (4 females and 7 males), with an average age of 22 years, average height of 177 cm, and average weight of 76 kg . Data was collected using a isokinetic dynamometer (Biodex System 3) available at the Gymnastic Pavilion of the University of Évora. Isokinetic dynamometer was used to assess both SSP and JPS.

The evaluation of SSP was implemented prior to JPS at 60° (3 reps) and 180° (20 reps) as described on the protocol for Biodex shoulder assessment of internal and external rotation. Assessment position started with glenohumeral joint at 90° of abduction (Figure 1 and 2).

JPS was evaluated using active positioning and passive determined position at three given external shoulder rotation amplitudes (75°, 35° and 20°). For both active and passive JPS an upper limb inflatable sleeve was used. Participants put a blindfold when determining passive JPS (Figure 3 and 4).



Fig 1. SSP Evaluation: Shoulder at 90° abduction with 90° of external rotation




Fig 2. SSP Evaluation: Shoulder at 90° abduction with 0° of external rotation




Fig 3. JPS Evaluation: Shoulder at 90° abduction with 35° of external rotation



Fig 3. JPS Evaluation: Shoulder at 90° abduction with 75° of external rotation

Results:

For the right upper limb, on passive movement 6.7 ± 5.8 (75°), 1.7 ± 2.4 (35°) and 0 ± 2.3 (20°). In active movement 1.3 ± 3.1 (75°), 1.0 ± 3.6 (35°) and 1.2 ± 2.7 (20°) were observed. In the left upper limb, we observed the following results, in passive movement: 4.7 ± 3.7 (75°), 1.3 ± 4.1 (35°) and 0.7 ± 3.5 (20°). Finally, in active movement we obtained the following results: 0.5 ± 3.4 (75°), 0.7 ± 2.0 (35°) and 2.7 ± 2.4 (20°). In the SSP the peak torque (PK), at 60° right upper limb, external rotation 41.4 ± 12.2 and internal rotation 40.6 ± 17.5 ; at 60° left upper limb, external rotation 34.3 ± 10.7 and internal rotation 42.0 ± 12.8 . PK at 180° right upper limb, external rotation 37.2 ± 12.2 and internal rotation 37.9 ± 15.0 ; at 180° left upper limb, external rotation 29.8 ± 10.8 and internal rotation 37.2 ± 11.6 . Range of Movement (ROM) at 60° (89.6 ± 3.2) and at 180° (88.8 ± 0.6).




Figure 1. Peak Torque




Figure 2. Range of Movement.




Figure 3. Joint Position Sense (JPS)

Discussion and Conclusions:

It was possible to conclude after analyzing the collected data, that dominant upper limb (JPS) obtained better results, compared to the non-dominant. So, we can affirm that in active movement athletes had a better perception of joint position, compared to passive movement. Also we can conclude that all athletes that had previous injuries had a decreased JPS (3).

Dominant limb performed greater ROM at 60° as 180° compared to non-dominant limb. as we can see in the figure 2 the accumulation of forces can cause reduction of the internal rotation and increase the movement of external rotation in comparison with uninjured shoulder4.

On average, athletes have a higher peak torque when performing internal than external rotation. It is also verified, based on recent literature, that the dominant member presents a higher peak torque in the internal rotation than external rotation at 60°, but at 180° the opposite occurs. In the non-dominant limb there is a higher peak torque in the internal rotation than external rotation in both range of motion. (60° and 180°) (5).

References

- (1) Karcher, C., & Buchheit, M. (2014). On-Court Demands of Elite Handball, with Special Reference to Playing Positions. *Sports Medicine*, 44(6), 797-814. <http://dx.doi.org/10.1007/s40279-014-0164-z>
- (2) Hm, L., Ji, L., Ch, C., Cm, T., & Jt, s. (2003). Evaluation of shoulder proprioception following muscle fatigue. *Clin Biomech*, 18(9).
- (3) Higson, E., Herrington, L., Butler, C., & Horsley, I. (2018). The short-term effect of swimming training load on shoulder rotational range of motion, shoulder joint position sense and pectoralis minor length. *Shoulder & Elbow*, 10(4), 285-291. doi: 10.1177/1758573218773539
- (4) Fieseler, G., Jungermann, P., Koke, A., Irlenbusch, L., Delank, K., & Schwesig, R. (2015). Glenohumeral range of motion (ROM) and isometric strength of professional team handball athletes, part III: changes over the playing season. *Archives Of Orthopaedic And Trauma Surgery*, 135(12), 1691-1700. doi: 10.1007/s00402-015-2308-5
- (5) Dover, G., & Powers, M. E. (2003). During Internal and External Rotation of the Shoulder. *Journal of Athletic Training*, 38(4), 304-310.