



Nutritional care in older adults: are we doing everything? An expert opinion review

Elisabet Sanchez-Garcia, Alfonso J. Cruz-Jentoft, Paula Ravasco, Merja Suominen & Prof Kaisu Pitkälä

To cite this article: Elisabet Sanchez-Garcia, Alfonso J. Cruz-Jentoft, Paula Ravasco, Merja Suominen & Prof Kaisu Pitkälä (2024) Nutritional care in older adults: are we doing everything? An expert opinion review, Current Medical Research and Opinion, 40:9, 1563-1576, DOI: [10.1080/03007995.2024.2380007](https://doi.org/10.1080/03007995.2024.2380007)

To link to this article: <https://doi.org/10.1080/03007995.2024.2380007>



© 2024 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group



Published online: 01 Aug 2024.



Submit your article to this journal [↗](#)



Article views: 5642



View related articles [↗](#)



View Crossmark data [↗](#)



Citing articles: 8 View citing articles [↗](#)

REVIEW ARTICLE

 OPEN ACCESS Check for updates

Nutritional care in older adults: are we doing everything? An expert opinion review

Elisabet Sanchez-Garcia^{a,b} , Alfonso J. Cruz-Jentoft^b , Paula Ravasco^c , Merja Suominen^d  and Prof Kaisu Pitkälä^d 

^aConsultant in Geriatric Medicine, Mater Private Network, Cork, Ireland; ^bServicio de Geriátria, Hospital Universitario Ramón y Cajal (IRYCIS), Madrid, Spain; ^cUniversidade Católica Portuguesa, Faculty of Medicine and Centre for Interdisciplinary Research in Health, Centre for Interdisciplinary Research Egas Moniz (CiiEM), Lisbon, Portugal; ^dDepartment of General Practice and Primary Health Care, University of Helsinki, and Helsinki University Hospital, Helsinki, Finland

ABSTRACT

Malnutrition is a prevalent, yet often underdiagnosed and undertreated, condition in older adults. It is characterized by weight loss and/or reduced muscle mass due to diminished caloric intake, inflammation, and/or disease burden. In return, malnutrition can lead to diminished skeletal muscle functionality and disability, among others. Malnutrition plays a crucial role in the pathogenesis of two prevalent geriatric syndromes, namely sarcopenia and frailty. The complex interplay between malnutrition, sarcopenia, and frailty significantly impacts the older population, leading to increased morbidity, mortality, hospitalization rates, quality-of-life, and healthcare costs. Given the prognostic significance of malnutrition in geriatric care, recent guidelines emphasized the role of nutritional support in vulnerable populations. A group of vulnerable populations to malnutrition, sarcopenia, and frailty are older patients with hip fractures, cancer patients, and those with sarcopenic dysphagia. This article highlights the importance of individualized nutritional assessment and treatment in the management of vulnerable populations such as older patients with hip fractures, cancer, and those suffering from sarcopenic dysphagia. It presents practical protocols and guidelines that can be instrumental in enhancing the nutritional care of these groups, thereby improving their overall health outcomes.

ARTICLE HISTORY

Received 28 May 2024
Revised 8 July 2024
Accepted 10 July 2024

KEYWORDS

Ageing; malnutrition; frailty; Sarcopenia; geriatrics; nutritional interventions

Introduction

Despite the considerable progress in geriatric care, malnutrition remains challenging for older adults. Previous reports suggested that 3.2% and 23.4% of community-dwelling older adults in Europe are malnourished or at risk of malnutrition, respectively¹. Additionally, the prevalence of malnutrition increases progressively with declined functional dependency, affecting nearly 40% of hospitalized patients and 50% of those in rehabilitation².

The development of malnutrition in older adults is multifaceted. Ageing is inherently associated with physiological, psychological, and social changes, directly affecting nutritional status and dietary needs³. Age-related physiological changes, such as altered digestive functions, changes in taste and smell, decreased physical activity, depletion of lean body mass, poor muscle health, and declining mental status, significantly impact dietary intake and lead to progressive nutritional deterioration^{4,5}. Furthermore, chronic inflammatory diseases and swallowing disorders, which are more common in older adults, often affect appetite and food intake and induce skeletal muscle catabolism, leading to malnutrition^{3,6}.

From a clinical standpoint, malnutrition in older adults is associated with an increased risk of mortality, prolonged hospitalization, impaired recovery from acute illness or surgery, falls and fractures, impaired immunity, and recurrent infections⁷. The humanistic burden of malnutrition is also substantial in older adults; previous reports showed that malnutrition increased the risk of poor health-related quality-of-life (HRQoL) and impaired individual independence^{8–10}. Additionally, malnutrition generates an economic burden due to increased hospitalizations and healthcare resource utilization (HCRUs)^{11,12}.

According to the Global Leadership Initiative on Malnutrition (GLIM) consensus, reduced muscle mass is one of the phenotypic criteria that defines malnutrition¹³. In return, malnutrition plays a crucial role in the pathogenesis of two distinct geriatric syndromes, namely sarcopenia and frailty. Sarcopenia was formally recognized as a disease in 2016 (ICD-10-CM code: M62.84) and is defined as an age-related loss of muscle mass and strength, affecting primarily functional outcomes^{14,15}. Sarcopenia is closely associated with malnutrition; the synergy between age-related loss of

CONTACT Elisabet Sanchez-Garcia  elisabet_sanchez@hotmail.com  Consultant in Geriatric Medicine, Mater Private Network in Cork, Cork, T12 K199, Ireland; Servicio de Geriátria, Hospital Universitario Ramón y Cajal (IRYCIS), Madrid, Spain

© 2024 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group
This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited, and is not altered, transformed, or built upon in any way. The terms on which this article has been published allow the posting of the Accepted Manuscript in a repository by the author(s) or with their consent.
www.cmrojournal.com

muscle mass and function (a hallmark of sarcopenia) and protein deficiency creates a vicious cycle of malnutrition-sarcopenia syndrome, in which reduced protein intake exacerbates muscle loss, further diminishing the capacity to consume and utilize nutrients effectively^{16,17}. Previous reports showed that nearly one-third of older adults at nutritional risk were sarcopenic¹⁸.

Likewise, frailty, a condition characterized by decreased physiological reserve and increased vulnerability to stressors, is both a consequence and a contributor to the cycle of malnutrition and sarcopenia. The interplay of malnutrition and sarcopenia significantly escalates the risk and severity of frailty¹⁹. The risk of malnutrition was reported in up to 64% of frail older adults and 66% of sarcopenic patients in the community setting^{20,21}. Malnutrition, sarcopenia, and frailty in older adults significantly increase the risk of serious consequences, including disability, prolonged hospitalization, falls and fractures, impaired HRQoL, and mortality^{22–27}. On the other hand, low physical activity plays a key role in the development of sarcopenia and frailty²⁸, and it is identified as a relevant modifiable risk factor²⁹. This is particularly relevant given the high burden of low physical activity globally and its association with mortality and disability-adjusted life years³⁰.

The interconnected nature of malnutrition, sarcopenia, and frailty, as well as their deleterious consequences, highlights the need for a multifaceted approach to the nutritional management of older adults, particularly those with chronic diseases or acute events exacerbating the risk of nutritional deficiencies. Various clinical guidelines recommend personalized nutritional assessment and treatment, including supplementation when necessary, for older adults experiencing malnutrition or at risk of it. These guidelines emphasize the significance of an integrated and tailored nutritional approach across different clinical contexts, underscoring its role in enhancing health outcomes^{31,32}. Still, malnutrition remains under-recognized and unaddressed in clinical practice. Furthermore, the availability of multidisciplinary teams, including dietitians, physiotherapists, and speech therapists, varies in different healthcare settings, presenting additional challenges. These disparities necessitate the customization of nutritional care plans tailored to the resources and circumstances of each setting. There is also a need to improve nutritional education and awareness among healthcare providers. Overcoming these challenges is imperative for clinicians committed to effectively addressing nutritional care. In this article, we review three clinical scenarios – hip fracture, cancer, and oropharyngeal dysphagia (OD) – which pose significant nutritional risks in older adults, and provide specific practical protocols to address these concerns.

Review development

This review article is based on the scientific contents of a satellite symposium conducted at the 19th Congress of the European Union Geriatric Medicine Society (EuGMS), Helsinki, Finland. The symposium was held on September 21, 2023, and aimed to discuss recent updates in nutritional care for

older adults at risk due to acute events or chronic diseases. During this symposium, the consequences of malnutrition and nutritional management protocols for the following vulnerable populations were discussed: older adults with hip fractures, cancer patients, and those with sarcopenic dysphagia.

An online bibliographic search was conducted in Medline *via* PubMed from its inception to November 2023 to support the development of this article. A MeSH-based search strategy was employed as follows: (“Geriatrics” [MeSH Terms]) AND (“Malnutrition” [MeSH Terms]) OR (“Sarcopenia” [MeSH Terms]) OR (frail older) OR “Geriatric Nutrition”) AND (“Chronic Disease” [MeSH Terms]) OR “hip fracture” OR “Cancer” OR “Sarcopenic dysphagia”). Additionally, supplementary keywords were included alongside MeSH terms, such as “elderly nutritional care”, “aging and dietary needs”, and “geriatric dietary interventions”. Clinical trials, real-world studies, reviews, and guidelines documents were reviewed, and relevant data were extracted.

Changing the patterns of nutritional care in the orthogeriatric wards

Hip fractures represent a major public health burden, with a global incidence of ten million cases per year. As the population ages, the incidence of hip fractures is expected to increase significantly, approaching 21 million cases by 2050^{33,34}. The current body of evidence shows a rising trend of hip fractures with advancing age; recent reports showed that the incidence of hip fractures among older adults ranged from 95–316 per 100,000 population, which is expected to double by 2050 as the population ages³⁵. While the incidence of hip fractures is constantly higher in adults older than 65 years compared to younger age groups, there are observed variations in the incidence of hip fractures among older adults based on sex (higher incidence in females due to osteoporosis³⁶) geographical location (increasing trends in Asia³⁷), comorbidities, and lifestyle³⁵. Several factors contribute to the high risk of falls and hip fractures with aging, including loss of bone mineral density, osteoporosis and osteopenia, thyroid disorders, and accidental falls from conditions impairing balance, such as Parkinson’s disease, stroke, and peripheral neuropathy^{38,39}.

Over the years, hip fractures have become a true geriatric syndrome, with significant medical and socioeconomic consequences due to the complexity of care required and the profound impact on mortality, morbidity, HRQoL, and HCRUs⁴⁰. The median age at the onset of hip fractures in developed countries was found to be 83–84 years old^{36,41}, a group that is inherently vulnerable to impaired recovery, complications, and a higher risk of mortality. Hip fractures are notably associated with increased mortality rates in older adults; the 1-year mortality rate following a hip fracture is particularly high, ranging from 14.4–34.8%^{35,42}. Patients with hip fractures have a 5–8-times greater mortality rate than patients without fractures within the first 3 months of the event⁴³.

Alongside the high mortality, hip fractures in older adults are associated with a substantial morbidity burden. Older adults with hip fractures exhibited an increased risk of serious complications, including recurrent infections, prolonged hospitalization, chronic pain, delirium, pressure sores or ulcers, and increased risk of new fractures⁴⁴. Notably, it was found that >50% of older adults who suffer a hip fracture never recover their baseline functional status⁴⁵. This decline in functionality often results in impaired mobility, leading to a loss of independence and the need for long-term care or assistance with daily activities. Previous reports showed that patients with a hip fracture are 4.2-times more likely to have mobility limitations in the community 2 years after fracture⁴⁶, while only 40–70% of patients return to their pre-fracture level of independence⁴⁷. Additionally, patients who experience a hip fracture are 5.6-times more likely to require institutionalization in the following year post-fracture⁴⁸. The overall HRQoL of these patients was also found to be significantly impacted⁴⁹.

The burden of malnutrition is another prevalent concern in older adults with hip fractures (Figure 1). In a recent systematic review of nine studies ($n = 1,665$ participants, mean age 79.9–86.1 years old), the reported prevalence of malnutrition, as diagnosed by the Mini Nutritional Assessment (MNA[®]), among hip fracture patients ranged from 4–39%⁵⁶. Another report showed that nearly 58% of those admitted to a rehabilitation ward were at risk of malnutrition⁵⁷. The relationship between malnutrition and the incidence of falls and hip fractures in older adults can be described as a cause-and-effect cycle that exacerbates both conditions. As previously mentioned, malnutrition contributes to the loss of skeletal muscle mass and functionality, mobility impairment, and accidental falls^{7,50}. On the other hand, patients with hip fractures suffer from mobility

impairment and prolonged hospitalization⁴⁶, which can exacerbate muscle loss and lead to poor nutritional intake⁵¹.

Malnutrition can have a profound negative impact on the outcomes of older adults with hip fractures. The cumulative body of evidence suggests that pre-existing malnutrition is an independent predictor of mortality^{58,59}. In a recent report of 318 hip fracture patients aged >50 years old, malnutrition was significantly associated with shorter survival time, and it independently predicted 6-month mortality⁶⁰. Additionally, among older adults with hip fractures, malnutrition significantly increased the risk of postoperative delirium⁶¹, impaired cognition⁵⁸, worse rehabilitation outcomes, poor functional status, impaired gait at discharge^{57,62,63}, impaired recovery to pre-fracture daily activities and walking ability⁵², and functional dependence⁶⁴.

Given these profound effects, it is not surprising that a growing number of published studies evaluated the impact of nutritional interventions on the outcomes of older adults with hip fractures and increased risk of malnutrition. Initially, a systematic review of 18 trials by the Cochrane Group found no significant impact of oral nutritional supplements (ONS) on mortality; however, the use of ONS was associated with a reduced risk of complications – such as infection and thrombosis – and unfavorable outcomes⁶⁵. Nonetheless, these findings were based on low-quality evidence, and most included studies had a high risk of selection bias⁶⁶. More recently, a systematic review by Takahashi et al. included ten trials that evaluated nutritional supplements in older adults with hip fractures ($n = 1119$ patients). The pooled analysis showed that nutritional supplements were associated with significantly lower risks of mortality (risk ratio [RR] = 0.61) and complications (RR = 0.67), as well as improved skeletal muscle functions⁶⁷. Another recent meta-analysis showed similar




Figure 1. Reported prevalence of risk of malnutrition and malnutrition in older adults with hip fracture, as assessed by MNA^{®50–55}.

findings⁶⁴. A systematic review concluded that nutritional therapy in older adults with hip fractures was cost-effective and improved functional recovery⁵⁹.

These findings reflect the crucial role of addressing malnutrition in the context of hip fractures. The management of hip fractures in geriatric patients is not only a matter of surgical intervention but also involves comprehensive postoperative care (i.e. orthogeriatric care), where nutrition plays a pivotal role. Several orthogeriatric care models, including nutritional intervention, were proposed for managing hip fracture patients, with generally favorable impacts on the mortality and morbidity of the patients^{68–70}. In a recent meta-analysis of 37 studies on patients with hip fractures, orthogeriatric care significantly reduced the length of stay (mean difference [MD] = −1.55 days), in-hospital mortality (risk difference [RD] = 28%), 1-year mortality (RD = 14%), and delirium (RD = 19%). The effect of orthogeriatric care was more prominent in the shared care model⁷¹.

In return, multiple clinical guidelines recommend nutritional therapy as part of multidisciplinary orthogeriatric care for at-risk patients with hip fractures. Integrated into a multi-dimensional, multidisciplinary management plan, the European Society for Clinical Nutrition and Metabolism (ESPEN) recommends postoperative ONS for geriatric patients with hip fractures to reduce the risk of complications³². Similar recommendations were provided by the Japanese Association of Rehabilitation Nutrition⁷², the Italian intersociety consensus⁷³, and the Finnish Current Care Guidelines⁷⁴, among others^{75,76}. However, strategies to address malnutrition and improve clinical outcomes as an integral part of orthogeriatric care of patients with hip fractures remain under-recognized in clinical practice. Addressing the challenge of malnutrition in older adults with hip fractures requires a proactive, multifaceted approach.


In the practice of one of the authors (AJCJ), we follow a nutritional pathway incorporating geriatrician-led

multidisciplinary integrated orthogeriatric care for older adults with hip fractures (Figure 2). This approach prioritizes early nutritional screening and dysphagia assessment on admission. We also recommend specific ONS (with high-quality protein, essential amino acids, calcium, and vitamin D) for 3 months for all patients, alongside assessing and treating the causes of malnutrition. The multidisciplinary team revises the nutritional status at 3 months to optimize the nutritional support for those who still have nutritional problems. Rehabilitation and physical therapy play a key role in functional and nutritional recovery. All nutritional plans should be part of a multifaceted intervention that includes physical therapy^{72,77}.

In this model, nutritional treatment plans are followed up at primary care to ensure their effectiveness and continuity, with primary care playing a crucial role in maintaining and adjusting the nutritional interventions according to the patient's evolving needs, ensuring a seamless transition from hospital to home care. This model may be helpful to others in changing the patterns of nutritional care in hip fracture care in a quest to improve patient outcomes.

Malnutrition, sarcopenia, and frailty: the cornerstone in the management of the oncogeriatric patient

Cancer is predominantly a disease of older individuals, with a significant proportion of cancer diagnoses occurring in those aged ≥65 years. Epidemiological data reveal that the incidence and mortality rates of cancer increase dramatically with age; nearly 60% of all new cancer cases and 70% of cancer-related deaths occur in older adults⁷⁸. Additionally, recent estimates projected a notable increase in cancer incidences among older adults parallel to the increase in the global older adult population. It was estimated that cancer incidence among older adults will increase from 6.7 million



*Specific ONS: High-quality protein, essential amino acids, calcium and vitamin D

Figure 2. A proposed nutritional pathway for hip fractures in a geriatrician-led integrated orthogeriatric unit. Abbreviations: MNA®-SF, Mini Nutritional Assessment–short form; MN, malnutrition; ONS, oral nutritional supplement.

cases in 2012 to 14 million cases by 2035, with the highest predicted increase in the Middle East and Northern Africa (+157%) and in China (+155%)⁷⁹. Other estimates revealed that, among adults aged ≥ 80 years, cancer incidence will increase from 2.3 million cases in 2018 to 6.9 million cases in 2050 (20.5% of all cancer cases)⁸⁰. The high burden of cancer incidence in older adults is attributed to various factors, including age-related changes in the immune system and the increased prevalence of other risk factors, such as obesity and smoking during their life course, in older individuals⁸¹.

Older adults with cancer often present with unique challenges. Previous reports demonstrated that 40% of older adults with cancer present with at least one comorbidity, particularly cardiovascular disease or diabetes⁸². Older adults usually present with more advanced disease stages, a lower probability of resectable disease, and a lower chance of receiving chemotherapy/radiotherapy⁸³. Compared to younger populations, older adults have a higher risk of cancer mortality and shorter survival time⁸⁴.

Growing evidence suggests an alarming prevalence of malnutrition in patients with cancer, particularly in older adults. Previous reports suggested that nearly 20–70% of patients with cancer are malnourished or at risk of developing malnutrition, with a higher incidence in patients with advanced stages and/or systemic inflammation⁸⁵. In older adults (mean age = 65 years) with cancer, the prevalence of malnutrition was found to be 34%; of these, 89% had moderate-to-severe malnutrition that requires urgent nutritional support⁸⁶. Other reports indicated that the prevalence of moderate-to-severe malnutrition was as high as 58% in patients with cancer aged ≥ 65 years⁸⁷. Alongside age-related changes, the development of malnutrition in patients with cancer is multifaceted and attributed to several cancer-related, treatment-related, and patient-specific factors⁸⁵. Cancer patients may have an impaired desire to eat or eating difficulties due to the local tumor effects or treatment toxicities. In addition, cancer-mediated metabolic alterations can disrupt normal nutrient metabolism, impacting the digestion and absorption of nutrients^{86,88}.

Malnutrition can have devastating consequences in older adults with cancer. Previous reports showed that malnutrition was significantly associated with poor survival outcomes. Malnourished patients with head and neck tumors (median age = 61 years) were found to have significantly shorter overall survival (OS) and disease-free survival (DFS) than well-nourished patients; in the adjusted Cox regression model, malnutrition was found to be an independent predictor of OS⁸⁹. Another multivariate analysis showed that weight loss $\geq 5\%$ was significantly associated with shorter progression-free survival (PFS)⁹⁰. Malnourished patients with cancer often experience higher toxicity rates and lower tolerance to chemotherapy, necessitating dose reductions or interruptions^{91,92}. Furthermore, malnourished patients were found to have a higher risk of infections, postoperative complications, prolonged hospitalization, and unplanned hospital visits^{92–94}.

Perhaps one of the major consequences of malnutrition in older adults with cancer is the increased risk of muscle-related conditions, i.e. sarcopenia and cachexia, due to the

potential synergistic adverse effects. In patients with head and neck cancer, it was found that 28.6%, 24.2%, and 43.8% of the patients with sarcopenia, frailty, and cachexia were malnourished, respectively⁹⁵. Malnutrition has a complex and often cyclic interplay with sarcopenia, frailty, and cachexia, attributed to the interconnected etiological factors^{96,97}, as inadequate nutritional intake can exacerbate the decline in muscle mass and function, worsening frailty and sarcopenia; both conditions are already common in cancer patients due to multimorbidity, chronic systemic inflammation, oxidative stress, age-related metabolic changes, and limited mobility⁹⁸. Likewise, poor nutrient intake and nutritional deterioration can worsen cachexia, a prevalent, life-threatening condition in cancer patients characterized by skeletal muscle and/or adipose tissue wasting due to a negative protein and energy balance and metabolic derangements⁹⁹. Malnutrition thus becomes more severe in patients with cachexia¹⁰⁰.

Sarcopenia and frailty in patients with cancer are associated with increased treatment toxicity, postoperative complications, and worse survival outcomes^{101,102}. These adverse effects can be further exacerbated by malnutrition; a previous report demonstrated that the concurrent presence of malnutrition risk and sarcopenia was associated with a 2-fold increase in mortality among hospitalized patients¹⁰³. Additionally, cachexia was found to be associated with worse OS¹⁰⁰. While it is plausible to predict a synergistic adverse effect when malnutrition co-occurs with cachexia, this cumulative effect remains unclear and needs further research.

Despite the profound impact of malnutrition on older adults with cancer, it remains largely under-diagnosed and untreated in clinical practice. Nutritional care should be an integral part of cancer care, and there is a need for a tailored multimodal care intervention in managing oncogeriatric patients with poor nutritional status, particularly considering the interplay between malnutrition, frailty, sarcopenia, and cachexia¹⁰⁴. In older adults with cancer, malnutrition, sarcopenia, frailty, and cachexia should not be seen as isolated conditions but as disorders that can co-exist or occur sequentially (31% of the patients had at least two concurrent conditions⁹⁵) in which malnutrition represents a common factor^{13,96}.

Thus, early nutritional support can potentially reduce the risk of muscle-related conditions (i.e. sarcopenia, cachexia, and frailty), treatment-related adverse events, treatment interruptions or discontinuation, and poor survival^{105–108}. Nutritional screening and intervention should start during the initial work-up, and the objectives of nutritional therapy are to recover or maintain adequate nutritional status, correct nutritional deficits, reduce the frequency and duration of treatment interruptions or rehospitalizations, and enhance the overall functional status of the patient¹⁰⁹. The ELCAPA (ELderly CAncer PATients) survey identified malnutrition in older adults with cancer by the presence of one of the following criteria: “at least 10% weight loss in 6 months or 5% in 1 month and/or body mass index less than 21 kg/m² and/or Mini-Nutritional Assessment score less than 17/30 and/or serum albumin level less than 35 g/L”. This profile can help

physicians select patients indicated for urgent nutritional intervention¹¹⁰.

Although the optimal nutritional support for older adults with cancer and malnutrition is still not defined, the guidelines of the European Society of Medical Oncology (ESMO) and European Society for Clinical Nutrition and Metabolism (ESPEN) stress the need for maintaining an adequate energy intake and assuring the correct amount of protein^{104,111}. Due to the expected anabolic resistance in older patients with cancer, a higher range of protein intake (1.2–1.5 g/kg/day) is recommended to promote muscle mass balance and energy (25–30 Kcal/kg/day)¹¹². Nutritional interventions can encompass dietary advice, treatment of underlying causes and symptoms, and offer high energy-dense ONS (≥ 2 kcal/ml) in symptomatic patients to improve compliance^{109,111}. Recently, the American Society for Clinical Oncology (ASCO), in collaboration with the Cancer and Aging Research Group (CARG) and the International Society of Geriatric Oncology (SIOG), developed a practical guideline with a multidimensional approach to provide care for all older patients with cancer, incorporating nutritional care¹¹³. In addition to nutritional care, rehabilitation and physical activity play important roles in the multidimensional approach¹¹¹. The current evidence shows that the combination of nutritional interventions and physical activity significantly improves functional status and HRQoL and potentially mitigates the risk of treatment-related adverse events^{72,114,115}.

A group of experts in nutrition, in a collaborative project between Catolica Medical School, Portuguese Institute of Oncology, and CUF Hospital, developed a practical protocol based on clinical evidence and clinical practice that allows healthcare professionals caring for patients with cancer to identify patients with or at risk of malnutrition and/or muscle depletion for subsequent evaluation and follow-up by a multidisciplinary team (Figure 3), currently submitted for publication. Cooperation between specialized professionals in cancer care is essential to assess and improve treatment efficiency and patient care. This cooperation should also involve primary care as an integral part of following up and monitoring the effectiveness of the nutritional plans.

As we look toward the future in the management of cancer patients, particularly in the context of malnutrition, it becomes evident that a one-size-fits-all approach is inadequate. Cancer patients represent a highly heterogeneous population with varying combinations of comorbidities, physiological reserves, nutritional deficiencies, and geriatric syndromes. This diversity necessitates individualized, multifaceted nutritional strategies tailored to each patient's unique needs.

Challenges in the care of sarcopenic dysphagia in older adults

OD is a highly prevalent condition in older adults, particularly among those aged ≥ 75 years. Previous reports demonstrated that nearly 46% of the geriatric population suffers from OD¹¹⁶, with higher prevalences in institutionalized individuals (40–60%^{117,118}), people in geriatric care units

(44%¹¹⁹), hospitalized patients aged ≥ 80 years old (82.4%¹²⁰), older adults with hip fractures (55%¹²¹), and patients with cancer (54%¹²²). In frail patients, the prevalence of OD was reported to be 47.4%¹²³. In fact, aging is characterized by anatomical and functional changes in the swallowing mechanism, referred to as presbyphagia, increasing the risk of OD¹²⁴. Patients with OD are at increased risk of recurrent respiratory infection and aspiration pneumonia (odds ratio [OR] = 2.07)^{116,120}; previous reports demonstrated that older adults aged ≥ 75 years old accounted for 76.0% of aspiration pneumonia-related deaths¹²⁵. Additionally, OD was found to significantly increase the risk of malnutrition (OR = 2.21), delirium (OR = 2.23), and mortality (OR = 2.73)¹¹⁶.

Despite its high prevalence and severe consequences, OD is poorly recognized in clinical settings and is usually diagnosed at a late stage when related complications happen¹²⁶.

OD is acknowledged as a geriatric syndrome, demanding a comprehensive and interdisciplinary evaluation to diagnose and treat the causes of the condition^{127,128}. OD in older adults can result from age-related neurodegenerative diseases; the most common conditions leading to OD include stroke (with reported prevalences of 51–55% with clinical testing and 64–78% with instrumental tools¹²⁹) or progressive neurologic disease (e.g. dementia, amyotrophic lateral sclerosis, Parkinson's disease)¹³⁰. Recent research has identified sarcopenia as one of the potential causes of oropharyngeal dysphagia in older adults, leading to the concept of sarcopenic dysphagia.

Sarcopenic dysphagia refers to swallowing difficulties due to sarcopenia of the swallowing and generalized skeletal muscles. A recent position statement proposed a diagnostic algorithm for sarcopenic dysphagia in older adults¹³¹. The Japanese Society of Dysphagia Rehabilitation diagnostic criteria confirm sarcopenic dysphagia if both OD and sarcopenia are present after excluding other causes of OD. The diagnostic workup for sarcopenic dysphagia starts with the assessment of physical function and total muscle mass to diagnose sarcopenia (Figure 4)¹³². Once sarcopenia is confirmed, swallow function should be assessed using any clinical or instrumental swallowing examinations. Muscle strength should be assessed by examining tongue pressure^{133,134}, considering a tongue pressure of ≤ 20 kPa to confirm a probable diagnosis^{133,134}. The oropharyngeal muscle mass can be measured through various imaging techniques such as ultrasound (US) and computed tomography (CT). These methods provide a visual representation of the muscles, allowing for an assessment of their size and condition^{135,136}. Although some studies suggest using these imaging tests to examine swallowing muscles, there is currently no established cut-off point to determine the loss of swallowing muscle mass. Consequently, evaluating the decrease in swallowing muscle mass using these imaging tests and making a definitive diagnosis for sarcopenic dysphagia, a condition characterized by muscle loss and swallowing difficulties, remains a challenging task.

A growing number of studies have investigated the prevalence of sarcopenic dysphagia in older adults. A recent report found that the prevalence of sarcopenic dysphagia




Figure 3. Early identification and intervention according to the nutritional risk in patients with cancer (submitted).

among hospitalized older adults was 22.8%¹³⁷. Other reports showed that 5.6% of older adults living in the community had sarcopenic dysphagia¹³⁸. Furthermore, nearly 32% of the patients undergoing dysphagia rehabilitation had sarcopenic dysphagia¹³⁹. In recent meta-analyses, the pooled OR for the association between sarcopenia and OD ranged from 3.1 to

4.06 ($p < 0.05$)^{116,140}. It should be noted that most of these studies were limited by a non-instrumental assessment of swallowing¹⁴¹.

Sarcopenic dysphagia may also be an important area due to the prognostic significance of sarcopenia in frail older adults. Previous reports showed that impaired tongue




Figure 4. A proposed diagnostic algorithm for sarcopenic dysphagia. Adapted from Kakehi et al.¹³². Abbreviations: BIA, bioelectrical impedance analysis; DXA, dual-energy X-ray absorptiometry; GS, gait speed; HS, hand grip strength.

strength was associated with poor recovery, a higher risk of respiratory infection, and worse life expectancy¹⁴². Recent reports also showed that sarcopenic dysphagia was associated with a 1.4-times higher risk of mortality¹⁴³, poor functional capacity, malnutrition, and impaired daily activities¹³². The diagnosis of sarcopenic dysphagia may also be significant as it appears to have prognostic implications in terms of swallowing function recovery¹³⁷. On the other hand, dysphagia results in decreased nutrient intake, leading to malnutrition and exacerbating overall muscle loss and weakness, including the muscles involved in swallowing^{144,145}.

Sarcopenic dysphagia can be unintentionally triggered or worsened by medical interventions. This is known as iatrogenic sarcopenia, which can occur during hospitalization and is a significant concern¹⁴⁶. It can be categorized into three types: activity-related (caused by unnecessary inactivity), nutrition-related (due to inappropriate nutritional care), and disease-related (stemming from iatrogenic diseases or adverse drug events)¹⁴⁶. Thus, clinicians involved in dysphagia care must emphasize preventive strategies as an integral part of their practice.

Early diagnosis and treatment may mitigate the consequences of OD, including sarcopenic dysphagia; treatment plans should be based on a multidisciplinary evaluation and tailored based on individual needs and capabilities, with active patient involvement in the decision-making process. Compensatory strategies, such as postural adjustments, swallowing maneuvers, and dietary modifications, represent a cornerstone in

preventing aspiration pneumonia (see Figure 5)¹⁴⁷. Additionally, rehabilitative techniques and muscle-strengthening methods are employed to enhance swallowing and tongue muscle functions¹⁴⁴. Recent case reports demonstrated that rehabilitative techniques, including resistance training of the swallowing muscles, lingual resistance exercises, and/or head-lift exercises, were associated with significant improvements in swallowing function, daily activities, and oral intake^{148–152}.

These rehabilitative techniques should be combined with nutritional therapy to ensure adequate energy and protein intake (Figure 5)¹³². There is some promising evidence, mostly based on clinical cases, that addresses the impact of nutrition on sarcopenic dysphagia. Previous reports showed that an average daily intake of ≥ 30 kcal/Kg, with a protein intake of ≥ 1.2 g/Kg/day, improved swallowing function and tongue strength in older adults with dysphagia^{153,154}. These findings align with the recommendations for older patients at risk of malnutrition according to European guidelines for older adults³². European guidelines recommend providing at least an energy intake of ≥ 30 –35 kcal/kg/day and a protein intake of ≥ 1.2 g/kg/day¹³¹. Several non-European consensuses also exist for the nutritional management of OD patients and support the same recommendations^{155,156}. According to the ESPEN guidelines in geriatrics and neurology, texture-modified enriched food, such as adding thickened fluids (TF)^{157,158}, should be offered as a compensatory strategy to older adults with signs of OD and malnutrition or risk of malnutrition^{32,159}.




Figure 5. A proposed algorithm for the management of sarcopenic dysphagia. Abbreviations: CGA, comprehensive geriatric assessment; IDDSI, international dysphagia diet standardization initiative; ONS, oral nutritional supplement; RH, rehabilitative techniques.

Thus, adding thickened fluids (TF) is a common practice to increase ONS viscosity and allow more time to control the swallowing process^{157,158}. However, modified diets are associated with lower intake and a higher risk of malnutrition, as non-compliance is common¹⁶⁰. Prethickened ONS can be a useful treatment for malnutrition in people with OD when requirements are not reached despite diet enrichment strategies. Prethickened ONS might be preferred as adding thickeners to calorie and protein supplements can sometimes result in a lumpy, less palatable beverage¹⁶¹. In addition to dietary modifications, comprehensive treatment of sarcopenic dysphagia should also encompass educational efforts, integrating primary care for monitoring nutritional plan effectiveness, maintaining oral hygiene, treating periodontal diseases¹⁶², managing xerosis, and regularly reviewing and adjusting medications¹⁴⁶.

Sarcopenic dysphagia has gained attention over the recent years due to its consequences on the clinical outcomes of older adults. Nutrition and rehabilitation emerge as essential aspects, so further research and clinical trials are essential for advancing our understanding and developing effective management strategies for this condition.

Conclusion

In conclusion, malnutrition presents a challenge in medical care, particularly for older adults with hip fractures, cancer, and sarcopenic dysphagia. In these groups, nutritional care is crucial in mitigating malnutrition risks and consequences, especially while recovering from acute illness. This publication discusses practical approaches for managing malnutrition in these clinical situations, including early screening for malnutrition, individualized nutrition plans, and post-intervention

monitoring and adjustment to optimize nutritional outcomes and assess the need for further plans. Integrating primary care is critical to follow-up on the treatment plan and ensure a seamless transition from hospital to home care. Additionally, rehabilitation should be an integral component of nutritional treatment in oncology, hip fracture, and dysphagia due to its multifaceted benefits on patient outcomes^{72,77,111,114}. This approach ensures that nutritional care is a foundational aspect of comprehensive geriatric care, offering a proactive pathway to enhance patient outcomes. By standardizing the incorporation of nutritional assessments and interventions in clinical settings, healthcare providers can more effectively address the nutritional needs of older adults. Involving and integrating the collaboration of both public and private resources may be necessary.

Nutritional management in geriatrics within developing countries presents unique challenges that require tailored strategies. To ensure adequate nutritional care for elderly patients in developing countries, it is crucial to implement early detection programs for malnutrition and chronic diseases specific to this population and train healthcare personnel in geriatric nutritional management. Developing standardized protocols in hospitals, community settings, and primary care settings, as well as forming multidisciplinary teams for comprehensive care, is essential.

As healthcare professionals, we must advocate for including dietitians and specialized rehabilitation therapists or physiotherapists in our healthcare systems. These specialists are essential for the nutritional treatment of elderly patients, particularly those with cancer, hip fractures, and dysphagia.

Continued research and clinical trials are essential for advancing our understanding and developing effective

management strategies for older adults with malnutrition or at risk of developing malnutrition.

Transparency

Declaration of funding

The author retains the editorial process, including the discussion, at all times. Content Ed Net Switzerland provided editorial and medical writing assistance for the preparation of this manuscript; this assistance was funded by Nestlé Health Science. All medical writing activities followed the 2022 Good Publication Practice (GPP 2022) guidelines update. The views and opinions expressed are those of the authors.

Declaration of financial/other relationships

All the authors received honoraria from Nestlé for chairing and presenting this information in a satellite symposium at the 19th Congress of the EuGMS, Helsinki, Finland. ACJ received payment or honoraria for education courses from Abbott Nutrition and Danone-Nutricia. ACJ received payment or honoraria for lectures from Nestlé Health Science and Fresenius-Kabi. PR received honoraria for lectures from Nestlé Health Science and Danone-Nutricia. ES received consulting fees from Nestlé Health Science, Danone-Nutricia, Abbot, and Fresenius-Kabi. ES payment or honoraria for lectures or presentations from Nestlé Health Science, Danone-Nutricia, Abbot, and Fresenius-Kabi.

A reviewer on this manuscript disclosed that they are employed by Danone (Nutricia) which manufactures products intended for disease-related malnutrition including those referred to as Oral Nutritional Supplements (ONS). Peer reviewers on this manuscript have no relevant financial or other relationships to disclose.

Author contributions

All authors contributed to the conceptualization, writing, and review & editing of the present opinion review. All authors have approved the manuscript before submission.

Acknowledgements

Medical writing and editorial support in preparing this paper were provided by Content Ed Net Switzerland.

ORCID

Elisabet Sanchez-Garcia  <http://orcid.org/0000-0003-0320-5207>
 Alfonso J. Cruz-Jentoft  <http://orcid.org/0000-0001-7628-4861>
 Paula Ravasco  <http://orcid.org/0000-0002-6056-8269>
 Merja Suominen  <http://orcid.org/0009-0008-1431-8542>
 Prof Kaisu Pitkälä  <http://orcid.org/0000-0001-9659-6985>

References

- Cereda E, Pedrolli C, Klersy C, et al. Nutritional status in older persons according to healthcare setting: a systematic review and meta-analysis of prevalence data using MNA[®]. *Clin Nutr*. 2016;35(6):1282–1290. doi: [10.1016/j.clnu.2016.03.008](https://doi.org/10.1016/j.clnu.2016.03.008).
- Kaiser MJ, Bauer JM, Rämisch C, et al. Frequency of malnutrition in older adults: a multinational perspective using the mini nutritional assessment. *J Am Geriatr Soc*. 2010;58(9):1734–1738. doi: [10.1111/j.1532-5415.2010.03016.x](https://doi.org/10.1111/j.1532-5415.2010.03016.x).
- Dent E, Wright ORL, Woo J, et al. Malnutrition in older adults. *Lancet*. 2023;401(10380):951–966. doi: [10.1016/S0140-6736\(22\)02612-5](https://doi.org/10.1016/S0140-6736(22)02612-5).
- O'Keeffe M, Kelly M, O'Herlihy E, et al. Potentially modifiable determinants of malnutrition in older adults: a systematic review. *Clin Nutr*. 2019;38(6):2477–2498. doi: [10.1016/j.clnu.2018.12.007](https://doi.org/10.1016/j.clnu.2018.12.007).
- Streicher M, van Zwienen-Pot J, Bardon L, et al. Determinants of incident malnutrition in community-dwelling older adults: a MaNuEL multicohort meta-analysis. *J Am Geriatr Soc*. 2018;66(12):2335–2343. doi: [10.1111/jgs.15553](https://doi.org/10.1111/jgs.15553).
- Dent E, Hoogendijk EO, Wright ORL. New insights into the anorexia of ageing: from prevention to treatment. *Curr Opin Clin Nutr Metab Care*. 2019;22(1):44–51. doi: [10.1097/MCO.0000000000000525](https://doi.org/10.1097/MCO.0000000000000525).
- Norman K, Haß U, Pirlich M. Malnutrition in older adults-recent advances and remaining challenges. *Nutrients*. 2021;13(8):2764. doi: [10.3390/nu13082764](https://doi.org/10.3390/nu13082764).
- Rasheed S, Woods RT. An investigation into the association between nutritional status and quality of life in older people admitted to hospital. *J Hum Nutr Diet*. 2014;27(2):142–151. doi: [10.1111/jhn.12072](https://doi.org/10.1111/jhn.12072).
- Kvamme JM, Olsen JA, Florholmen J, et al. Risk of malnutrition and health-related quality of life in community-living elderly men and women: the Tromsø study. *Qual Life Res*. 2011;20(4):575–582. doi: [10.1007/s11136-010-9788-0](https://doi.org/10.1007/s11136-010-9788-0).
- Rasheed S, Woods RT. Malnutrition and quality of life in older people: a systematic review and meta-analysis. *Ageing Res Rev*. 2013;12(2):561–566. doi: [10.1016/j.arr.2012.11.003](https://doi.org/10.1016/j.arr.2012.11.003).
- Khalatbari-Soltani S, Marques-Vidal P. The economic cost of hospital malnutrition in Europe; a narrative review. *Clin Nutr ESPEN*. 2015;10(3):e89–e94. doi: [10.1016/j.clnesp.2015.04.003](https://doi.org/10.1016/j.clnesp.2015.04.003).
- Deutz NEP, Ashurst I, Ballesteros MD, et al. The underappreciated role of low muscle mass in the management of malnutrition. *J Am Med Dir Assoc*. 2019;20(1):22–27. doi: [10.1016/j.jamda.2018.11.021](https://doi.org/10.1016/j.jamda.2018.11.021).
- Cederholm T, Jensen GL, Correia MITD, et al. GLIM criteria for the diagnosis of malnutrition – a consensus report from the global clinical nutrition community. *Clin Nutr*. 2019;38(1):1–9. doi: [10.1016/j.clnu.2018.08.002](https://doi.org/10.1016/j.clnu.2018.08.002).
- Cruz-Jentoft AJ, Gonzalez MC, Prado CM. Sarcopenia ≠ low muscle mass. *Eur Geriatr Med*. 2023;14(2):225–228. doi: [10.1007/s41999-023-00760-7](https://doi.org/10.1007/s41999-023-00760-7).
- Anker SD, Morley JE, von Haehling S. Welcome to the ICD-10 code for sarcopenia. *J Cachexia Sarcopenia Muscle*. 2016;7(5):512–514. doi: [10.1002/jcsm.12147](https://doi.org/10.1002/jcsm.12147).
- Cruz-Jentoft AJ, Kiesswetter E, Drey M, et al. Nutrition, frailty, and sarcopenia. *Aging Clin Exp Res*. 2017;29(1):43–48. doi: [10.1007/s40520-016-0709-0](https://doi.org/10.1007/s40520-016-0709-0).
- Vandewoude MFJ, Alish CJ, Sauer AC, et al. Malnutrition-sarcopenia syndrome: Is this the future of nutrition screening and assessment for older adults? *J Aging Res*. 2012;2012:651570–651578. doi: [10.1155/2012/651570](https://doi.org/10.1155/2012/651570).
- Tan VMH, Pang BWJ, Lau LK, et al. Malnutrition and sarcopenia in community-dwelling adults in Singapore: Yishun Health Study. *J Nutr Health Aging*. 2021;25(3):374–381. doi: [10.1007/s12603-020-1542-x](https://doi.org/10.1007/s12603-020-1542-x).
- AlMohaisen N, Gittins M, Todd C, et al. What is the overlap between malnutrition, frailty and sarcopenia in the older population? Study protocol for cross-sectional study using UK Biobank. *PLoS One*. 2022;17(12):e0278371. doi: [10.1371/journal.pone.0278371](https://doi.org/10.1371/journal.pone.0278371).
- Boulos C, Salameh P, Barberger-Gateau P. Malnutrition and frailty in community dwelling older adults living in a rural setting. *Clin Nutr*. 2016;35(1):138–143. doi: [10.1016/j.clnu.2015.01.008](https://doi.org/10.1016/j.clnu.2015.01.008).
- Beaudart C, Rizzoli R, Bruyère O, et al. Sarcopenia: burden and challenges for public health. *Arch Public Heal*. 2014;72:45.
- Thompson MQ, Yu S, Tucker GR, et al. Frailty and sarcopenia in combination are more predictive of mortality than either condition alone. *Maturitas*. 2021;144:102–107. doi: [10.1016/j.maturitas.2020.11.009](https://doi.org/10.1016/j.maturitas.2020.11.009).

- [23] Hsu YH, Liang CK, Chou MY, et al. Association of cognitive impairment, depressive symptoms and sarcopenia among healthy older men in the veterans retirement community in southern Taiwan: a cross-sectional study. *Geriatr Gerontol Int*. 2014;14(51):102–108. doi: [10.1111/ggi.12221](https://doi.org/10.1111/ggi.12221).
- [24] Kojima G, Iliffe S, Jivraj S, et al. Association between frailty and quality of life among community-dwelling older people: a systematic review and meta-analysis. *J Epidemiol Community Health*. 2016;70(7):716–721. doi: [10.1136/jech-2015-206717](https://doi.org/10.1136/jech-2015-206717).
- [25] Kojima G. Quick and simple FRAIL scale predicts incident Activities of Daily Living (ADL) and Instrumental ADL (IADL) disabilities: a systematic review and meta-analysis. *J Am Med Dir Assoc*. 2018;19(12):1063–1068. doi: [10.1016/j.jamda.2018.07.019](https://doi.org/10.1016/j.jamda.2018.07.019).
- [26] Hirani V, Blyth F, Naganathan V, et al. Sarcopenia is associated with incident disability, institutionalization, and mortality in community-dwelling older men: the concord health and ageing in men project. *J Am Med Dir Assoc*. 2015;16(7):607–613. doi: [10.1016/j.jamda.2015.02.006](https://doi.org/10.1016/j.jamda.2015.02.006).
- [27] Kojima G. Frailty as a predictor of future falls among community-dwelling older people: a systematic review and meta-analysis. *J Am Med Dir Assoc*. 2015;16(12):1027–1033. doi: [10.1016/j.jamda.2015.06.018](https://doi.org/10.1016/j.jamda.2015.06.018).
- [28] da Silva VD, Tribess S, Meneguci J, et al. Association between frailty and the combination of physical activity level and sedentary behavior in older adults. *BMC Public Health*. 2019;19(1):709. <https://bmcpublihealth.biomedcentral.com/articles/10.1186/s12889-019-7062-0>. doi: [10.1186/s12889-019-7062-0](https://doi.org/10.1186/s12889-019-7062-0).
- [29] Sobhani A, Sharifi F, Fadayevatan R, et al. Low physical activity is the strongest factor associated with frailty phenotype and frailty index: data from baseline phase of Birjand Longitudinal Aging Study (BLAS). *BMC Geriatr*. 2022;22(1):498. <https://bmgeriatr.biomedcentral.com/articles/10.1186/s12877-022-03135-y>. doi: [10.1186/s12877-022-03135-y](https://doi.org/10.1186/s12877-022-03135-y).
- [30] Xu YY, Xie J, Yin H, et al. The Global Burden of Disease attributable to low physical activity and its trends from 1990 to 2019: an analysis of the Global Burden of Disease study. *Front Public Heal*. 2022;10:1018866.
- [31] Dent E, Morley JE, Cruz-Jentoft AJ, et al. Physical frailty: ICFSR international clinical practice guidelines for identification and management. *J Nutr Health Aging*. 2019;23(9):771–787. doi: [10.1007/s12603-019-1273-z](https://doi.org/10.1007/s12603-019-1273-z).
- [32] Volkert D, Beck AM, Cederholm T, et al. ESPEN practical guideline: clinical nutrition and hydration in geriatrics. *Clin Nutr*. 2022; 41(4):958–989. doi: [10.1016/j.clnu.2022.01.024](https://doi.org/10.1016/j.clnu.2022.01.024).
- [33] Wu AM, Bisignano C, James SL, et al. Global, regional, and national burden of bone fractures in 204 countries and territories, 1990–2019: a systematic analysis from the Global Burden of Disease Study 2019. *Lancet Heal Longev*. 2021;2(9):e580–e592. doi: [10.1016/S2666-7568\(21\)00172-0](https://doi.org/10.1016/S2666-7568(21)00172-0).
- [34] Gullberg B, Johnell O, Kanis JA. World-wide projections for hip fracture. *Osteoporos Int*. 1997;7(5):407–413. doi: [10.1007/pl00004148](https://doi.org/10.1007/pl00004148).
- [35] Sing CW, Lin TC, Bartholomew S, et al. Global epidemiology of hip fractures: secular trends in incidence rate, post-fracture treatment, and all-cause mortality. *J Bone Miner Res*. 2023;38(8): 1064–1075. doi: [10.1002/jbmr.4821](https://doi.org/10.1002/jbmr.4821).
- [36] Longo UG, Viganò M, de Girolamo L, et al. Epidemiology and management of proximal femoral fractures in Italy between 2001 and 2016 in older adults: analysis of the national discharge registry. *Int J Environ Res Public Health*. 2022;19:16985.
- [37] Ballane G, Cauley JA, Luckey MM, et al. Secular trends in hip fractures worldwide: opposing trends east versus west. *J Bone Miner Res*. 2014;29(8):1745–1755. doi: [10.1002/jbmr.2218](https://doi.org/10.1002/jbmr.2218).
- [38] Papaioannou I, Pantazidou G, Baikousis A, et al. Polypharmacy and falls of elderly with hip fracture: a retrospective study and literature review. *Osteoporos Int*. 2020;31: S265–S266. <https://www.cochranelibrary.com/central/doi/10.1002/central/CN-02261693/full?highlight=Abstract=falls%7Cfall%7Cpolypharmacy%7Cpolypharmac%0Ahttps://www.cochranelibrary.com/central/doi/10.1002/central/CN-02261693/full>.
- [39] Marks R, Algrante JP, MacKenzie CR, et al. Hip fractures among the elderly: causes, consequences and control. *Ageing Res Rev*. 2003;2(1):57–93. doi: [10.1016/S1568-1637\(02\)00045-4](https://doi.org/10.1016/S1568-1637(02)00045-4).
- [40] Deutschbein J, Lindner T, Möckel M, et al. Health-related quality of life and associated factors after hip fracture. Results from a six-month prospective cohort study. *PeerJ*. 2023;11:e14671. doi: [10.7717/peerj.14671](https://doi.org/10.7717/peerj.14671).
- [41] Viganò M, Pennestrì F, Listorti E, et al. Proximal hip fractures in 71,920 elderly patients: incidence, epidemiology, mortality and costs from a retrospective observational study. *BMC Public Health*. 2023;23(1):1963. <https://bmcpublihealth.biomedcentral.com/articles/10.1186/s12889-023-16776-4>. doi: [10.1186/s12889-023-16776-4](https://doi.org/10.1186/s12889-023-16776-4).
- [42] Downey C, Kelly M, Quinlan JF. Changing trends in the mortality rate at 1-year post hip fracture – a systematic review. *World J Orthop*. 2019;10(3):166–175. doi: [10.5312/wjo.v10.i3.166](https://doi.org/10.5312/wjo.v10.i3.166).
- [43] Schnell S, Friedman SM, Mendelson DA, et al. The 1-year mortality of patients treated in a hip fracture program for elders. *Geriatr Orthop Surg Rehabil*. 2010;1(1):6–14. doi: [10.1177/2151458510378105](https://doi.org/10.1177/2151458510378105).
- [44] Carpintero P, Caeiro JR, Carpintero R, et al. Complications of hip fractures: a review. *World J Orthop*. 2014;5(4):402–411. doi: [10.5312/wjo.v5.i4.402](https://doi.org/10.5312/wjo.v5.i4.402).
- [45] Tang VL, Sudore R, Cenzer IS, et al. Rates of recovery to pre-fracture function in older persons with hip fracture: an observational study. *J Gen Intern Med*. 2017;32(2):153–158. doi: [10.1007/s11606-016-3848-2](https://doi.org/10.1007/s11606-016-3848-2).
- [46] Norton R, Butler M, Robinson E, et al. Declines in physical functioning attributable to hip fracture among older people. A follow-up study of case-control participants. *Disabil Rehabil*. 2000; 22(8):345–351. doi: [10.1080/096382800296584](https://doi.org/10.1080/096382800296584).
- [47] Dyer SM, Crotty M, Fairhall N, et al. A critical review of the long-term disability outcomes following hip fracture. *BMC Geriatr*. 2016;16(1):158. doi: [10.1186/s12877-016-0332-0](https://doi.org/10.1186/s12877-016-0332-0).
- [48] Autier P, Haentjens P, Bontin J, et al. Costs induced by hip fractures: a prospective controlled study in Belgium. *Belgian Hip Fracture Study Group*. *Osteoporos Int*. 2000;11(5):373–380. <http://www.ncbi.nlm.nih.gov/pubmed/10912837>. doi: [10.1007/s001980070102](https://doi.org/10.1007/s001980070102).
- [49] Griffin XL, Parsons N, Achten J, et al. Recovery of health-related quality of life in a United Kingdom hip fracture population: the Warwick hip trauma evaluation – a prospective cohort study. *Bone Joint J*. 2015;97-B(3):372–382. doi: [10.1302/0301-620X.97B3.35738](https://doi.org/10.1302/0301-620X.97B3.35738).
- [50] Torres MJ, Féart C, Samieri C, et al. Poor nutritional status is associated with a higher risk of falling and fracture in elderly people living at home in France: the Three-City cohort study. *Osteoporos Int*. 2015;26(8):2157–2164. doi: [10.1007/s00198-015-3121-2](https://doi.org/10.1007/s00198-015-3121-2).
- [51] Groenendijk I, Kramer CS, Den Boeft LM, et al. Hip fracture patients in geriatric rehabilitation show poor nutritional status, dietary intake and muscle health. *Nutrients*. 2020;12(9):2528. doi: [10.3390/nu12092528](https://doi.org/10.3390/nu12092528).
- [52] Li HJ, Cheng HS, Liang J, et al. Functional recovery of older people with hip fracture: Does malnutrition make a difference? *J Adv Nurs*. 2013;69(8):1691–1703. doi: [10.1111/jan.12027](https://doi.org/10.1111/jan.12027).
- [53] Goisser S, Schrader E, Singler K, et al. Malnutrition according to mini nutritional assessment is associated with severe functional impairment in geriatric patients before and up to 6 months after hip fracture. *J Am Med Dir Assoc*. 2015;16(8):661–667. <https://linkinghub.elsevier.com/retrieve/pii/S1525861015001814>. doi: [10.1016/j.jamda.2015.03.002](https://doi.org/10.1016/j.jamda.2015.03.002).
- [54] Drevet S, Bioteau C, Mazière S, et al. Prevalence of protein-energy malnutrition in hospital patients over 75 years of age admitted for hip fracture. *Orthop Traumatol Surg Res*. 2014; 100(6):669–674. doi: [10.1016/j.otsr.2014.05.003](https://doi.org/10.1016/j.otsr.2014.05.003).
- [55] Murphy MC, Brooks CN, New SA, et al. The use of the Mini-Nutritional Assessment (MNA) tool in elderly orthopaedic patients. *Eur J Clin Nutr*. 2000;54(7):555–562. doi: [10.1038/sj.ejcn.1601055](https://doi.org/10.1038/sj.ejcn.1601055).

- [56] Foo MXE, Wong GJY, Lew CCH. A systematic review of the malnutrition prevalence in hospitalized hip fracture patients and its associated outcomes. *JPEN J Parenter Enteral Nutr.* 2021;45(6):1141–1152. doi: [10.1002/jpen.2211](https://doi.org/10.1002/jpen.2211).
- [57] Wong AM, Xu BY, Low LL, et al. Impact of malnutrition in surgically repaired hip fracture patients admitted for rehabilitation in a community hospital: a cohort prospective study. *Clin Nutr ESPEN.* 2021;44:188–193. doi: [10.1016/j.clnesp.2021.06.024](https://doi.org/10.1016/j.clnesp.2021.06.024).
- [58] Zanetti M, Gortan Cappellari G, Ratti C, et al. Poor nutritional status but not cognitive or functional impairment per se independently predict 1 year mortality in elderly patients with hip fracture. *Clin Nutr.* 2019;38(4):1607–1612. doi: [10.1016/j.clnu.2018.08.030](https://doi.org/10.1016/j.clnu.2018.08.030).
- [59] Malafarina V, Reginster JY, Cabrerizo S, et al. Nutritional status and nutritional treatment are related to outcomes and mortality in older adults with hip fracture. *Nutrients.* 2018;10(5):555. doi: [10.3390/nu10050555](https://doi.org/10.3390/nu10050555).
- [60] Franz K, Deutschbein J, Riedlinger D, et al. Malnutrition is associated with six-month mortality in older patients admitted to the emergency department with hip fracture. *Front. Med.* 2023;10:28. doi: [10.3389/fmed.2023.1173528](https://doi.org/10.3389/fmed.2023.1173528).
- [61] Mazzola P, Ward L, Zazzetta S, et al. Association between pre-operative malnutrition and postoperative delirium after hip fracture surgery in older adults. *J Am Geriatr Soc.* 2017;65(6):1222–1228. doi: [10.1111/jgs.14764](https://doi.org/10.1111/jgs.14764).
- [62] Inoue T, Misu S, Tanaka T, et al. Pre-fracture nutritional status is predictive of functional status at discharge during the acute phase with hip fracture patients: a multicenter prospective cohort study. *Clin Nutr.* 2017;36(5):1320–1325. doi: [10.1016/j.clnu.2016.08.021](https://doi.org/10.1016/j.clnu.2016.08.021).
- [63] Kobayashi H, Inoue T, Ogawa M, et al. Malnutrition diagnosed by the Global Leadership Initiative on Malnutrition criteria as a predictor of gait ability in patients with hip fracture. *Injury.* 2022;53(10):3394–3400. doi: [10.1016/j.injury.2022.08.004](https://doi.org/10.1016/j.injury.2022.08.004).
- [64] Chen B, Zhang JH, Duckworth AD, et al. Effect of oral nutritional supplementation on outcomes in older adults with hip fractures and factors influencing compliance: a systematic review and meta-analysis of randomized controlled trials. *Bone Joint J.* 2023;105-B(SUPP_6):3–3. doi: [10.1302/1358-992X.2023.6.002](https://doi.org/10.1302/1358-992X.2023.6.002).
- [65] Avenell A, Smith TO, Curtain JP, et al. Nutritional supplementation for hip fracture aftercare in older people. *Cochrane Database Syst. Rev.* 2016;11(11):CD001880.
- [66] Morrison CA, Morrison MM. For you were hungry and I gave you food: the prevalence and treatment of malnutrition in patients with acute hip fracture. *Nutr Clin Pract.* 2022;37(1):59–67. doi: [10.1002/ncp.10809](https://doi.org/10.1002/ncp.10809).
- [67] Takahashi K, Momosaki R, Yasufuku Y, et al. Nutritional therapy in older patients with hip fractures undergoing rehabilitation: a systematic review and meta-analysis. *J Am Med Dir Assoc.* 2020;21(9):1364–1364.e6. doi: [10.1016/j.jamda.2020.07.005](https://doi.org/10.1016/j.jamda.2020.07.005).
- [68] Vidán M, Serra JA, Moreno C, et al. Efficacy of a comprehensive geriatric intervention in older patients hospitalized for hip fracture: a randomized, controlled trial. *J Am Geriatr Soc.* 2005;53(9):1476–1482. doi: [10.1111/j.1532-5415.2005.53466.x](https://doi.org/10.1111/j.1532-5415.2005.53466.x).
- [69] González-Montalvo JL, Mauleón JL, Gil-Garay E, et al. The orthogeriatric unit for acute patients: a new model of care that improves efficiency in the management of patients with hip fracture. *Hip Int.* 2010;20(2):229–235. doi: [10.1177/112070001002000214](https://doi.org/10.1177/112070001002000214).
- [70] Naglie G, Tansey C, Kirkland JL, et al. Interdisciplinary inpatient care for elderly people with hip fracture: a randomized controlled trial. *C Can Med Assoc J.* 2002;167:25–32.
- [71] Van Heghe A, Mordant G, Dupont J, et al. Effects of orthogeriatric care models on outcomes of hip fracture patients: a systematic review and meta-analysis. *Calcif Tissue Int.* 2022;110(2):162–184. doi: [10.1007/s00223-021-00913-5](https://doi.org/10.1007/s00223-021-00913-5).
- [72] Nishioka S, Aragane H, Suzuki N, et al. Clinical practice guidelines for rehabilitation nutrition in cerebrovascular disease, hip fracture, cancer, and acute illness: 2020 update. *Clin Nutr ESPEN.* 2021;43:90–103. doi: [10.1016/j.clnesp.2021.02.018](https://doi.org/10.1016/j.clnesp.2021.02.018).
- [73] De Vincentis A, Behr AU, Bellelli G, et al. Orthogeriatric co-management for the care of older subjects with hip fracture: recommendations from an Italian intersociety consensus. *Aging Clin Exp Res.* 2021;33(9):2405–2443. doi: [10.1007/s40520-021-01898-9](https://doi.org/10.1007/s40520-021-01898-9).
- [74] Society D of the FMS and the FO. Finnish Current Care Guidelines I Hip Fracture [Internet]. 2011. Available from: <https://www.kaypahoito.fi/hoi50040?tab=suositus>
- [75] Khan AA, AbuAlrob H, Al-Alwani H, et al. Post hip fracture orthogeriatric care – a Canadian position paper addressing challenges in care and strategies to meet quality indicators. *Osteoporos Int.* 2023;34(6):1011–1035. doi: [10.1007/s00198-022-06640-3](https://doi.org/10.1007/s00198-022-06640-3).
- [76] Ranhoff AH, Saltvedt I, Frihagen F, et al. Interdisciplinary care of hip fractures.: orthogeriatric models, alternative models, interdisciplinary teamwork. *Best Pract Res Clin Rheumatol.* 2019;33(2):205–226. doi: [10.1016/j.berh.2019.03.015](https://doi.org/10.1016/j.berh.2019.03.015).
- [77] Wakabayashi H, Sakuma K. Rehabilitation nutrition for sarcopenia with disability: a combination of both rehabilitation and nutrition care management. *J Cachexia Sarcopenia Muscle.* 2014;5(4):269–277. <https://onlinelibrary.wiley.com/doi/10.1007/s13539-014-0162-x>. doi: [10.1007/s13539-014-0162-x](https://doi.org/10.1007/s13539-014-0162-x).
- [78] Estape T. Cancer in the elderly: challenges and barriers. *Asia-Pacific J Oncol Nurs.* 2018;5(1):40–42. doi: [10.4103/apjon.apjon_52_17](https://doi.org/10.4103/apjon.apjon_52_17).
- [79] Pilleron S, Sarfati D, Janssen-Heijnen M, et al. Global cancer incidence in older adults, 2012 and 2035: a population-based study. *Int J Cancer.* 2019;144(1):49–58. doi: [10.1002/ijc.31664](https://doi.org/10.1002/ijc.31664).
- [80] Pilleron S, Soto-Perez-de-Celis E, Vignat J, et al. Estimated global cancer incidence in the oldest adults in 2018 and projections to 2050. *Int J Cancer.* 2021;148(3):601–608. doi: [10.1002/ijc.33232](https://doi.org/10.1002/ijc.33232).
- [81] Barbé-Tuana F, Funchal G, Schmitz CRR, et al. The interplay between immunosenescence and age-related diseases. *Semin Immunopathol.* 2020;42(5):545–557. doi: [10.1007/s00281-020-00806-z](https://doi.org/10.1007/s00281-020-00806-z).
- [82] Edwards BK, Noone A, Mariotto AB, et al. Annual Report to the Nation on the status of cancer, 1975–2010, featuring prevalence of comorbidity and impact on survival among persons with lung, colorectal, breast, or prostate cancer. *Cancer.* 2014;120(9):1290–1314. <https://acsjournals.onlinelibrary.wiley.com/doi/10.1002/cncr.28509>. doi: [10.1002/cncr.28509](https://doi.org/10.1002/cncr.28509).
- [83] Alejandro Rauh-Hain J, Pepin KJ, Meyer LA, et al. Management for elderly women with advanced-stage, high-grade endometrial cancer. *Obstet Gynecol.* 2015;126(6):1198–1206. doi: [10.1097/AOG.0000000000001140](https://doi.org/10.1097/AOG.0000000000001140).
- [84] Pham J, Conron M, Wright G, et al. Excess mortality and undertreatment in elderly lung cancer patients: treatment nihilism in the modern era? *ERJ Open Res.* 2021;7(2):00393–2020. doi: [10.1183/23120541.00393-2020](https://doi.org/10.1183/23120541.00393-2020).
- [85] Arends J. Malnutrition in cancer patients: causes, consequences and treatment options. *Eur J Surg Oncol.* 2023;50(5):107074. <https://linkinghub.elsevier.com/retrieve/pii/S0748798323007126>.
- [86] Trabulo C, Lopes J, da Silva Dias D, et al. Assessment of nutritional status of oncology patients at hospital admission: a Portuguese real-world study. *Front Nutr.* 2022;9:972525. doi: [10.3389/fnut.2022.972525](https://doi.org/10.3389/fnut.2022.972525).
- [87] de Pinho NB, Martucci RB, Rodrigues VD, et al. High prevalence of malnutrition and nutrition impact symptoms in older patients with cancer: results of a Brazilian multicenter study. *Cancer.* 2020;126(1):156–164. doi: [10.1002/cncr.32437](https://doi.org/10.1002/cncr.32437).
- [88] Huhmann MB, Cunningham RS. Importance of nutritional screening in treatment of cancer-related weight loss. *Lancet Oncol.* 2005;6(5):334–343. doi: [10.1016/S1470-2045\(05\)70170-4](https://doi.org/10.1016/S1470-2045(05)70170-4).
- [89] Orell HK, Pohju AK, Osterlund P, et al. GLIM in diagnosing malnutrition and predicting outcome in ambulatory patients with head and neck cancer. *Front Nutr.* 2022;9:1030619. doi: [10.3389/fnut.2022.1030619](https://doi.org/10.3389/fnut.2022.1030619).
- [90] Mitry E, Douillard JY, Van Cutsem E, et al. Predictive factors of survival in patients with advanced colorectal cancer: an individual data analysis of 602 patients included in irinotecan phase III

- trials. *Ann Oncol.* 2004;15(7):1013–1017. doi: [10.1093/annonc/mdh267](https://doi.org/10.1093/annonc/mdh267).
- [91] Ravasco P, Monteiro-Grillo I, Vidal PM, et al. Nutritional deterioration in cancer: the role of disease and diet. *Clin Oncol.* 2003; 15(8):443–450. doi: [10.1016/s0936-6555\(03\)00155-9](https://doi.org/10.1016/s0936-6555(03)00155-9).
- [92] Findlay M, White K, Brown C, et al. Nutritional status and skeletal muscle status in patients with head and neck cancer: impact on outcomes. *J Cachexia Sarcopenia Muscle.* 2021;12(6): 2187–2198. doi: [10.1002/jcsm.12829](https://doi.org/10.1002/jcsm.12829).
- [93] Riad A, Knight SR, Ghosh D, et al. Impact of malnutrition on early outcomes after cancer surgery: an international, multicentre, prospective cohort study. *Lancet Glob Heal.* 2023;11(3): e341–e349. doi: [10.1016/S2214-109X\(22\)00550-2](https://doi.org/10.1016/S2214-109X(22)00550-2).
- [94] Muresan BT, Núñez-Abad M, Artero A, et al. Relation of malnutrition and nosocomial infections in cancer patients in hospital: an observational study. *J Nutr Metab.* 2022;2022:5232480–5232488. doi: [10.1155/2022/5232480](https://doi.org/10.1155/2022/5232480).
- [95] Gingrich A, Volkert D, Kiesswetter E, et al. Prevalence and overlap of sarcopenia, frailty, cachexia and malnutrition in older medical inpatients. *BMC Geriatr.* 2019;19(1):120. doi: [10.1186/s12877-019-1115-1](https://doi.org/10.1186/s12877-019-1115-1).
- [96] Prado CM, Landi F, Chew STH, et al. Advances in muscle health and nutrition: a toolkit for healthcare professionals. *Clin Nutr.* 2022;41(10):2244–2263. doi: [10.1016/j.clnu.2022.07.041](https://doi.org/10.1016/j.clnu.2022.07.041).
- [97] Ryan AM, Power DG, Daly L, et al. Cancer-associated malnutrition, cachexia and sarcopenia: the skeleton in the hospital closet 40 years later. *Proc Nutr Soc.* 2016;75(2):199–211. doi: [10.1017/S002966511500419X](https://doi.org/10.1017/S002966511500419X).
- [98] Stumph J, König F. Sarcopenia and Frailty in Lung Cancer. *Lung Cancer Rehabil.* 2022;2022:139–149.
- [99] Ni J, Zhang L. Cancer cachexia: definition, staging, and emerging treatments. *Cancer Manag Res.* 2020;12:5597–5605. doi: [10.2147/CMAR.S261585](https://doi.org/10.2147/CMAR.S261585).
- [100] Orell-Kotikangas H, Österlund P, Mäkitie O, et al. Cachexia at diagnosis is associated with poor survival in head and neck cancer patients. *Acta Otolaryngol.* 2017;137(7):778–785. doi: [10.1080/00016489.2016.1277263](https://doi.org/10.1080/00016489.2016.1277263).
- [101] da Silva Dias D, Machado M, Trabulo C, et al. Impact of Body Composition on Prognosis and Dose-Limiting Toxicities on Metastatic Colorectal Cancer. *Front Nutr.* 2021;8:671547. doi: [10.3389/fnut.2021.671547](https://doi.org/10.3389/fnut.2021.671547).
- [102] Feliciano EMC, Kroenke CH, Meyerhardt JA, et al. Metabolic dysfunction, obesity, and survival among patients with early-stage colorectal cancer. *J Clin Oncol.* 2016;34(30):3664–3671. doi: [10.1200/JCO.2016.67.4473](https://doi.org/10.1200/JCO.2016.67.4473).
- [103] Hu X, Zhang L, Wang H, et al. Malnutrition-sarcopenia syndrome predicts mortality in hospitalized older patients. *Sci Rep.* 2017; 7(1):3171. doi: [10.1038/s41598-017-03388-3](https://doi.org/10.1038/s41598-017-03388-3).
- [104] Arends J, Strasser F, Gonella S, et al. Cancer cachexia in adult patients: ESMO Clinical Practice Guidelines. *ESMO Open.* 2021; 6(3):100092. doi: [10.1016/j.esmoop.2021.100092](https://doi.org/10.1016/j.esmoop.2021.100092).
- [105] Baldwin C, Spiro A, Ahern R, et al. Oral nutritional interventions in malnourished patients with cancer: a systematic review and meta-analysis. *J Natl Cancer Inst.* 2012;104(5):371–385. doi: [10.1093/jnci/djr556](https://doi.org/10.1093/jnci/djr556).
- [106] Ravasco P, Monteiro-Grillo I, Camilo M. Individualized nutrition intervention is of major benefit to colorectal cancer patients: long-term follow-up of a randomized controlled trial of nutritional therapy. *Am J Clin Nutr.* 2012;96(6):1346–1353. doi: [10.3945/ajcn.111.018838](https://doi.org/10.3945/ajcn.111.018838).
- [107] Ravasco P, Monteiro-Grillo I, Vidal PM, et al. Dietary counseling improves patient outcomes: a prospective, randomized, controlled trial in colorectal cancer patients undergoing radiotherapy. *J Clin Oncol.* 2005;23(7):1431–1438. doi: [10.1200/JCO.2005.02.054](https://doi.org/10.1200/JCO.2005.02.054).
- [108] Kim J-M, Sung M-K. The Efficacy of Oral Nutritional Intervention in Malnourished Cancer Patients: A Systemic Review. *Clin Nutr Res.* 2016;5(4):219–236. doi: [10.7762/cnr.2016.5.4.219](https://doi.org/10.7762/cnr.2016.5.4.219).
- [109] Muscaritoli M, Arends J, Aapro M. From guidelines to clinical practice: a roadmap for oncologists for nutrition therapy for cancer patients. *Ther Adv Med Oncol.* 2019;11:1758835919880084. doi: [10.1177/1758835919880084](https://doi.org/10.1177/1758835919880084).
- [110] Ferrat E, Audureau E, Paillaud E, et al. Four distinct health profiles in older patients with cancer: latent class analysis of the prospective elcapa cohort. *J Gerontol A Biol Sci Med Sci.* 2016; 71(12):1653–1660. doi: [10.1093/gerona/glw052](https://doi.org/10.1093/gerona/glw052).
- [111] Muscaritoli M, Arends J, Bachmann P, et al. ESPEN practical guideline: clinical Nutrition in cancer. *Clin Nutr.* 2021;40(5):2898–2913. doi: [10.1016/j.clnu.2021.02.005](https://doi.org/10.1016/j.clnu.2021.02.005).
- [112] Maddocks M. Physical activity and exercise training in cancer patients. *Clin Nutr ESPEN.* 2020;40:1–6. doi: [10.1016/j.clnesp.2020.09.027](https://doi.org/10.1016/j.clnesp.2020.09.027).
- [113] Dale W, Klepin HD, Williams GR, et al. Practical assessment and management of vulnerabilities in older patients receiving systemic cancer therapy: ASCO Guideline Update. *J Clin Oncol.* 2023;41(26):4293–4312. <https://ascopubs.org/doi/10.1200/JCO.23.00933>. doi: [10.1200/JCO.23.00933](https://doi.org/10.1200/JCO.23.00933).
- [114] Prado CM, Laviano A, Gillis C, et al. Examining guidelines and new evidence in oncology nutrition: a position paper on gaps and opportunities in multimodal approaches to improve patient care. *Support Care Cancer.* 2022;30(4):3073–3083. doi: [10.1007/s00520-021-06661-4](https://doi.org/10.1007/s00520-021-06661-4).
- [115] Kakehi S, Wakabayashi H, Inuma H, et al. Rehabilitation nutrition and exercise therapy for Sarcopenia. *World J Mens Health.* 2021; 40(1):1–10. doi: [10.5534/wjmh.200190](https://doi.org/10.5534/wjmh.200190).
- [116] Banda KJ, Chu H, Chen R, et al. Prevalence of oropharyngeal dysphagia and risk of pneumonia, malnutrition, and mortality in adults aged 60 years and older: a meta-analysis. *Gerontology.* 2022;68(8):841–853. doi: [10.1159/000520326](https://doi.org/10.1159/000520326).
- [117] Lin LC, Wu SC, Chen HS, et al. Prevalence of impaired swallowing in institutionalized older people in Taiwan. *J Am Geriatr Soc.* 2002;50(6):1118–1123. doi: [10.1046/j.1532-5415.2002.50270.x](https://doi.org/10.1046/j.1532-5415.2002.50270.x).
- [118] Barczy SR, Sullivan PA, Robbins JA. How should dysphagia care of older adults differ? Establishing optimal practice patterns. *Semin Speech Lang.* 2000;21(4):347–361. doi: [10.1055/s-2000-8387](https://doi.org/10.1055/s-2000-8387).
- [119] Cabré M, Almirall J, Clavé P. Aspiration pneumonia: management in Spain. *Eur Geriatr Med.* 2011;2(3):180–183. doi: [10.1016/j.eurger.2011.03.004](https://doi.org/10.1016/j.eurger.2011.03.004).
- [120] Mateos-Nozal J, Montero-Erassquín B, Sánchez García E, et al. High prevalence of oropharyngeal dysphagia in acutely hospitalized patients aged 80 years and older. *J Am Med Dir Assoc.* 2020;21(12):2008–2011. doi: [10.1016/j.jamda.2020.04.032](https://doi.org/10.1016/j.jamda.2020.04.032).
- [121] Mateos-Nozal J, García ES, Rodríguez ER, et al. Oropharyngeal dysphagia in older patients with hip fracture. *Age Ageing.* 2021; 50(4):1416–1421. doi: [10.1093/ageing/afab032](https://doi.org/10.1093/ageing/afab032).
- [122] Frowen J, Hughes R, Skeat J. The prevalence of patient-reported dysphagia and oral complications in cancer patients. *Support Care Cancer.* 2020;28(3):1141–1150. doi: [10.1007/s00520-019-04921-y](https://doi.org/10.1007/s00520-019-04921-y).
- [123] Carrión S, Cabré M, Monteis R, et al. Oropharyngeal dysphagia is a prevalent risk factor for malnutrition in a cohort of older patients admitted with an acute disease to a general hospital. *Clin Nutr.* 2015;34(3):436–442. doi: [10.1016/j.clnu.2014.04.014](https://doi.org/10.1016/j.clnu.2014.04.014).
- [124] Feng H-Y, Zhang P-P, Wang X-W. Presbyphagia: dysphagia in the elderly. *World J Clin Cases.* 2023;11(11):2363–2373. doi: [10.12998/wjcc.v11.i11.2363](https://doi.org/10.12998/wjcc.v11.i11.2363).
- [125] Gupte T, Knack A, Cramer JD. Mortality from aspiration pneumonia: incidence, trends, and risk factors. *Dysphagia.* 2022;37(6): 1493–1500. doi: [10.1007/s00455-022-10412-w](https://doi.org/10.1007/s00455-022-10412-w).
- [126] Clavé P, Shaker R. Dysphagia: current reality and scope of the problem. *Nat Rev Gastroenterol Hepatol.* 2015;12(5):259–270. doi: [10.1038/nrgastro.2015.49](https://doi.org/10.1038/nrgastro.2015.49).
- [127] Clavé P, Rofes L, Arreola V, et al. Diagnosis and management of oropharyngeal dysphagia and its nutritional and respiratory complications in the elderly. *Gastroenterol Res Pract.* 2011;2011: 979. doi: [10.1155/2011/818979](https://doi.org/10.1155/2011/818979).
- [128] Baijens LWJ, Clavé P, Cras P, et al. European society for swallowing disorders – European union geriatric medicine society white

- paper: oropharyngeal dysphagia as a geriatric syndrome. *Clin Interv Aging*. 2016;11:1403–1428. doi: [10.2147/CIA.S107750](https://doi.org/10.2147/CIA.S107750).
- [129] Balcerak P, Corbiere S, Zubal R, et al. Post-stroke Dysphagia: prognosis and treatment – a systematic review of RCT on interventional treatments for dysphagia following subacute stroke. *Front Neurol*. 2022;13:823189. doi: [10.3389/fneur.2022.823189](https://doi.org/10.3389/fneur.2022.823189).
- [130] Daniels S. Neurological disorders affecting oral, pharyngeal swallowing. *GI Motil Online*. 2006;2210. Available from <http://www.nature.com/gimo/contents/pt1/full/gimo34.html>.
- [131] Fujishima I, Fujiu-Kurachi M, Arai H, et al. Sarcopenia and dysphagia: position paper by four professional organizations. *Geriatr Gerontol Int*. 2019;19(2):91–97. doi: [10.1111/ggi.13591](https://doi.org/10.1111/ggi.13591).
- [132] Kakehi S, Isono E, Wakabayashi H, et al. Sarcopenic dysphagia and simplified rehabilitation nutrition care process: an update. *Ann Rehabil Med*. 2023;47(5):337–347. <http://www.e-arm.org/journal/view.php?doi=> doi: [10.5535/arm.23101](https://doi.org/10.5535/arm.23101).
- [133] Mori T, Wakabayashi H, Ogawa N, et al. The mass of geniohyoid muscle is associated with maximum tongue pressure and tongue area in patients with sarcopenic dysphagia. *J Nutr Health Aging*. 2021;25(3):356–360. doi: [10.1007/s12603-020-1528-8](https://doi.org/10.1007/s12603-020-1528-8).
- [134] Maeda K, Akagi J. Decreased tongue pressure is associated with sarcopenia and sarcopenic dysphagia in the elderly. *Dysphagia*. 2015;30(1):80–87. doi: [10.1007/s00455-014-9577-y](https://doi.org/10.1007/s00455-014-9577-y).
- [135] Sporns KB, Hanning U, Schmidt R, et al. Volumetric assessment of swallowing muscles: a comparison of CT and MRI segmentation. *Rofo*. 2018;190(5):441–446. doi: [10.1055/s-0043-120529](https://doi.org/10.1055/s-0043-120529).
- [136] Ogawa N, Mori T, Fujishima I, et al. Ultrasonography to Measure swallowing muscle mass and quality in older patients with sarcopenic dysphagia. *J Am Med Dir Assoc*. 2018;19(6):516–522. doi: [10.1016/j.jamda.2017.11.007](https://doi.org/10.1016/j.jamda.2017.11.007).
- [137] Nagano A, Onaka M, Maeda K, et al. Prevalence and characteristics of the course of dysphagia in hospitalized older adults. *Nutrients*. 2023;15(20):4371. <https://www.mdpi.com/2072-6643/15/20/4371>. doi: [10.3390/nu15204371](https://doi.org/10.3390/nu15204371).
- [138] Cha S, Kim WS, Kim KW, et al. Sarcopenia is an independent risk factor for dysphagia in community-dwelling older adults. *Dysphagia*. 2019;34(5):692–697. doi: [10.1007/s00455-018-09973-6](https://doi.org/10.1007/s00455-018-09973-6).
- [139] Wakabayashi H, Takahashi R, Murakami T. The prevalence and prognosis of sarcopenic dysphagia in patients who require dysphagia rehabilitation. *J Nutr Health Aging*. 2019;23(1):84–88. doi: [10.1007/s12603-018-1117-2](https://doi.org/10.1007/s12603-018-1117-2).
- [140] Zhao WT, Yang M, Wu HM, et al. Systematic review and meta-analysis of the association between sarcopenia and dysphagia. *J Nutr Health Aging*. 2018;22(8):1003–1009. doi: [10.1007/s12603-018-1055-z](https://doi.org/10.1007/s12603-018-1055-z).
- [141] Calles M, Wirth R, Labeit B, et al. Sarcopenic dysphagia revisited: a cross-sectional study in hospitalized geriatric patients. *Nutrients*. 2023;15(12):15. doi: [10.3390/nu15122662](https://doi.org/10.3390/nu15122662).
- [142] Nagano A, Ueshima J, Tsutsumiuchi K, et al. Effect of tongue strength on clinical outcomes of patients: a systematic review: tongue strength and clinical outcomes. *Arch Gerontol Geriatr*. 2022;102:104749. doi: [10.1016/j.archger.2022.104749](https://doi.org/10.1016/j.archger.2022.104749).
- [143] Campo-Rivera N, Ocampo-Chaparro JM, Carvajal-Ortiz R, et al. Sarcopenic dysphagia is associated with mortality in institutionalized older adults. *J Am Med Dir Assoc*. 2022;23(10):1720.e11–1720.e17. doi: [10.1016/j.jamda.2022.06.016](https://doi.org/10.1016/j.jamda.2022.06.016).
- [144] Chen KC, Jeng Y, Wu WT, et al. Sarcopenic dysphagia: a narrative review from diagnosis to intervention. *Nutrients*. 2021;13(11):4043. doi: [10.3390/nu13114043](https://doi.org/10.3390/nu13114043).
- [145] de Sire A, Ferrillo M, Lippi L, et al. Sarcopenic dysphagia, malnutrition, and oral frailty in elderly: a comprehensive review. *Nutrients*. 2022;14(5):982. doi: [10.3390/nu14050982](https://doi.org/10.3390/nu14050982).
- [146] Nagano A, Nishioka S, Wakabayashi H. Rehabilitation nutrition for iatrogenic sarcopenia and sarcopenic dysphagia. *J Nutr Health Aging*. 2019;23(3):256–265. doi: [10.1007/s12603-018-1150-1](https://doi.org/10.1007/s12603-018-1150-1).
- [147] Bath PM, Lee HS, Everton LF. Swallowing therapy for dysphagia in acute and subacute stroke. *Cochrane Database Syst. Rev*. 2018;10(10):CD000323.
- [148] Maeda K, Akagi J. Treatment of sarcopenic dysphagia with rehabilitation and nutritional support: a comprehensive approach. *J Acad Nutr Diet*. 2016;116(4):573–577. doi: [10.1016/j.jand.2015.09.019](https://doi.org/10.1016/j.jand.2015.09.019).
- [149] Wakabayashi H, Uwano R. Rehabilitation nutrition for possible sarcopenic dysphagia after lung cancer surgery. *Am J Phys Med Rehabil*. 2016;95(6):e84–e89. doi: [10.1097/PHM.0000000000000458](https://doi.org/10.1097/PHM.0000000000000458).
- [150] Hashida N, Shamoto H, Maeda K, et al. Rehabilitation and nutritional support for sarcopenic dysphagia and tongue atrophy after glossectomy: a case report. *Nutrition*. 2017;35:128–131. doi: [10.1016/j.nut.2016.11.003](https://doi.org/10.1016/j.nut.2016.11.003).
- [151] Nakayama E, Tohara H, Sato M, et al. Time course and recovery of the movements of hyoid bone and thyroid cartilage during swallowing in a patient with sarcopenic dysphagia. *Am J Phys Med Rehabil*. 2020;99(5):E64–E67. doi: [10.1097/PHM.0000000000001211](https://doi.org/10.1097/PHM.0000000000001211).
- [152] Yamada Y, Shamoto H, Maeda K, et al. Home-based combined therapy with rehabilitation and aggressive nutrition management for a Parkinson's disease patient with sarcopenic dysphagia: a case report. *Prog Rehabil Med*. 2018;3(0):20180019. doi: [10.2490/prm.20180019](https://doi.org/10.2490/prm.20180019).
- [153] Shimizu A, Fujishima I, Maeda K, et al. Nutritional management enhances the recovery of swallowing ability in older patients with sarcopenic dysphagia. *Nutrients*. 2021;13:596. doi: [10.3390/nu13020596](https://doi.org/10.3390/nu13020596).
- [154] Nagano A, Maeda K, Koike M, et al. Effects of physical rehabilitation and nutritional intake management on improvement in tongue strength in sarcopenic patients. *Nutrients*. 2020;12(10):3104. doi: [10.3390/nu12103104](https://doi.org/10.3390/nu12103104).
- [155] Chinese Expert Consensus Group of Dysphagia and Nutrition Management. Chinese expert consensus on food and nutrition management for dysphagia (2019 version). *Asia Pac J Clin Nutr*. 2020;29(2):434–444.
- [156] Ueshima J, Shimizu A, Maeda K, et al. Nutritional management in adult patients with dysphagia: position paper from Japanese working group on integrated nutrition for dysphagic people. *J Am Med Dir Assoc*. 2022;23(10):1676–1682. doi: [10.1016/j.jamda.2022.07.009](https://doi.org/10.1016/j.jamda.2022.07.009).
- [157] Dantas RO, Kern MK, Massey BT, et al. Effect of swallowed bolus variables on oral and pharyngeal phases of swallowing. *Am J Physiol – Gastrointest Liver Physiol*. 1990;258(5 Pt 1):G675–81.
- [158] Clavé P, De Kraa M, Arreola V, et al. The effect of bolus viscosity on swallowing function in neurogenic dysphagia. *Aliment Pharmacol Ther*. 2006;24(9):1385–1394. doi: [10.1111/j.1365-2036.2006.03118.x](https://doi.org/10.1111/j.1365-2036.2006.03118.x).
- [159] Burgos R, Bretón I, Cereda E, et al. ESPEN guideline clinical nutrition in neurology. *Clin Nutr*. 2018;37(1):354–396. doi: [10.1016/j.clnu.2017.09.003](https://doi.org/10.1016/j.clnu.2017.09.003).
- [160] O'Keeffe ST. Use of modified diets to prevent aspiration in oropharyngeal dysphagia: Is current practice justified? *BMC Geriatr*. 2018;18(1):167. doi: [10.1186/s12877-018-0839-7](https://doi.org/10.1186/s12877-018-0839-7).
- [161] Garcia-Almeida JM, López Alarcón PL, Layola-Briás M, et al. Adherence, tolerance and consumption habits of a texture-adapted oral nutritional supplement in a population of malnourished patients with dysphagia. *Clin Nutr ESPEN*. 2023;54:555. doi: [10.1016/j.clnesp.2022.09.287](https://doi.org/10.1016/j.clnesp.2022.09.287).
- [162] van der Maarel-Wierink CD, Vanobbergen JNO, Bronkhorst EM, et al. Oral health care and aspiration pneumonia in frail older people: a systematic literature review. *Gerodontology [Internet]*. 2013;30(1):3–9. <https://onlinelibrary.wiley.com/doi/10.1111/j.1741-2358.2012.00637.x>. doi: [10.1111/j.1741-2358.2012.00637.x](https://doi.org/10.1111/j.1741-2358.2012.00637.x).