

Rita Rodrigues¹, Zilda Morais¹, Raquel Lisboa^{1,2}, Davide Mendes³ and Marco Silva³

¹ CiiEM, Egas Moniz Cooperativa de Ensino Superior, Almada, Portugal.

² Departamento de Biologia, Universidade de Aveiro; Portugal.

³ Departamento de Química, Faculdade de Ciências e Tecnologia, Universidade Nova de Lisboa, Almada, Portugal.

rodrigues.rfb@gmail.com

Introduction

Due to the variety of organisms with different physiologies, the marine environment is considered the main source for new drugs.

The most promising appear to be microorganisms, invertebrates and algae, respectively, being the lipids a chemical class with high potential, especially terpenoids, steroids and eicosanoids [1].

The objective of this study was to initiate the characterization of the lipidic compounds present in the gonads of both sexes of the scyphozoan *Catostylus tagi*, native of the Tagus and Sado estuaries [2] (Fig. 1).




Fig. 1. Left: *C. tagi* in medusa stage. Up: gonads.

Stearic, palmitic and oleic, respectively, were the three most abundant fatty acids in both sexes.

Considering sterols, cholesterol, 24-methylenecholesterol, campesterol and β -sitosterol were unequivocally detected both in male and female.

The [(3β)-cholestca-5,24-dien-3-yloxy], RT 78.094, was confirmed only in male (Fig. 3).

Isofucosterol (Stigmasta-5,24(28)-dien-3-ol, (3β ,24Z)) reported for a jellyfish [3] was not detected.

These findings are in agreement with lipids reported for another cnidarian [4]. Studies focused at unambiguous chemical differentiation of each sex are now being planned.

References

- [1] Rocha J, Calado R, Leal M. *In: Springer Handbook of Marine Biotechnology*. Berlin Heidelberg: Springer; 2015. p. 823-49.
- [2] Morais Z, Schiariti A & Morandini A. *In: Jellyfish, Ecology, Distribution Patterns and Human Interactions*. New York: Nova Science Publishers; 2017. p. 91-145.
- [3] Ballantine JA, Roberts JC, Morris RJ. *Biological Mass Spectrometry*. 1976;3:14-20.
- [4] Revel J et al. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*. 2016;191:140-51.

Materials and Methods

Chloroform soluble compounds were obtained by soxhlet extraction from the lyophilized gonads of *C. tagi* (Fig. 2).



Fig. 2. Sample preparation for GC-MS and previous analysis.

Chromatographic conditions: Samples were injected as trimethylsilyl derivatives. T_i : 250°C; Split ratio: 1:20; gas rate (He): 1mL/min. Column: VF5-ms, 30m, 0.25 nm ID, 0.25 mm film; T_c : 60°C (1min), 2°C/min until 90°C (0 min), 3°C/min until 280°C (40.67 min). Equipment GC/MS: GC Agilent 6890N - MS Thermo DSQ.

Results




Fig. 3. GC-MS of chloroform extracts of gonads of *C. tagi*. Up: female, Down: male.

Acknowledgements

We thank the fishermen, Vítor Reis and Olímpio Simões, Rute Pinheiro from ITN and Luz Fernandes from UNL. This study was funded by Egas Moniz Coop Ensino Superior CRL, project EM-ZM 03/04.