

Home- vs gym-based exercise delivery modes of two multicomponent intensity training regimes on cardiorespiratory fitness and arterial stiffness in adults with intellectual and developmental disability during the COVID-19 pandemic – a randomized controlled trial

Journal of Intellectual Disabilities
2025, Vol. 29(1) 66–85
© The Author(s) 2024



Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/17446295241242507
journals.sagepub.com/home/jid



Xavier Melo

Centro de Investigação Interdisciplinar Egas Moniz (CiiEM), Egas Moniz School of Health & Science, Monte da Caparica, Portugal

Centro Interdisciplinar de Estudo da Performance Humana (CIPER), Faculdade de Motricidade Humana – Universidade de Lisboa, Lisboa, Portugal

Bruno Simão and Catarina Catela

Centro Interdisciplinar de Estudo da Performance Humana (CIPER), Faculdade de Motricidade Humana – Universidade de Lisboa, Lisboa, Portugal

Research & Development Department, GCP Lab, Ginásio Clube Português, Lisboa, Portugal

Isabel Oliveira

Research & Development Department, GCP Lab, Ginásio Clube Português, Lisboa, Portugal

Faculdade de Ciências da Saúde e do Desporto, Universidade Europeia, Lisboa, Portugal

Sara Planche and Ana Louseiro

Research & Development Department, GCP Lab, Ginásio Clube Português, Lisboa, Portugal

João Luís Marôco

Integrative Human Physiology Laboratory, Manning College of Nursing & Health Sciences, University of Massachusetts Boston, Boston, MA, USA

Guillermo R Oviedo

Faculty of Psychology, Education and Sport Science Blanquerna, University Ramon Llull, Barcelona, Spain

Bo Fernhall

Integrative Human Physiology Laboratory, Manning College of Nursing & Health Sciences, University of Massachusetts Boston, Boston, MA, USA

Helena Santa-Clara

Centro Interdisciplinar de Estudo da Performance Humana (CIPER), Faculdade de Motricidade Humana – Universidade de Lisboa, Lisboa, Portugal

Research & Development Department, GCP Lab, Ginásio Clube Português, Lisboa, Portugal

Corresponding author:

Xavier Melo, Egas Moniz School of Health & Science, Caparica 2829-511 Portugal.
Email: xmelo@egasmoniz.edu.pt

Date accepted: 11 March 2024

Abstract

Background: We compared the effects of home- vs gym-based delivery modes of two 8-week supervised multicomponent intensity training regimes on cardiorespiratory fitness and arterial stiffness in 17 adults with intellectual and developmental disability during the COVID-19 pandemic. **Methods:** Participants were assigned to sprint interval training or continuous aerobic training, both incorporating resistance training. The intervention started with 8-weeks of online training (M1-M2), 1-month of detraining, plus 8-weeks of gym-based training (M3-M4). **Results:** Peak oxygen uptake decreased from M1-M2 and increased from M2-M4. Central arterial stiffness decreased between M1-M2, and M1-M4, along with peripheral arterial stiffness. Central systolic blood pressure decreased from M1-M2 only with sprint interval training. **Conclusion:** Home-based training minimized the negative impact of the lockdown on central arterial stiffness and central blood pressure, but it did not match the benefits on cardiorespiratory fitness and peripheral arterial stiffness of a gym-based intervention, irrespective of the multicomponent intensity training regime. Registered in ClinicalTrials.gov NCT05701943.

Keywords

intellectual developmental disability, covid-19, blood pressure, arterial stiffness, cardiorespiratory fitness, physical exercise, multicomponent exercise, delivery mode

Introduction

The COVID-19 pandemic has cast a spotlight on the intensified challenges faced by individuals with deficits in intellectual and adaptive functioning in the conceptual, social and practical domain, referred to hereafter as intellectual and developmental disabilities ([American Psychiatric Association, 2022](#)), exacerbating their health issues, mental well-being, and societal disadvantages ([Emerson and Hatton, 2008](#); [Kirby, 2020](#); [Qiao, 2020](#); [WHO, 2020](#)).

While the mandatory lockdowns have presented substantial hurdles for these individuals in maintaining physical activity levels ([Fitzgerald et al., 2022](#)), the aftermath of the pandemic brings to the forefront the persistent need for flexible and accessible exercise interventions ([Theis et al., 2021](#)). Beyond the immediate risk of infection, this population continues to grapple with multifaceted barriers that curtail their engagement in community-based exercise programs ([Jacinto et al., 2021](#)). It is evident that people with intellectual and developmental disabilities have long faced challenges in meeting physical activity requirements and maintaining optimal cardiorespiratory fitness ([Gawlik et al., 2017](#); [Hilgenkamp and Baynard, 2018](#)). Existing research underscores their susceptibility to cardiovascular risk factors ([Gawlik et al., 2017](#); [Hsieh et al., 2014](#)), highlighting the pressing need for effective interventions that can mitigate these risks. The increased vulnerability of this population to infection and COVID-19 complications is a paramount concern ([Zbinden-Foncea et al., 2020](#)), emphasizing the significance of measures that can both enhance cardiovascular well-being and reduce the risk of infectious diseases.

Prolonged periods of reduced physical activity, a byproduct of lockdowns, can lead to changes in the material properties of the arterial wall ([Bohn et al., 2017](#)). These alterations, in turn, have functional consequences for the artery, influencing the way pressure, blood flow, and arterial diameter change with each heartbeat ([Salvi, 2016](#)). This concept is known as arterial stiffness and can further amplify the risk

of cardiovascular disease (Cavalcante et al., 2011; Hansen et al., 2006; Naqvi and Lee, 2014). Arterial stiffness can be measured in various types of arteries, including muscular and elastic ones, and assessed in cross-section, longitudinally along the vessel, or in both directions. Typically, arterial stiffness is evaluated by measuring the velocity of the pulse-wave travelling in a specific segment, commonly carotid-femoral pulse wave velocity (cfPWV). The pulse wave travels through the aorta, and its velocity is inversely related to the distensibility of the arterial wall. In simpler terms, the higher cfPWV, the lower the vascular distensibility, providing insights into the rigidity of the aorta. However, while the association of time spent in physical activities and sedentary behaviors with cfPWV has been studied in the general population (Germano-Soares et al., 2018), research in people with intellectual and developmental disabilities has remained limited, primarily focusing on individuals with Down syndrome (Rodrigues et al., 2011). This underscores the necessity for comprehensive investigations encompassing various etiologies within this diverse population.

Exercise training, whether aerobic-only (Boer et al., 2014; Melo et al., 2021) or multicomponent, also incorporating resistance training (Mendonca et al., 2011; Oviedo et al., 2014), improves cardiorespiratory fitness (2 to 6 mL.kg.min⁻¹) and brachial systolic blood pressure (bSBP) (-15 to -6 mmHg) in people with intellectual and developmental disabilities. Specifically, Calders et al. (2011) demonstrated greater reductions in bSBP with a 12-week multicomponent training regime compared to an aerobic-only training group. Both the multicomponent and aerobic-only training groups (n = 15 each) exhibited similar increases in aerobic capacity. These findings collectively suggest that exercise training, particularly a multicomponent regime, is beneficial for people with intellectual and developmental disabilities. However, the nuanced effects of exercise intensity and its impact on arterial stiffness in this population remain ambiguous, as similar results have been reported with continuous aerobic training and high-intensity interval training for central and peripheral arterial stiffness (Kim, 2017; Melo et al., 2021). Importantly, these observations have predominantly stemmed from supervised gym or laboratory-based interventions, avenues that became largely inaccessible during the COVID-19 lockdown.

As we navigate the post-pandemic landscape, home-based exercise interventions takes on even greater significance. Challenges like travel time, financial constraints, and anxiety associated with novel environments have perennially hindered the involvement of people with intellectual and developmental disabilities in community-based programs (Jacinto et al., 2021). Online or virtual interventions hold promise in addressing these barriers and providing a more inclusive, accessible, and tailored approach to exercise engagement (Fjellstrom et al., 2022). This assumption is reinforced as the shadow of potential infectious disease outbreaks looms large (Smith, 2021), emphasizing the need for interventions that ensure sustained and consistent access to exercise.

Hence, in view of these complex and evolving considerations, this study endeavored to compare the effectiveness of two distinct 8-week supervised multicomponent intensity training regimes, delivered either at home or at the gym, in improving cardiorespiratory fitness and managing arterial stiffness in people with intellectual and developmental disabilities. We hypothesized that home- and gym-based exercise delivery modes hold equal potential in augmenting cardiorespiratory fitness and overall arterial health in this population, irrespective of the multicomponent intensity training regimes.

Methodology

Design

This study was a 16-week parallel group randomized controlled trial in which participants were randomly assigned in a 1:1 allocation ratio to a multicomponent exercise training regime, either with

sprint interval training or continuous aerobic training. Randomization was done using a random-block scheme obtained from <https://www.randomizer.org/>. Both groups exercised for 60 minutes, 3 times per week. The intervention started with 8 weeks of online training via Google Meets at their homes, followed by 1 month of detraining with no exercise besides those available by the SPORTS4ALL program, and another 8 weeks of on-site training at the gym in GCP. Participants' exercise adherence was evaluated using an exercise diary maintained by the exercise physiologists. The study timeline, intervention, and assessments are presented in [Figure 1](#), and all related trials were retrospectively registered on [ClinicalTrials.gov](https://clinicaltrials.gov) under NCT05701943 due to human, time, and logistical constraints faced during the COVID-19 pandemic. The primary outcomes were cardiorespiratory fitness (peak VO_2) and central arterial stiffness (cfPWV), assessed at GCP laboratory on 4 occasions over a 20-week period: before (M1) and after (M2) the home-based intervention, and before (M3) and after (M4) the gym-based intervention. All laboratory tests were conducted by the same researchers at a room temperature of 22–24°C, in the morning, after a fasting period of 3 hours, no alcohol for 24 hours, no caffeine for 8 hours, and no vigorous exercise for 48 hours before data collection. On the day of testing, participants had their body composition measured, brachial arterial pressure assessed, arterial stiffness evaluated by applanation tonometry, and completed an incremental test to exhaustion.

Participants

Based upon a small effect size of 0.173 derived from a target increase of 0.3 L.min⁻¹ and a SD of 0.4 L.min⁻¹ in peak VO_2 ([Boer and Moss, 2016](#)), 15 participants were required per group, assuming a 95% confidence level, 5% alpha error, and 20% beta error, with a 1:1 ratio between groups. To account for a 20% dropout rate, a total of 36 participants (18 per group) were recruited from the social project SPORTS4ALL by Ginásio Clube Português (GCP). These people have any type of disability or limitation, and the project ensures equality of opportunities in well-being, leisure, and culture through sports. Study advertising took place through invitation by the research team. Thirty-six participants met the following inclusion criteria: ≥ 18 and ≤ 55 years; diagnosed with mild to moderate intellectual and developmental disabilities; exercised at least 1 d/week in the last 2 months; able to participate in group exercise activities with ≥ 8 people; able to walk independently; and able to perform all physical fitness assessments. Four participants were excluded because they had 1 of the following: cardiovascular disease, respiratory disorder, metabolic disease, atlantoaxial instability, severe intellectual and developmental disabilities, smoking, and/or use of heart rate and blood pressure altering or non-steroidal anti-inflammatory medications, inability to comply with guidelines for participation in exercise testing and training ([Riebe et al., 2018](#)). Informed written consent was signed by all participants and parent(s) and legal guardian(s), and participants or parents/legal guardians filled-out the Physical Activity Readiness Questionnaire for Everyone, and a sociodemographic questionnaire. By the time the mandatory COVID-19 home confinement was declared by the Portuguese government, only 17 participants had completed laboratory exercise testing. Only these participants took part in this study. This study was approved by the Ethical Committee of Faculdade de Motricidade Humana – Universidade de Lisboa (46/2021).

Exercise interventions

The study implemented exercise programs for a duration of 8 weeks, which were delivered to participants in their homes via Google Meets, as well as at the gym. The sessions were designed by




Figure 1. Participant flow diagram. Abbreviations: SIT = sprint interval training; CAT = continuous aerobic exercise training.

exercise physiologists and rehabilitation technicians in adherence to the American College of Sports Medicine (ACSM) guidelines (Riebe et al., 2018) and the National Strength and Conditioning Association (NSCA) (Haff et al., 2016). Exercise physiologists underwent standardized training and a 2-month familiarization period. Both multicomponent intensity training regimes comprising aerobic, resistance, balance, and flexibility exercises, were divided into 2 groups, each with 8-9 participants, maintaining a participant-exercise physiologist ratio of 3:1. In each exercise session, whether home- or gym-based, at least one rehabilitation technician was also present. Both programs were divided into three phases, which were adapted from Oviedo et al., (2020): the initial phase (weeks 1-3), the improvement phase (weeks 4-6), and the maintenance phase (weeks 7-8). The multicomponent intensity training regimes were matched for total duration, and all sessions included a standardized 5-minute warm-up and cool-down exercises. During the washout period, 14 participants underwent 3 sessions, each corresponding to a phase of the home-based intervention. These sessions were conducted while participants wore heart rate monitors, providing a proxy for exercise intensity during each phase of the home-based intervention. Both multicomponent intensity

training regimes comprised a standard resistance training program, consisting of 6 exercises targeting various muscle groups. For the upper limbs, these exercises comprised seated biceps curls, triceps extensions, and frontal shoulder raises. For the torso, seated rows and side bends were included, while chair squats were performed for the lower limbs. All exercises were executed using either 2-4 kg dumbbells or the participants' own body weight, based on individual capabilities. To monitor resistance training intensity, the 0-10 points Rate of Perceived Exertion OMNI-Resistance Exercise Scale (RPE OMNI-RES) (Gearhart et al., 2009) was employed. The intensity was kept constant at 2 x 12 repetitions at RPE OMNI-RES 8 throughout the study.

Multicomponent continuous aerobic training regime. The multicomponent continuous aerobic training regime, henceforth referred to as continuous aerobic training, encompassed a comprehensive set of exercises, incorporating aerobic, resistance, balance, and flexibility exercises (see [Supplement 1](#)). For the aerobic component, participants engaged in moderate intensity, steady-state exercises for an extended period, tailored to their intervention location. In the home-based intervention, participants engaged in bodyweight exercises, such as low-impact jumping jacks, slow standing box, slow side shift with floor touch, slow high knees, half burpee (without jump), and hook box (Borrega-Mouquinho et al., 2021). In the gym-based intervention, participants used cycle ergometers (Star Trac Spinner Blade ION 7220, Vancouver, WA) for their aerobic component. Each session began with a 5-minute warm-up, followed by three continuous cycling bouts at a steady-state intensity. The duration of each bout (5-10 minutes) and the intensity (55-85% heart rate reserve) were progressively increased in each phase. Participants exercised at the target intensity using the OMNI scale 4-6 (Stanish and Aucoin, 2007) during home-based intervention and an heart rate chest band (H10 Polar, Electro, Kempele, Finland) during gym-based intervention.

Multicomponent sprint interval training regime. The multicomponent sprint interval training regime, henceforth referred to as "sprint interval training", a form of high intensity interval training, also encompassed a comprehensive set of exercises, incorporating aerobic, resistance, balance, and flexibility exercises ([Supplement 2](#)). However, the aerobic component featured short bursts of high-intensity sprints followed by periods of lower-intensity activity or rest, performed differently depending on the location of the intervention. In the home-based intervention, bodyweight exercises such as jumping jacks, standing box, side shift with floor touch, high knees, half burpee, and hook box were used (Borrega-Mouquinho et al., 2021). In the gym-based intervention, cycle ergometers (Star Trac Spinner Blade ION 7220, Vancouver, WA) were utilized instead. The aerobic component began with a 5-minute warm-up, followed by bouts of 5-10 minutes of exercise, consisting of 5-20 second all-out sprints followed by 15-45 seconds of low cadence recovery (1:3-1:2 work-rest ratio). The duration of sprints and active recovery were adjusted throughout the program. The OMNI scale 8-10 (Stanish and Aucoin, 2007) was used to ensure that participants exercised at the target intensity during the home intervention, and an heart rate chest band (H10 Polar, Electro, Kempele, Finland) was used during the gym intervention.

Cardiopulmonary exercise test

Participants completed a ramp incremental cycle ergometer test to exhaustion on a calibrated electronically braked cycle ergometer (Monark 839 E, Ergomedic; Monark, Vansbro, Sweden) at a pedal cadence of 70-75 rev.min⁻¹ with initial and incremental workloads of 10-20 watts. Respiratory gases were continuously analyzed, with a mixing-chamber method using a portable gas analyzer (K5, Cosmed, Rome, Italy), calibrated according to the manufacturer's instructions. HR was

monitored continuously (Garmin, US) and data were evaluated in 10-s averages. Peak VO_2 was defined as the highest 20-s value attained in the last minute of effort when 2 of the following criteria were met: (1) $\sim 90\%$ of predicted maximal HR: $210 - 0.56(\text{age}) - 15.5$ (Down Syndrome) (Fernhall et al., 2001); (2) Plateau in VO_2 with an increase in workload ($< 2.0 \text{ mL} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$); (3) Respiratory exchange ratio ≥ 1.1 ; and/or (4) subjective judgment that the participant could no longer continue. Results were compared to age and sex normative data (Pescatello et al., 2014). Chronotropic responses were calculated as HR reserve/ (predicted/maximal HR - HR at rest)*100 (Azarbal et al., 2004). Chronotropic incompetence was defined as the failure to reach 80% of the chronotropic response (Brubaker and Kitzman, 2011). HR recovery was calculated as the difference in heart rate after 1-min of recovery in relation to the peak heart rate. An abnormal heart rate recovery was defined as reductions of less than $12 \text{ b} \cdot \text{min}^{-1}$ (Cole et al., 1999). The first ventilatory threshold (VT1) was identified as 1) the first increase in the ventilatory equivalent for O_2 (VE/VO_2), without any increases in the ventilatory equivalent for CO_2 (VE/VCO_2), and 2) the first increase in the expiratory fraction of O_2 . The first increase in the ventilatory equivalent for CO_2 (VE/VCO_2) and the first reduction in the expiratory fraction of CO_2 were identified as the second ventilatory threshold (VT2) (Meyer et al., 2005).

Regional arterial stiffness and blood pressure

The study utilized piezoelectric pressure mechanotransducers to measure arterial stiffness through PWV (Complior Analyse, ALAM Medical, Paris, France). The pressure waveform travel distance (d) was measured between the carotid-femoral (cf), and carotid-distal posterior tibial arteries (cd) using tape measures, and the Pulse transit time (PTT) was calculated using the intersect tangent algorithm of the foot-to-foot method. The measurements were taken by a single operator on the right side of the body after 10 high quality pulse waveforms ($> 90\%$). The cfPWV and cdPWV were used as indices of central and peripheral arterial stiffness, respectively (Salvi et al., 2019; Townsend et al., 2015). The central SBP (cSBP) was assessed using the piezoelectric pressure mechanotransducers placed in the carotid artery, calibrated from the brachial mean arterial pressure (bMAP), and the mean values were extracted from a 15-s window of acquisition. The cfPWV was compared to established European reference values (The Reference Values for Arterial Stiffness' Collaboration, 2010a). The inter-day CV for cfPWV and cdPWV were 1.12% and 3.02%, respectively.

Body composition

Height, waist circumference, and weight were measured to the nearest 0.1 cm and nearest 0.1 kg, respectively, on a scale with an attached stadiometer (model 770, Seca; Hamburg, Deutschland). Body composition was measured using a seca mBCA 515 using 8 electrodes.

Statistical analyses

All statistical analyzes were conducted with a significant level (α) of 0.05 using the R software, version 4.1.2. We performed an intention-to-treat analysis using all randomized participants. The data are presented as mean and standard deviation. To compare baseline characteristics Welch's independent t-tests were used.

Changes in the incidence of participants with low cardiorespiratory fitness, high bSBP, cfPWV, and BMI throughout the study period were tested using generalized estimating equation models.

The models were fitted following a Poisson distribution using the *geepack* package (Halekoh et al., 2006).

The changes in the cardiorespiratory fitness and vascular outcomes were examined using linear mixed models fitted with restricted maximum likelihood and applying Satterthwaite's method for approximating degrees of freedom for the F test from the *R lmerTest* package (Kuznetsova et al., 2017). The fixed effects were time (home- and gym-based) and multicomponent intensity training regime (continuous aerobic training and sprint interval training), whereas the random intercept was defined for each participant. All models' residuals were tested for normality and homogeneity of variances using both Shapiro-Wilk and Breusch-Pagan tests, respectively, and via QQ plot inspection using the *R performance* package (Lüdecke, Makowski, et al., 2020b) Using the *R sjstats* package (Lüdecke, Ben-Shachar, et al., 2020a), partial eta squares (η^2) were calculated for main effects and interaction and interpreted using the benchmarks suggested by Cohen (1988) defining small ($\eta^2 < 0.05$), medium ($\eta^2 < 0.25$), and large ($\eta^2 > 0.25$) effect sizes. In the presence of significant main effects and interactions, post-hoc comparisons using Tukey's HSD test were conducted using the *R emmeans* package (Lenth R, 2020).

Results

Characterization of the exercise training programs

The baseline characteristics of the participants are presented in [Table 1](#). The home-based intervention attendance rate was 81% (sprint interval training = 92%; continuous aerobic training = 69%) with a mean of 19 (sprint interval training = 22 sessions; continuous aerobic training = 17 sessions) out of 24 training sessions offered. Four participants in continuous aerobic training and 1 participant in sprint interval training attended less than 75% of the home-based sessions. The gym-based intervention attendance rate was 79% (sprint interval training = 77%; continuous aerobic training = 80%) with a mean of 19 (sprint interval training = 19; continuous aerobic training = 19) out of 24 training sessions offered. Three participants in sprint interval training attended less than 75% of the gym-based intervention (38%, 63%, and 71%). Reasons behind the insufficient attendance rates included illness (n = 1), schedule mismatch (n = 5), and technical issues during the home-based sessions (n = 2). No instances of harm, adverse events, or SARS-CoV-2 virus infections were reported throughout the duration of the study.

Home-based and gym-based exercise physiological demands are depicted in [Supplement 4](#). Mean heart rate (difference (d) = 15, 95% CI: -1 to 31 b.min⁻¹, $p = 0.06$) and %heart rate reserve (d = 2, 95% CI: -6 to 10 b.min⁻¹, $p = 0.54$) during sprint interval training and continuous aerobic training in the home-based intervention did not differ. In the gym-based intervention, the mean heart rate (d = 25 b.min⁻¹, 95% CI: 9 to 40 b.min⁻¹, $p = 0.01$), but not the mean %heart rate reserve (d = 5, 95% CI: -3 to 13 b.min⁻¹, $p = 0.17$), was higher during sprint interval training compared to continuous aerobic training, respectively. There were no differences in heart rate and %heart rate reserve between exercise delivery modes ([Supplement 3](#))

Cardiorespiratory fitness

No baseline differences were observed between multicomponent intensity training regimes in cardiorespiratory fitness indices ([Table 2](#)). All participants remained classified as unfit from M1 to M4. Abnormal heart rate recovery and chronotropic incompetence were present in 24% and 18% of the participants, respectively. A main effect of time was observed for peak VO_2 [F (3,39) = 4.68; $p = 0.007$;

Table 1. Baseline characteristics of the participants by multicomponent intensity training regime.

Variables	SIT (n=8)	CAT (n=9)	(p-value)
Characteristics			
Age (years)	26±8	29±13	$p = 0.46$
Sex (male/female)	6/2	5/4	$p = 0.43$
Resting heart rate (b.min ⁻¹)	80±11	79±14	$p = 0.88$
Etiology			
Autism spectrum disorder	3	0	$p = 0.04$
Down Syndrome	2	3	$p = 0.33$
Global Development Delay	1	4	$p = 0.17$
Intellectual Disability	2	1	$p = 0.48$
Williams Syndrome	0	1	$p = 0.36$

Data are presented as mean ± standard deviation. Abbreviations: SIT = sprint interval training; CAT = continuous aerobic exercise training.

$\eta^2 = 0.26$], suggesting a decrease from M1 to M2 ($d = -1.8$, 95% CI: -0.2 to -3.4 mL.kg.min⁻¹, $p = 0.02$), and an increase from M2 to M4 ($d = 2.1$, 95% CI: 0.4 to 3.8 mL/kg/min, $p = 0.01$; [Figure 2](#)), independent of age, sex, and fat mass. Main effects of time were also observed for power output [F (3,37) = 5.07; $p = 0.005$; $\eta^2 = 0.29$] and for time-to-exhaustion [F (3,38) = 3.37; $p = 0.03$; $\eta^2 = 0.21$]. These suggested decreases from M1 to M2 ($d = -27$, 95% CI: -49 to 5 W, $p = 0.01$), but increases ($d = 26$, 95% CI: 3 to 50, $p = 0.02$) from M2 to M4 in power output, together with CPET duration ($d = 104$, 95% CI: 14 to 194 s, $p = 0.02$). A multicomponent intensity training regime-by-time interaction effect was observed for VT2 [F (3,39) = 3.93; $p = 0.02$; $\eta^2 = 0.23$], suggesting that VT2 decreased from M1 to M2 in sprint interval training ($d = -5.2$, 95% CI: -8.8 to -1.6 mL.kg.min⁻¹, $p < 0.01$), and from M2 to M3, M2 to M4 in continuous aerobic training ($d = 3.8$, 95% CI: 0.1 to 7.6 mL/kg/min, $p = 0.04$ and $d = 4.7$, 95% CI: 0.9 to 8.4 mL.kg.min⁻¹, $p = 0.01$, respectively), independent of age, sex, and peak VO₂.

Arterial stiffness

No baseline differences were observed between multicomponent intensity training regimes in arterial stiffness indices ([Figure 2](#)). A main effect of time was observed in the incidence of participants with cfPWV above +2SD of the age- and blood pressure-specific mean reference values ($\chi^2 = 9617$, $p < 0.001$). At baseline, 24% of participants exhibited high central arterial stiffness, at M2 only 12% ($p = 0.383$), at M3 only 7% ($p = 0.260$) and at M4 ($p < 0.001$) none of the participants.

Main effects of time were observed for cfPWV [F (3,38) = 5.15; $p = 0.004$; $\eta^2 = 0.29$] and for cdPWV [F (3,39) = 3.32; $p = 0.03$; $\eta^2 = 0.29$]. These suggested a reduction in cfPWV from M1 to M2 ($d = -0.61$, 95% CI: -1.1 to -0.1 m.s⁻¹, $p = 0.01$), and in both cfPWV ($d = -0.63$, 95% CI: -1.1 to -0.1 m.s⁻¹, $p = 0.01$) and cdPWV ($d = -0.72$, 95% CI: -1.4 to -0.1 m/s, $p = 0.03$; [Figure 3](#)) from M1 to M4. Changes in cfPWV, but not in cdPWV, remained significant after adjustment for bMAP, peak VO₂, height, fat mass, resting heart rate, and age.

Blood pressure

No baseline differences were observed between multicomponent intensity training regimes in hemodynamic parameters ([Table 3](#)). The incidence of elevated bSBP ([Casey et al., 2019](#)) was 65%

Table 2. Comparison of indices of cardiorespiratory fitness among delivery exercise modes and multicomponent intensity training regimes.

Cardiorespiratory fitness	SIT	CAT	Main effect of time (p-value; partial eta square)	Main effect of multicomponent intensity training regimes (p-value; partial eta square)	Interaction (p-value; partial eta square)
Time to peak power (s)			($p = 0.02$; $\eta^2 = 0.21$)	($p = 0.19$; $\eta^2 = 0.13$)	($p = 0.98$; $\eta^2 = 0$)
M1	557±132	625±193	-	-	-
M2	497±75	553±92	M2 < M4	-	-
M3	540±79	623±90	-	-	-
M4	595±176	663±72	-	-	-
Peak Power (W)			($p < 0.01$; $\eta^2 = 0.29$)	($p = 0.71$; $\eta^2 = 0.00$)	($p = 0.71$; $\eta^2 = 0.04$)
M1	205±25	192±39	M1 > M2	-	-
M2	168±25	174±26	M2 < M4	-	-
M3	184±20	177±35	-	-	-
M4	200±24	195±45	-	-	-
Heart rate max (b.min ⁻¹)			($p = 0.03$; $\eta^2 = 0.19$)	($p = 0.10$; $\eta^2 = 0.17$)	($p = 0.44$; $\eta^2 = 0.07$)
M1	159±13	151±29	-	-	-
M2	158±17	143±26	-	-	-
M3	146±21	139±19	-	-	-
M4	143±19	143±24	-	-	-
RER			($p = 0.18$; $\eta^2 = 0.12$)	($p = 0.10$; $\eta^2 = 0.17$)	($p = 0.46$; $\eta^2 = 0.06$)
M1	1.16±0.14	1.00±0.10	-	-	-
M2	1.11±0.16	1.07±0.10	-	-	-
M3	1.11±0.14	1.05±0.09	-	-	-
M4	1.12±0.18	1.00±0.10	-	-	-
Heart rate recovery 1 st Minute (b.min ⁻¹)			($p = 0.48$; $\eta^2 = 0.06$)	($p = 0.83$; $\eta^2 = 0.00$)	($p = 0.57$; $\eta^2 = 0.05$)
M1	19±11	18±6	-	-	-
M2	21±8	24±8	-	-	-
M3	22±8	18±5	-	-	-
M4	22±9	20±11	-	-	-
VT1 (mL.kg.min ⁻¹)			($p = 0.06$; $\eta^2 = 0.17$)	($p = 0.23$; $\eta^2 = 0.10$)	($p = 0.52$; $\eta^2 = 0.05$)
M1	16.8±5.0	13.7±5.0	-	-	-
M2	12.7±4.7	10.4±6.0	-	-	-
M3	16.2±9.0	10.8±2.0	-	-	-
M4	15.5±6.2	15.1±5.0	-	-	-
VT2 ((mL.kg.min ⁻¹))			($p < 0.01$; $\eta^2 = 0.31$)	($p = 0.19$; $\eta^2 = 0.11$)	($p = 0.02$; $\eta^2 = 0.23$)

(continued)

Table 2. (continued)

Cardiorespiratory fitness	SIT	CAT	Main effect of time (p-value; partial eta square)	Main effect of multicomponent intensity training regimes (p-value; partial eta square)	Interaction (p-value; partial eta square)
M1	26.2±6.2	18.6±6.5	M1 > M2	-	SIT M1 > SIT M2
M2	21.0±5.7	16.4±5.1	-	-	-
M3	23.1±6.6	19.7±7.2	M3 > M2	-	CAT M3 > CAT M2
M4	23.0±6.0	20.5±5.5	M4 > M2	-	CAT M4 > CAT M2

Data are presented as mean ± standard deviation. Abbreviations: RER = respiratory exchange ratio; VT1 = first ventilatory threshold; VT2 = second ventilatory threshold; SIT = sprint interval training; CAT = continuous aerobic exercise training; M = moment.




Figure 2. Comparison of central (cfPWV) and peripheral arterial (cdPWV) stiffness among delivery exercise modes and multicomponent intensity training regimes.




Figure 3. Comparison of peak oxygen uptake and diastolic blood pressure among delivery exercise modes and multicomponent intensity training regimes.

at M1, 47% at M2, 35% at M3, and 35% at M4 but these changes did not reach statistical significance ($\chi^2 = 2.40, p = 0.49$). Main effects of time were observed for bDBP [$F(3,39) = 5.32; p = 0.004; \eta^2 = 0.29$, Figure 1] and bMAP [$F(3,39) = 5.28; p = 0.004; \eta^2 = 0.29$]. These supported a decrease from M1 to M4 (bDBP: $d = -6$, 95% CI: -10 to -2 mmHg, $p = 0.002$; bMAP: $d = -6$, 95% CI: -10 to -2 mmHg, $p = 0.003$), from M2 and M4 (bDBP: $d = -5$, 95% CI: -9 to -1 mmHg, $p = 0.03$; bMAP: $d = -4$, 95% CI: -9 to -1 mmHg, $p = 0.03$), independently of age, sex, cfPWV, peak VO_2 , and fat mass. A multicomponent intensity training regime-by-time interaction effect was observed for cSBP [$F(3,39) = 2.85; p = 0.04; \eta^2 = 0.18$], suggesting a decrease from M1 to M2 in sprint interval training ($d = -17$, 95 to -1 mmHg, $p = 0.03$), independently of age, sex, and peak VO_2 , but not cfPWV and fat mass.

Body composition

No baseline differences were observed between multicomponent intensity training regimes in body composition outcomes at M1. At baseline, 41% of the participants were overweight and 12% obese, while at M3, 24% were overweight and 12% obese ($\chi^2 = 1.08, p = 0.78$) based on body mass index (BMI). Increased central fat accumulation was observed in 65% of the participants at M1, and 50% at M3. A main effect of time was observed for %fat mass [$F(3,38) = 3.76; p = 0.02; \eta^2 = 0.23$],

Table 3. Comparison of hemodynamics indices among delivery exercise modes and multicomponent intensity training regimes.

Hemodynamics	SIT	CAT	Main effect of time (p-value; partial eta square)	Main effect of multicomponent intensity training regimes (p-value; partial eta square)	Interaction (p-value; partial eta square)
bSBP (mmHg)			($p = 0.23$; $\eta^2 = 0.11$)	($p = 0.81$; $\eta^2 = 0.00$)	($p = 0.19$; $\eta^2 = 0.11$)
	M1	125±13	122±19	-	-
	M2	122±12	124±18	-	-
	M3	120±13	121±11	-	-
	M4	115±12	123±7	-	-
bDBP (mmHg)			($p < 0.01$; $\eta^2 = 0.29$)	($p = 0.82$; $\eta^2 = 0.00$)	($p = 0.28$; $\eta^2 = 0.09$)
	M1	76±4	80±12	M1 > M4	-
	M2	80±5	77±12	M2 > M4	-
	M3	76±8	74±7	-	-
	M4	71±8	72±5	-	-
bMAP (mmHg)			($p < 0.01$; $\eta^2 = 0.29$)	($p = 0.80$; $\eta^2 = 0.00$)	($p = 0.45$; $\eta^2 = 0.06$)
	M1	92±5	94±13	M1 > M4	-
	M2	91±7	92±13	M2 > M4	-
	M3	91±8	89±7	-	-
	M4	85±9	89±5	-	-
bPP (mmHg)			($p = 0.87$; $\eta^2 = 0.02$)	($p = 0.94$; $\eta^2 = 0.00$)	($p = 0.09$; $\eta^2 = 0.15$)
	M1	49±13	42±14	-	-
	M2	46±9	47±13	-	-
	M3	44±11	46±11	-	-
	M4	45±8	50±7	-	-
cSBP (mmHg)			($p = 0.36$; $\eta^2 = 0.08$)	($p = 0.39$; $\eta^2 = 0.05$)	($p = 0.04$; $\eta^2 = 0.18$)
	M1	140±30	118±20	-	SIT M1 > SIT M2
	M2	123±21	123±15	-	-
	M3	131±30	121±12	-	-
	M4	124±32	120±8	-	-

Data are presented as mean ± standard deviation. Abbreviations: bSBP = brachial systolic blood pressure; bDBP = brachial diastolic blood pressure; bMAP = brachial mean arterial pressure; bPP = brachial pulse pressure; cSBP = central systolic blood pressure; SIT = sprint interval training; CAT = continuous aerobic exercise training; M = moment.

suggesting a decrease from M2 to M4 ($d = -2.0$, 95% CI: -3.0 to -0.2%, $p = 0.01$), independently of age and sex (Supplement 4)

Discussion

The study found that home and gym-based exercise are equally effective at reducing central arterial stiffness, but only gym-based exercise led to improvements in peripheral arterial stiffness, cardiorespiratory fitness, bDBP, and fat mass. Both multicomponent intensity training regimes increased cardiorespiratory fitness and reduced central and peripheral arterial stiffness, but only sprint interval training reduced cSBP. Thus, while the home-based intervention may have mitigated COVID-19's harmful effects on arterial health, gym-based exercise provided greater cardiovascular benefits for people with intellectual and developmental disabilities, regardless of the multicomponent intensity training regime.

Cardiorespiratory fitness

Aerobic training improves cardiorespiratory fitness in adults with intellectual and developmental disabilities (Boer et al., 2014; Boer and Moss, 2016; Melo et al., 2021). High-intensity aerobic exercise is more effective in improving cardiorespiratory fitness than low-intensity exercise when the volume is controlled (Martin-Smith et al., 2020). Sprint interval training yielded greater gain magnitude compared to continuous aerobic training in adolescents and young adults with intellectual and developmental disabilities (Boer et al., 2014) and adding 3 months of HIIT to 12 months of continuous aerobic training reduced the cardiorespiratory fitness non-response rate from 60% to 20% in adults with intellectual and developmental disease (Melo et al., 2021). In the present study, sprint interval training and continuous aerobic training increased peak VO_2 comparably, consistent with previous reports (Su et al., 2019).

Our home-based intervention did not mitigate the negative impact of the 8-week mandatory home confinement on cardiorespiratory fitness. Peak VO_2 typically experiences a 7-15% reduction after 2 weeks of bed rest (Pišot et al., 2016; Schwendinger and Pocecco, 2020). Our study observed similar decreases in peak VO_2 over the home-confinement period, potentially increasing the risk of cardiovascular disease and mortality (Kodama et al., 2009). Gym-based exercise resulted in an 8% increase in peak VO_2 , possibly reducing all-cause mortality risk (Lee et al., 2011). High cardiorespiratory fitness is associated with lower risk of severe COVID-19 and related complications such as hospitalization, intensive care, and death (Af Geijerstam et al., 2021; Christensen et al., 2021; Ekblom-Bak et al., 2021; Ritter and Karargas, 2021). Fat mass reductions noted only after gym-based training may have contributed to increases in cardiorespiratory fitness, suggesting that gym-based training induces greater energy expenditure compared to home-based training.

Blood pressure and arterial stiffness

The effects of aerobic training on blood pressure (BP) and arterial stiffness in people with and without intellectual and developmental disabilities depend on exercise intensity and duration, with larger reductions observed in those with higher BP and stiffer arteries (Ashor et al., 2014; Huang et al., 2016; Zhang et al., 2018). However, the effectiveness of the exercise intervention appears to vary, as both sprint interval training and continuous aerobic training were equally effective in reducing regional stiffness and DBP, as reported in other studies (Hasegawa et al., 2018; Way et al., 2019). The discrepancies in findings may relate to differences in exercise intervention duration and intensity between studies. Both home- and gym-based exercise modes were effective in reducing central arterial stiffness, but decreases in peripheral arterial stiffness were only evident after the gym-based intervention. This suggests that remote multicomponent intensity training regimes using body weight, even without objective control of the intensity, are effective at controlling traditional cardiovascular risk factors and delivering vascular benefits. The findings also agree with those indicating that peripheral arterial stiffness is less amenable to change in short-exercise training interventions (Hayashi et al., 2005). Changes in central and peripheral arterial stiffness were associated with changes in MAP, confirming high BP as the main determinant of stiffer arteries through the promotion of matrix remodeling (Safar et al., 2018). However, the gym-based intervention led to a decrease in bDBP, suggesting that other mechanisms contribute to reductions in central arterial stiffness, including reductions in oxidative stress. Importantly, this exercise-dependent diastolic antihypertensive benefit is clinically relevant (> 5 mmHg), but the home-based intervention did not produce the same benefit. Nonetheless, the home-based intervention,

particularly sprint interval training, reduced cSBP, an important predictor of cardiovascular disease (Lamarche et al., 2021).

Limitations

This study is not without limitations. Our statistical inferences may be underpowered because we only sampled half of the intended participants due to COVID-19 constraints. However, we would like to emphasize that our conclusions are supported by large effect sizes observed, as reported for the main effects of time and multicomponent intensity training regimes -by-time interaction effects across various parameters. Still, readers should interpret these findings as preliminary. We used a cycle ergometer to measure cardiorespiratory fitness, which may not translate to other activities commonly prescribed to this population. Exercise testing validity and reliability, particularly with cycle ergometry, have been questioned in this population, but most CPETs in sprint interval training were maximal. However, only six CPETs were maximal in the continuous aerobic training. We could not objectively monitor exercise intensity during the home intervention because we did not have the means or authorization to deliver heart rate monitors to participants at home during the lockdown. Participants in the sprint interval training group did not achieve the target intensity in both delivery modes. The training regimes were not isocaloric, making effective comparisons challenging. We did not control dietary intake, which likely increased during the lockdown and could have affected study outcomes. Finally, our study included persons with intellectual and developmental disabilities who were familiar with exercise training, which limits the generalization of our findings to the sedentary population.

Practical implications

The study's practical implications hold substantial significance for practitioners and policymakers directing their focus toward cardiovascular health in people with intellectual and developmental disabilities. The findings underscore the critical importance of tailoring exercise interventions to individual needs, with a noteworthy emphasis on the pivotal influence of the delivery mode over the multicomponent intensity training regimes. Gym-based exercises exhibited superior outcomes, affirming their effectiveness. However, the study also accentuates the value of adaptable home-based interventions, particularly in mitigating challenges such as central arterial stiffness during periods like lockdowns. These insights advocate for a balanced approach, urging practitioners to consider both home and gym-based strategies for the development of personalized and inclusive exercise programs, irrespective of the multicomponent intensity training regime. Policymakers can leverage this information to shape inclusive fitness policies, ensuring access to suitable facilities and fostering awareness of cardiovascular health within this population.

Conclusion

A home-based exercise intervention minimized the negative physiological impact of a mandatory lockdown on central arterial stiffness and blood pressure but does not match the benefits in cardiorespiratory fitness, peripheral arterial stiffness, and fat mass of an iso-temporal gym-based intervention, irrespective of the multicomponent intensity training regime.

Acknowledgements

The authors would like to extend their gratitude and acknowledgments to all study participants and their families for their time and energy spent on this program. We would also like to thank JP and AS for their logistical and personal contributions to the successful completion of this program.

Declaration of conflicting interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

Ethical statement

Ethical approval

Ethical Committee of Faculdade de Motricidade Humana – Universidade de Lisboa (4/6/2021).

Disclosures

HSC, SP, and AL contributed to the conception and design, critically reviewed the manuscript, and gave final approval. XM, BS, CC, and IO contributed to the acquisition, analysis, interpretation, drafted and revised the manuscript, and gave final approval. BF, GO, and JLM contributed to the interpretation, critically reviewed the manuscript, and gave final approval.

Registry

ClinicalTrials.gov NCT05701943.

ORCID iD

Xavier Melo  <https://orcid.org/0000-0003-3934-3986>

Data availability statement

The data set associated with this manuscript is deposited here: 10.17632/93w6dmhptd.1

Supplemental Material

Supplemental material for this article is available online.

References

- Af Geijerstam A, Mehlig K, Börjesson M, et al. (2021) Fitness, strength and severity of COVID-19: A prospective register study of 1 559 187 Swedish conscripts. *BMJ Open* 11(7). BMJ Publishing Group.
- American Psychiatric Association (2022) Diagnostic and Statistical Manual of Mental Disorders. In *Diagnostic and Statistical Manual of Mental Disorders*. American Psychiatric Association Publishing. Epub ahead of print 18 March 2022. DOI: [10.1176/APPLBOOKS.9780890425787](https://doi.org/10.1176/APPLBOOKS.9780890425787).
- Ashor AW, Lara J, Siervo M, et al. (2014) Effects of exercise modalities on arterial stiffness and wave reflection: A systematic review and meta-analysis of randomized controlled trials. *PLoS ONE* 9(10): 42. Public Library of Science.

- Azarbal B, Hayes SW, Lewin HC, et al. (2004) The incremental prognostic value of percentage of heart rate reserve achieved over myocardial perfusion single-photon emission computed tomography in the prediction of cardiac death and all-cause mortality. *Journal of the American College of Cardiology* 44(2): 423–430.
- Boer PH and Moss SJ (2016) Effect of continuous aerobic vs. interval training on selected anthropometrical, physiological and functional parameters of adults with Down syndrome. *Journal of Intellectual Disability Research* 60(4). Blackwell Publishing Ltd: 322–334.
- Boer PH, Meeus M, Terblanche E, et al. (2014) The influence of sprint interval training on body composition, physical and metabolic fitness in adolescents and young adults with intellectual disability: A randomized controlled trial. *Clinical Rehabilitation* 28(3): 221–231.
- Bohn L, Ramoa A, Silva G, et al. (2017) Sedentary Behavior and Arterial Stiffness in Adults with and without Metabolic Syndrome. *International Journal of Sports Medicine* 38(5). Georg Thieme Verlag: 396–401.
- Borrega-Mouquinho Y, Sánchez-Gómez J, Fuentes-García JP, et al. (2021) Effects of High-Intensity Interval Training and Moderate-Intensity Training on Stress, Depression, Anxiety, and Resilience in Healthy Adults During Coronavirus Disease 2019 Confinement: A Randomized Controlled Trial. *Frontiers in Psychology* 12, Frontiers Media S.A.
- Brubaker PH and Kitzman DW (2011) Chronotropic Incompetence. *Circulation* 123(9): 1010–1020.
- Calders P, Elmahgoub S, de Mettelinge TR, et al. (2011) Effect of combined exercise training on physical and metabolic fitness in adults with intellectual disability: A controlled trial. *Clinical Rehabilitation* 25(12): 1097–1108.
- Casey DE, Thomas RJ, Bhalla V, et al. (2019) 2019 AHA/ACC Clinical Performance and Quality Measures for Adults With High Blood Pressure: A Report of the American College of Cardiology/American Heart Association Task Force on Performance Measures. *Circulation. Cardiovascular quality and outcomes* 12(11). Circ Cardiovasc Qual Outcomes.
- Cavalcante JL, Lima JAC, Redheuil A, et al. (2011) Aortic stiffness: Current understanding and future directions. *Journal of the American College of Cardiology*. 57: 1511.
- Christensen RAG, Arneja J, Cyr KS, et al. (2021) The association of estimated cardiorespiratory fitness with COVID-19 incidence and mortality: A cohort study. *PLoS ONE* 16(5 May). Public Library of Science.
- Cohen J (1988) *Statistical Power Analysis for the Behavioral Sciences*. New York: Routledge Academic.
- Cole CR, Blackstone EH, Pashkow FJ, et al. (1999) Heart-Rate Recovery Immediately after Exercise as a Predictor of Mortality. *New England Journal of Medicine* 341(18): 1351–1357.
- Ekblom-Bak E, Väisänen D, Ekblom B, et al. (2021) Cardiorespiratory fitness and lifestyle on severe COVID-19 risk in 279,455 adults: a case control study. *International Journal of Behavioral Nutrition and Physical Activity* 18(1). BioMed Central Ltd.
- Emerson E and Hatton C (2008) Socioeconomic disadvantage, social participation and networks and the self-rated health of English men and women with mild and moderate intellectual disabilities: Cross sectional survey. *European Journal of Public Health* 18(1): 31–37.
- Fernhall B, Mccubbin Jeffrey A, Pitetti Kenneth H, et al. (2001) Prediction of maximal heart rate in individuals with mental retardation. *Med Sci Sports Exerc*. Available at: <https://www.acsm-msse.org>
- Fitzgerald H, Stride A and Drury S (2022) COVID-19, lockdown and (disability) sport. *Managing Sport and Leisure* 27(1–2). Routledge: 26–33.
- Fjellstrom S, Hansen E, Hölttä J, et al. (2022) Web-based training intervention to increase physical activity level and improve health for adults with intellectual disability. *Journal of intellectual disability research : JIDRJ Intellect Disabil Res*. 66(12): 967–977.
- Gawlik K, Zwierzchowska A, Manowska B, et al. (2017) Aerobic capacity of adults with intellectual disabilities. *Annals of Agricultural and Environmental Medicine* 24(1). Institute of Agricultural Medicine: 117–120.

- Gearhart RF, Lagally KM, Riechman SE, et al. (2009) STRENGTH TRACKING USING THE OMNI RESISTANCE EXERCISE SCALE IN OLDER MEN AND WOMEN. *Journal of Strength and Conditioning Research* 23(3): 1011–1015.
- Germano-Soares AH, Andrade-Lima A, Meneses AL, et al. (2018) Association of time spent in physical activities and sedentary behaviors with carotid-femoral pulse wave velocity: A systematic review and meta-analysis. *Atherosclerosis* 269. Elsevier: 211–218.
- Haff G and Triplett NT, National Strength & Conditioning Association (U.S.) (2016) *Essentials of Strength Training and Conditioning*. 4 ed. Champaign, IL: Human Kinetics.
- Halekoh U, Højsgaard S and Yan J (2006) The R Package *geepack* for Generalized Estimating Equations. In *JSS Journal of Statistical Software*. Available at: <https://www.jstatsoft.org/>
- Hansen TW, Staessen JA, Torp-Pedersen C, et al. (2006) Prognostic value of aortic pulse wave velocity as index of arterial stiffness in the general population. *Circulation* 113(5): 664–670.
- Hasegawa N, Fujie S, Horii N, et al. (2018) Effects of Different Exercise Modes on Arterial Stiffness and Nitric Oxide Synthesis. *Medicine and Science in Sports and Exercise* 50(6). Lippincott Williams and Wilkins: 1177–1185.
- Hayashi K, Sugawara J, Komine H, et al. (2005) Effects of aerobic exercise training on the stiffness of central and peripheral arteries in middle-aged sedentary men. *Japanese Journal of Physiology* 55(4): 235–239.
- Hilgenkamp TIM and Baynard T (2018) Do individuals with intellectual disability have a lower peak heart rate and maximal oxygen uptake? *Journal of Applied Research in Intellectual Disabilities* 31(5). Blackwell Publishing Ltd: 785–791.
- Hsieh K, Rimmer JH and Heller T (2014) Obesity and associated factors in adults with intellectual disability. *Journal of Intellectual Disability Research* 58(9). Blackwell Publishing Ltd: 851–863.
- Huang C, Wang J, Deng S, et al. (2016) The effects of aerobic endurance exercise on pulse wave velocity and intima media thickness in adults: A systematic review and meta-analysis. In *Scandinavian Journal of Medicine and Science in Sports*. Blackwell Munksgaard.
- Jacinto M, Vitorino AS, Palmeira D, et al. (2021) Perceived Barriers of Physical Activity Participation in Individuals with Intellectual Disability—A Systematic Review. *Healthcare* 9(11). Multidisciplinary Digital Publishing Institute (MDPI).
- Kim S-S (2017) Effects of endurance exercise and half-bath on body composition, cardiorespiratory system, and arterial pulse wave velocity in men with intellectual disabilities. *The Journal of Physical Therapy Science* 29: 1216–1218.
- Kirby T (2020) Efforts escalate to protect homeless people from COVID-19 in UK. *The Lancet Respiratory Medicine* 8(5). Lancet Publishing Group: 447–449.
- Kodama S, Saito K, Tanaka S, et al. (2009) Cardiorespiratory Fitness as a Quantitative Predictor of All-Cause Mortality and Cardiovascular Events in Healthy Men and Women A Meta-analysis. *JAMA* 301: 2024–2035.
- Kuznetsova A, Brockhoff PB and Christensen RHB (2017) **lmerTest** Package: Tests in Linear Mixed Effects Models. *Journal of Statistical Software* 82(13).
- Lamarque F, Agharazii M, Madore F, et al. (2021) Prediction of Cardiovascular Events by Type I Central Systolic Blood Pressure: A Prospective Study. *Hypertension* 77(2). Lippincott Williams and Wilkins: 319–327.
- Lee D-C, Sui X, Artero EG, et al. (2011) Epidemiology and Prevention Long-Term Effects of Changes in Cardiorespiratory Fitness and Body Mass Index on All-Cause and Cardiovascular Disease Mortality in Men The Aerobics Center Longitudinal Study. *Circulation* 124(23): 2483.
- Lenth R (2020) *emmeans: Estimated Marginal Means, aka Least-Squares Means*. R package version 0.9.6.
- Lüdecke D, Makowski D, Waggoner P, et al. (2020b) *Assessment of Regression Models Performance*. CRAN R package.

- Lüdecke D, Ben-Shachar M, Patil I, et al. (2020a) Extracting, Computing and Exploring the Parameters of Statistical Models using R. *Journal of Open Source Software. The Open Journal*. 5(53): 2445.
- Martin-Smith R, Cox A, Buchan DS, et al. (2020) High intensity interval training (HIIT) improves cardio-respiratory fitness (CRF) in healthy, overweight and obese adolescents: A systematic review and meta-analysis of controlled studies. *International Journal of Environmental Research and Public Health*. MDPI AG.
- Melo X, Pinto R, Angarten V, et al. (2021) Training responsiveness of cardiorespiratory fitness and arterial stiffness following moderate-intensity continuous training and high-intensity interval training in adults with intellectual and developmental disabilities. *Journal of Intellectual Disability Research*. Epub ahead of print 28 October 2021. DOI: [10.1111/jir.12894](https://doi.org/10.1111/jir.12894).
- Mendonca G V., Pereira FD and Fernhall B (2011) Effects of combined aerobic and resistance exercise training in adults with and without Down syndrome. *Archives of physical medicine and rehabilitation. Arch Phys Med Rehabil*. 92(1): 37–45.
- Meyer T, Lucía A, Earnest CP, et al. (2005) A Conceptual Framework for Performance Diagnosis and Training Prescription from Submaximal Gas Exchange Parameters ± Theory and Application. *Int J Sports Med* 26(1).
- Naqvi TZ and Lee MS (2014) Carotid intima-media thickness and plaque in cardiovascular risk assessment. In *JACC: Cardiovascular Imaging*. Elsevier Inc.
- Oviedo GR, Guerra-Balic M, Baynard T, et al. (2014) Effects of aerobic, resistance and balance training in adults with intellectual disabilities. *Research in Developmental Disabilities* 35(11). Elsevier Inc: 2624–2634.
- Oviedo GR, Javierre C, Font-Farré M, et al. (2020) Intellectual disability, exercise and aging: The IDEA study: Study protocol for a randomized controlled trial. *BMC Public Health* 20(1). BioMed Central.
- Pescatello LS, Arena R, Riebe D, et al. (2014) *ACSM'S Guidelines for Exercise Testing and Prescription*. Wolters Kluwer/Lippincott Williams & Wilkins Health. Epub ahead of print 2014.
- Pišot R, Marusic U, Biolo G, et al. (2016) Greater loss in muscle mass and function but smaller metabolic alterations in older compared with younger men following 2 wk of bed rest and recovery. *Journal of Applied Physiology* 120(8). American Physiological Society: 922–929.
- Qiao J (2020) What are the risks of COVID-19 infection in pregnant women? *The Lancet* 395.
- Riebe D, Ep-c A, Ehrman JK, et al. (2018) *ACSM's guidelines for exercise testing and prescription*. 10th ed. Philadelphia.
- Ritter O and Kararigas G (2021) Reply — COVID-19, the Female Immune Advantage, and Cardiovascular Impact. In *Mayo Clinic Proceedings*. Elsevier Ltd.
- Rodrigues AN, Cesar Coelho L, Goncalves WLS, et al. (2011) Stiffness of the large arteries in individuals with and without Down syndrome. *Vascular Health and Risk Management* 7(1). DOVE Medical Press Ltd: 375–381.
- Safar ME, Asmar R, Benetos A, et al. (2018) Interaction between hypertension and arterial stiffness an expert reappraisal. In *Hypertension*. Lippincott Williams and Wilkins.
- Salvi P (2016) *Pulse Waves: How Vascular Hemodynamics Affects Blood Pressure*. 2nd ed. Heidelberg, Germany: Springer International Publishing.
- Salvi P, Scalise F, Rovina M, et al. (2019) Noninvasive Estimation of Aortic Stiffness Through Different Approaches. *Hypertension* 74(1). Lippincott Williams & Wilkins Hagerstown, MD : 117–129.
- Schwendinger F and Pocecco E (2020) Counteracting physical inactivity during the COVID-19 pandemic: Evidence-based recommendations for home-based exercise. *International Journal of Environmental Research and Public Health* 17(11). MDPI AG.
- Smith J (2021) *Q&A: Future pandemics are inevitable, but we can reduce the risk*.
- Stanish HI and Aucoin M (2007) Usefulness of a Perceived Exertion Scale for Monitoring Exercise Intensity in Adults with Intellectual Disabilities. In *Education and Training in Developmental Disabilities*.

- Su LQ, Fu JM, Sun SL, et al. (2019) Effects of HIIT and MICT on cardiovascular risk factors in adults with overweight and/or obesity: A meta-analysis. *Plos One* 14(1).
- The Reference Values for Arterial Stiffness' Collaboration (2010a) Determinants of pulse wave velocity in healthy people and in the presence of cardiovascular risk factors: 'establishing normal and reference values'. *European Heart Journal*. Online.
- Theis N, Campbell N, De Leeuw J, et al. (2021) The effects of COVID-19 restrictions on physical activity and mental health of children and young adults with physical and/or intellectual disabilities. *Disability and Health Journal* 14(3). Elsevier: 101064.
- Townsend RR, Wilkinson IB, Schiffrin EL, et al. (2015) Recommendations for Improving and Standardizing Vascular Research on Arterial Stiffness: A Scientific Statement from the American Heart Association. *Hypertension* 66(3). Lippincott Williams and Wilkins: 698–722.
- Way KL, Sultana RN, Sabag A, et al. (2019) The effect of high Intensity interval training versus moderate intensity continuous training on arterial stiffness and 24 h blood pressure responses: A systematic review and meta-analysis. *Journal of Science and Medicine in Sport*. Elsevier Ltd. 22: 385.
- WHO (2020) *Mental health and psychosocial considerations during the COVID-19 outbreak*.
- Zbinden-Foncea H, Francaux M, Deldicque L, et al. (2020) *Does High Cardiorespiratory Fitness Confer Some Protection Against Proinflammatory Responses After Infection by SARS-CoV-2? Obesity*. Blackwell Publishing Inc.
- Zhang Y, Qi L, Xu L, et al. (2018) Effects of exercise modalities on central hemodynamics, arterial stiffness and cardiac function in cardiovascular disease: Systematic review and meta-analysis of randomized controlled trials. *PLoS ONE* 13(7). Public Library of Science.