

Identification of Drug Related Problems in a sample of Portuguese nursing homes

Silvestre, L., Cavaco-Silva, P., Oliveira, P., Carneiro, C., Fernandes, A.I., Costa, F.A.

CiiEM, Instituto Superior de Ciências da Saúde Egas Moniz (ISCSEM), Portugal

Campus Universitário, Quinta da Granja. Monte de Caparica. 2829 - 511 Caparica (alvesdacosta.f@gmail.com)


Introduction

The numerous comorbidities associated with ageing often lead to treatment with multiple drugs. Pharmacist-led medication review of polypharmacy patients may play an important role in the detection of Drug Related Problems (DRP), thus contributing to their prevention or solution.

Objectives

To determine the prevalence of real and potential DRP in polypharmacy elderly residing in nursing homes; and to describe the nature of DRP; and to explore acceptance of pharmacists' reports.

Methods



II Granada Consensus ⁽¹⁾:

Necessity	DRP 1: untreated condition DRP 2: unnecessary medicine
Effectiveness	DRP 3: non-quantitative ineffectiveness DRP 4: quantitative ineffectiveness
Safety	DRP 5: non-quantitative unsafety DRP 6: quantitative unsafety


Statistical Analysis: Univariate. Bivariate: Mann-Whitney, Student's T-Test; Pearson's correlation coefficient. 95% CI.

References


Results

Table 1 presents the sample main characteristics at baseline. The number of medicines directly correlated with the number of DRP detected (Fig 1). Flowchart 1 summarises interventions made and results obtained.

Table 1	Total	IG	CG	p
Age	126	85, 16 (6, 302)	84, 46 (5, 978)	0, 525
Medicines	1332	10, 78 (4, 556)	10, 37 (3, 858)	0, 584
DRP	2109	13, 50 (9, 19)	16 (7, 510)	0, 252



Flowchart 1



Discussion/Conclusions

Pharmacist-led medication review proved useful in identifying DRP in elderly polypharmacy patients. The acceptance rate of pharmacist's reports is far from ideal but may be justified by this project's unexpected impact on media coverage, which led to nursing homes being less receptive to admit flaws in their system. Meetings will be held with the facility that withheld information with the purpose of developing stronger bonds between healthcare providers involved.