



Review

Surgical site infection prevention care bundles in colorectal surgery: a scoping review

T. Cunha^{a,b,*}, S. Miguel^c, J. Maciel^{d,e}, C. Zagalo^{f,g}, P. Alves^h

^a Faculty of Health Sciences and Nursing, Universidade Católica Portuguesa, Lisbon, Portugal

^b UL-PPCIRA, Instituto Português de Oncologia Francisco Gentil, Lisbon, Portugal

^c Centre for Interdisciplinary Research in Health, Faculty of Health Sciences and Nursing, School of Nursing, Universidade Católica Portuguesa, Lisbon, Portugal

^d Faculdade de Medicina da Universidade de Lisboa, Lisbon, Portugal

^e Department of General Surgery, Instituto Português de Oncologia de Lisboa Francisco Gentil, Lisbon, Portugal

^f Clinical Research Unit, Centro de Investigação Interdisciplinar Egas Moniz, Egas Moniz – Cooperativa de Ensino Superior, Companhia de Responsabilidade Limitada, Almada, Portugal

^g Department of Head and Neck Surgery, Instituto Português de Oncologia de Lisboa Francisco Gentil, Lisbon, Portugal

^h Centre for Interdisciplinary Research in Health, Faculty of Health Sciences and Nursing, School of Nursing, Universidade Católica Portuguesa, Porto, Portugal

ARTICLE INFO

Article history:

Received 25 January 2024

Accepted 22 October 2024

Available online 31 October 2024

Keywords:

Surgical site infections

Patient care bundles

Colorectal surgery

Scoping review



SUMMARY

Background: Surgical site infection (SSI) prevention bundles have been used to reduce infection rates in most types of surgery. Bundles tailored to colorectal surgery have been used with success.

Aim: To identify and review the individual interventions that constitute each SSI prevention care bundle in colorectal surgery, and the reduction in SSI rate associated with their implementation.

Methods: A scoping review was conducted in PubMed, CINAHL, Web of Science Core Collection and Scopus in December 2022.

Results: This review analysed 48 of 164 identified studies on SSI prevention in colorectal surgery from 2011 to 2022. It revealed an average of 11 interventions per study, primarily in the pre-operative [mechanical bowel preparation, oral antibiotic bowel decontamination, hair removal, chlorhexidine gluconate (CHG) shower, normoglycaemia], intra-operative (antibiotic prophylaxis, normothermia, CHG skin preparation, antibiotic prophylaxis re-dosing, gown/glove change) and postoperative (normothermia, normoglycaemia, dressing removal, oxygen optimization, incision cleansing) periods. Despite these interventions, SSI rates remain high, indicating a need for further research to optimize intervention bundles and improve compliance across surgical stages.

* Corresponding author. Address: Faculty of Health Sciences and Nursing, Universidade Católica Portuguesa, Palma de Cima, 1649-023 Lisboa, Portugal.

E-mail address: tiagocunha@sapo.pt (T. Cunha).

Conclusions: The implementation of SSI prevention bundles, tailored to colorectal surgery, has shown a reduction in SSI rates and costs. Grouping interventions according to the peri-operative phase may increase compliance.

© 2024 The Authors. Published by Elsevier Ltd on behalf of The Healthcare Infection Society. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Introduction

Surgical site infections (SSIs) are among the most common healthcare-associated infections (HAIs) worldwide, accounting for more than 30% of all HAIs, and lead to increased morbidity, length of stay, readmission rates, and healthcare costs [1–3]. SSIs can cost up to \$3.2 billion each year, and place a burden on healthcare systems, limiting their response and increasing surgical waiting lists [4]. In 2014, the average cost of all SSIs was found to be £5239 in an UK hospital [2]. These costs are not only financial; SSIs also have a negative effect on patients' health-related quality of life, and occupational and societal commitments [2].

Colorectal cancer affects more than 1.9 million patients each year, causing 0.94 million deaths per year [5]. This accounts for 10% of all cases of cancer globally and 9.4% of cancer deaths, ranking it the third most common cancer in recent years, and the second leading cause of cancer death globally [5]. Colorectal surgery is paramount for the treatment of these tumours [5]. However, SSI rates in colorectal surgery are disproportionately high, ranging from 15% to 30%, versus 2.5% in all surgical procedures [1,3,6]. However, when compared with other clean-contaminated surgeries, such as gynaecological surgery (2–9.2%) [7–9] or head and neck surgery (6.6–41.8%) [10], the SSI rates are comparable. This may be attributed to the risk of wound contamination with bowel content [2].

In Europe, SSI rates in colorectal surgery varied from 2.9% to 12.5% for laparoscopic colon surgery, and from 4.1% to 16.9% for open colon surgery in 2017 [11]. For the period between 2018 and 2020, these rates increased to 3.5–16.4% and 3.6–26.8%, respectively [12]. These are the highest SSI rates amongst all types of surgical procedure surveyed in the European Centre for Disease Prevention and Control's HAI-Net SSI programme, and show a rising tendency [11,12].

The increased costs associated with colorectal SSI range from \$36.429 to \$144.809 per patient in the USA, according to a study which claimed that the true economic burden is under-recognized in most studies. [4]. In the UK, as colorectal surgery is a common elective procedure, the disproportionately high SSI rate leads to a substantial economic burden, with an associated 35% increase in direct healthcare costs [6]. As such, a reduction in the SSI rate for colorectal surgery has become a major objective of quality improvement initiatives [1]. In order to prevent SSIs, patient care bundles have been developed and implemented with some success. Different systematic reviews and meta-analyses have shown significant reductions in the risk of SSIs in patients who received an infection prevention care bundle compared with patients who received standard care [13,14]. A risk reduction of 40% ($P<0.001$) was recorded in one study [14].

Despite the implementation of SSI prevention bundles, the decrease in SSIs in colorectal surgery has been slow [12]. Some

centres have developed and implemented patient care bundles tailored to prevent SSIs in colorectal surgery [15].

The primary aim of this study was to identify and review the individual interventions that constitute each infection prevention care bundle in colorectal surgery. The secondary aims of this study were: to identify the peri-operative period in which each intervention is performed; and to identify the SSI reduction rate related to implementation of each care bundle.

Methods

Protocol and registration

This study was based on the guidance framework for conducting scoping reviews developed by the Joanna Briggs Institute [16]. It has been reported using the Preferred Reporting Items for Systematic Reviews and Meta-Analyses Extension for Scoping Reviews [17,18], and is registered on the OSF Database (<https://doi.org/10.17605/OSF.IO/64VEY>).

Eligibility criteria and search strategy

The selection of studies for this scoping review was based on the question, 'What individual interventions constitute each infection prevention care bundle in colorectal surgery?' Studies were eligible for inclusion in this scoping review if they met all of the following criteria: (i) original publication; (ii) reported on SSI prevention care bundles in patients undergoing colorectal surgery; and (iii) included adult patients alone.

The following exclusion criteria were selected: population – all studies without patients undergoing colorectal surgery or patients aged <18 years were excluded; study design – all revisions were excluded; intervention – all studies which only mentioned isolated interventions that were not part of a care bundle were excluded; and outcome – all studies that analysed interventions in the context of guidelines and not infection prevention care bundles were excluded.

The literature search was conducted in December 2022 using Web of Science Core Collection, PubMed, CINAHL and Scopus. It employed a comprehensive strategy using Medical Subject Heading terms and natural language keywords to ensure a thorough search of relevant articles. The search terms, which are outlined in [Supplementary File 1](#), were applied to the titles and abstracts of publications in the database. The detailed process of study selection, including the number of articles screened, assessed for eligibility and included in the review, is depicted in the PRISMA flow diagram ([Figure 1](#)). This approach was developed to capture the broadest possible range of relevant research.




Figure 1. PRISMA flow diagram.

Extraction of data and charting

Two independent authors (TC and SM) extracted and charted data from the included studies, removed duplicates using Rayyan software, and applied the eligibility criteria, with blind screening of titles and abstracts. All disagreements passed to the next step, and full-text screening was undertaken independently by the same reviewers (TC and SM). A third reviewer (PA) was available to resolve disagreements when necessary, through discussion with the other authors. The following information was tabulated: author; country; year of publication; aims; methodology; participants; healthcare context; interventions SSI bundles; and study design. After that, interventions were grouped by period: pre-operative, intra-operative and postoperative.

Results

In total, 164 papers were retrieved. Following the removal of duplicates, the titles and abstracts of 114 papers were screened, and the full-text of 55 papers was assessed for eligibility. Forty-eight papers met the inclusion criteria as shown in the PRISMA flow diagram [17] (Figure 1).

Characteristics of included studies

Forty-eight studies met the inclusion criteria [1,2,4,6,19–62]. The included studies were published between 2011 and 2022. Sixty-four percent (N=30) of the studies were published between 2018 and 2022. The studies were performed in the USA (N=29), UK (N=5), Canada (N=2), Australia (N=2), Italy (N=2), Spain (N=3), Netherlands (N=1), Saudi Arabia (N=1), Switzerland (N=1), South Korea (N=1) and Thailand (N=1). The number of participants in the studies ranged from

86 to 107,665, with an aggregate number of participants of 229,707. In one study, it was not possible to determine the number of participants [44].

The studies included 528 interventions, with a mean of 11 interventions per study, ranging between two and 22 interventions. Details about the 48 included studies can be viewed in [Supplementary File II](#). Regarding the study context, they can be divided into studies conducted at single hospitals (N=33) and studies conducted in multiple hospitals (N=15). In terms of study design, they can be divided into: cohort studies (N=17); retrospective cohort studies (N=13); prospective cohort studies (N=12); quasi-experimental studies (N=3); case-control studies (N=1); pre-post cohort studies (N=1); and a pragmatic study involving three phases (N=1).

A reduction in SSI rate was achieved in 43 studies. However, in two of these studies, the reduction was not significant. In three studies, an increase in SSI rate was witnessed. Two studies made no comparisons between the pre- and post-bundle implementation periods. These results can be found in [Tables I, II and III](#).

The interventions were divided into three phases: pre-operative, intra-operative and postoperative ([Tables I, II and III](#)). In the pre-operative phase, 158 interventions were identified. The mean number of interventions per study was three, ranging between zero and eight interventions. In the intra-operative phase, 272 interventions were identified. The mean number of interventions per study was six, ranging between zero and 12 interventions. Finally, in the postoperative phase, 96 interventions were identified. The mean number of interventions per study was two, ranging between zero and six.

The most common interventions in each phase are shown in [Table IV](#). Some interventions were present in all phases of the peri-operative period, such as normothermia and normoglycaemia.

Table I

Individual interventions and surgical site infection (SSI) rate reductions in the pre-operative period

[illegible]

CHG, chlorhexidine gluconate; ICU, intensive care unit; MRSA, methicillin-resistant *Staphylococcus aureus*; OR, operating room; PACU, postanesthesia care unit.

Discussion

The current scoping review reports findings from 48 studies, published between 2011 and 2022, identified through a systematic literature search. These studies focused on the individual interventions that constitute each infection prevention care bundle in colorectal surgery.

Implementation of SSI prevention bundles

The World Health Organization (WHO) [63] and several other national entities, including the National Institute for Health and Care Excellence (NICE) [64] and the Centers for Disease Control and Prevention (CDC) [65], have designed and implemented SSI prevention bundles. Despite these efforts, SSI rates in colorectal surgery remain high. This persistent issue underscores the complexity and challenges of reducing SSI rates effectively in clinical practice. The relatively high number of articles published in the last 5 years (2018–2022), with 64% ($N=29$) of the studies in this review falling into this period, suggests growing interest in this subject. This increase in research may reflect both a recognition of the ongoing problem, and a commitment to improving patient outcomes in colorectal surgery.

Composition and complexity of care bundles

One of the mechanisms to translate available research evidence into clinical practice is the implementation of care bundles [66]. These bundles are structured sets of evidence-based practices that, when performed collectively and reliably, have been shown to improve patient outcomes. This scoping review found a mean of 11 interventions per study.

which is higher than expected according to care bundle design guidelines. This discrepancy raises questions about the practical implementation of these bundles.

Guidelines for the design of care bundles recommend that they should contain only three to five interventions with strong clinical agreement [66,67]. This recommendation is based on evidence suggesting that simpler bundles with fewer components are more likely to be adhered to by clinical staff, thereby improving their effectiveness. Bundles with a smaller number of elements tend to have higher compliance rates. However, bundles with more than six interventions are not uncommon [66]. Such complex care bundles can pose significant challenges to adherence, particularly for multi-disciplinary teams that must coordinate and implement multiple practices [68].

Effectiveness of bundles with multiple interventions

A recent meta-analysis [69] showed that no increase in effectiveness was found when more interventions were combined. In fact, SSI prevention care bundles with a higher proportion of evidence-based interventions usually presented fewer interventions. This finding aligns with the recommendations by the Institute for Healthcare Improvement, which state that care bundles should consist of three to five evidence-based interventions [15,67]. This highlights the importance of not overcomplicating care bundles, as excessive complexity can hinder their practical application and reduce compliance.

Phase-based division of interventions

In this review, these bundles were divided into three operative phases: pre-operative, intra-operative and postoperative.

Table II

Individual interventions and surgical site infection (SSI) rate reductions in the intra-operative period

Intra-operative period		Dixon et al. [2]	Arroyo-Garcia et al. [19]	Antonacci et al. [20]	Clayman et al. [21]	Schlick et al. [22]	Bradjich et al. [23]	Jurt et al. [24]	Faraglier et al. [25]	Vicentini et al. [26]	Falconer et al. [6]	Zarain-Orador [27]	Loisriwat [28]	D'Andrea et al. [29]	Martinez et al. [30]	Leaper et al. [4]	Park et al. [31]	Hajrawala et al. [32]	Dean et al. [33]	D'Souza et al. [34]	Guzman-Pineda et al. [35]	Magee et al. [36]	Albert et al. [37]	Huang et al. [38]	Russell et al. [39]	Lin et al. [40]	Weiser et al. [42]	Gachabayov et al. [43]	Harris et al. [44]	Gorgun et al. [45]	Jaffe et al. [46]	Klan et al. [47]	Ohman et al. [48]	Bert et al. [49]	Hewitt et al. [50]	Kwaan et al. [51]	Berlitz et al. [52]	Rumberger et al. [53]	Tanner et al. [54]	Keenan et al. [55]	Shuman et al. [56]	Pérez-Blanco et al. [57]	Keenan et al. [1]	Watts et al. [58]	Cima et al. [59]	Crofta et al. [60]	Lutifiya et al. [61]	Bull et al. [62]	Total						
SSI rate reduction		14.9%	8.6%	1.7%	8.5%	n/a	n/a	0.8%	9.5%	6.0%	7.0%	1.4%	6.5%	9.2%	10.6%	n/a	5.7%	14.5%	10.0%	6.4%	11.0%	3.1%	7.2%	6.2%	n/a	2.8%	7.5%	6.9%	12.3%	55.5%	5.2%	7.8%	4.0%	11.5%	3.4%	9.2%	0.8%	5.3%	1.1%	4.0%	1.4%	10.6%	13.6%	15.5%	5.8%	2.8%	14.5%	8.0%	Total						
Intra-operative	Antibiotic prophylaxis (type, timing and discontinuation)																																																						38
	Normothermia																																																						29
	Skin prep with 2% CHG																																																						24
	Antibiotic re-dosing																																																						23
	Gown and glove change before fascial closure																																																						23
	Dedicated closure tray																																																						20
	Wound protector																																																						19
	Normoglycaemia																																																						17
	Intracavity lavage																																																						11
	Oxygen delivery (FiO ₂ ≥80%)																																																						8
	Triclosan-coated sutures																																																						7
	Limited OR traffic																																																						6
	Changing of suction and electrocautery																																																						5
	Laparoscopic surgery																																																						4
	Closure strategy																																																						3
	Sterile occlusive incisional wound dressing placed in OR																																																						3
	Surgical hand hygiene with hydroalcoholic solutions																																																						3
	Hair removal when appropriate with clippers																																																						3
	Altemeier wound classification																																																						2
	Glove change after each intra-operative digital rectal exam																																																						

CHG, chlorhexidine gluconate; CO₂, carbon dioxide; FiO₂, fraction of inspired oxygen; ICU, intensive care unit; OR, operating room; PACU, postanaesthesia care unit.

This division is in line with SSI prevention guidelines from WHO [63], NICE [64] and CDC [65,70]. Such a structured approach can help streamline the implementation process, and improve compliance rates by making the steps more manageable for healthcare providers.

Following this analysis of the different interventions and allocation to the three operative phases by the study authors, some inconsistencies were identified. As such, the present authors decided to group some interventions and reorder them in the operative phase that seemed most suitable. For this endeavour, it was necessary to define each operative phase in time. The pre-operative phase was considered as the period from admission until patient handover to the operating room staff, the intra-operative phase was considered as the period from patient handover to the operating room staff until return of the patient to the surgical ward, and the postoperative phase was considered as the period from return of the patient to the surgical ward until discharge. The outcome of this analysis is shown in Table V.

Compliance and outcomes

Infection prevention care bundles have been proven to be valid tools to reduce SSIs in colorectal surgery. However, it is important to note that three of the selected studies reported an increase in SSI rates. For instance, Jurt *et al.* [24] postulated that the absence of a decrease in SSI rate could be associated with low compliance with individual bundle interventions. Tanner *et al.* [54] found that full compliance with the care bundle was only achieved in 19% of patients, resulting in similar SSI rates between the compliant and non-compliant groups. Rumberger *et al.* [53] reported that, while their SSI rate did not improve after care bundle implementation, other outcomes in colorectal surgery were enhanced.

These findings emphasize the critical role of compliance in the effectiveness of care bundles. Without high levels of adherence to all components of a care bundle, the anticipated benefits may not be fully realized. This highlights the need for ongoing education, training and support for clinical staff to ensure that all aspects of care bundles are implemented consistently.

Table III

Individual interventions and surgical site infection (SSI) rate reductions in the postoperative period

		Postoperative period		Total	
Postoperative	SSI rate reduction	-14.9%			17
	Normoglycaemia	-5.6%			16
	Normothermia	-1.7%			10
	Removal of sterile dressing within 48 h	-8.5%			6
	Oxygen delivery (FIO ₂ ≥80%)	n/a			3
	Washing of incisions with CHG	n/a			3
	Hand hygiene	-3.8%			3
	SSI assessment and documentation	-9.5%			3
	Protect primary closure incisions for 5 days with silver-impregnated or polyhexamethylene biguanide dressing	-6.0%			3
	Urinary catheter removal within postoperative 2 days	-1.4%			2
	Standardized wound care orders	-6.5%			2
	Wound surveillance	-9.2%			2
	Patient education	-10.8%			2
	Patient shower with CHG after dressing removal	n/a			1
	Iodopovidone wound wash	-5.7%			1
	Short operative duration (incision to closure) <100 min	-14.5%			1
	Atteimer wound class documented	-10.8%			1
	PATOS documented	-5.4%			1
	ASA score documented	-11.0%			1
	Standard handoffs between pre-op, OR, PACU and/or ICU	-3.1%			1
	Standardized debriefing at the end of the case	-7.2%			1
	'Enhanced' recovery after surgery interventions	-5.2%			1
	Systolic blood pressure ≥90 mmHg	n/a			1
	Stoma bag applied before dressing if required	-2.8%			1
	Padded tegaderm dressing left in place for 5 days	-7.5%			1
	Update and assessment of surgical antibiotic prophylaxis	-12.3%			1
	Wound care service consultation obtained for all ostomies and complex wounds	-5.5%			1
	Patients asked to ambulate evening of surgery with staff	-5.2%			1
	Start of regular diet in evening of surgery	-7.8%			1
	Patients are maintained on a minimal narcotic regimen	-4.0%			1
	Discharge is upon laxation	-11.5%			1
	Low volume negative pressure to aid in healing	-3.4%			1
	Shower on postoperative day 2	-9.2%			1
Do not leave OR in scrubs	-0.8%			1	
Dressing changes using sterile technique	5.3%			1	
Review of wound by attending surgeon post discharge	1.1%			1	
	-4.0%			3	
	-9.8%			4	
	-1.4%			0	
	-10.6%			2	
	-13.6%			5	
	-15.5%			2	
	-5.8%			2	
	-2.8%			1	
	-14.5%			3	
	-8.0%			3	
	Total	0	5	0	17
		2022	2022	2022	2022
		Dixon et al. [2]	Arroyo-Garcia et al. [19]	Antonacci et al. [20]	Cleyghan et al. [21]
					Schlick et al. [22]
					Braych et al. [23]
					Jurt et al. [24]
					Faragher et al. [25]
					Vicentini et al. [26]
					Falconer et al. [6]
					Zarain-Obrador [27]
					Lohsiriwat [28]
					D'Andrea et al. [29]
					Martinez et al. [30]
					Leaper et al. [4]
					Park et al. [31]
					Hajiravala et al. [32]
					Dean et al. [33]
					D'Souza et al. [34]
					Guzman-Punke et al. [35]
					Mcgee et al. [36]
					Albert et al. [37]
					Hoang et al. [38]
					Russell et al. [39]
					Viv et al. [40]
					Lin et al. [41]
					Weiser et al. [42]
					Gachabayov et al. [43]
					Harris et al. [44]
					Gargun et al. [45]
					Jeffer et al. [46]
					Kilan et al. [47]
					Orman et al. [48]
					Bert et al. [49]
					Hewitt et al. [50]
					Kwaan et al. [51]
					Benlice et al. [52]
					Rumberger et al. [53]
					Tanier et al. [54]
					Keenan et al. [55]
					Keenan et al. [56]
					Perez-Blanco et al. [67]
					Keenan et al. [1]
					Wells et al. [58]
					Cima et al. [59]
					Croila et al. [60]
					Luffiya et al. [61]
					Bull et al. [62]
					2011

ASA, American Society of Anaesthesiologists; CHG, chlorhexidine gluconate; FIO₂, fraction of inspired oxygen; ICU, intensive care unit; OR, operating room; PACU, postanaesthesia care unit; PATOS, (infection) present at time of surgery.

Table IV

Most common interventions in each peri-operative phase

Pre-operative	Intra-operative	Postoperative
Mechanical bowel preparation	Antibiotic prophylaxis (type, timing and discontinuation)	Normothermia
Oral antibiotic bowel decontamination	Normothermia	Normoglycaemia
Hair removal with clippers when appropriate	Skin preparation with 2% CHG alcoholic solution	Removal of sterile dressing within 48 h
Shower with 2% CHG	Antibiotic re-dosing	Oxygen delivery ($\text{FiO}_2 \geq 80\%$)
Normoglycaemia	Gown and glove change before fascial closure	Washing of incisions with 2% CHG alcoholic solution
	Dedicated closure trays	
	Wound protectors	
	Normoglycaemia	

CHG, chlorhexidine gluconate; FiO₂, fraction of inspired oxygen.

Strengths and limitations

This scoping review has several strengths, including a comprehensive search strategy and robust inclusion criteria, ensuring a thorough examination of the literature on SSI prevention

bundles in colorectal surgery. However, there are limitations to this research that should be acknowledged.

Firstly, the search was limited to four databases (PubMed, CINAHL, Web of Science Core Collection and Scopus). While these databases are extensive and widely used, the exclusion

Table V

Distribution of interventions by peri-operative phase

Pre-operative	N		N
• CHG shower/cloths	28	• Checklist to determine patient readiness for OR	1
• Mechanical bowel preparation	26	• SSI risk assessment provided to surgeon	1
• Oral antibiotic bowel decontamination	24	• PATOS documented	1
• Hair removal with clippers when appropriate	24	• MRSA screening and decolonization	1
• Normoglycaemia	16	• Iodophor nares swabs to decolonize the nose for 24 h	1
• Normothermia	10	• Medical handwashing	1
• Patient education	5	• Maintenance of oxygen saturation at least 92%	1
• Clear liquid diets on day prior	3	• Optimizing nutrition	1
• Encourage smoking cessation 30 days before surgery	3	• Oral carbohydrate loading	1
• Standard handoffs between pre-op, OR, PACU and/or ICU	1		
Intra-operative	N		N
• Antibiotic prophylaxis (type, timing and discontinuation) and re-dosing	44	• Absorbable subcuticular suture and skin glue	2
• Skin prep with 2% CHG	31	• Use of warmed CO ₂ during laparoscopic procedures	1
• Normothermia	29	• Elimination of 'flash' steam sterilization of instruments	1
• Drapes, gown and glove change before fascial closure	25	• Standard handoffs between pre-op, OR, PACU and/or ICU	1
• Dedicated closure tray	20	• Short operative duration (<100 min)	1
• Wound protector	19	• Systolic blood pressure ≥90 mmHg	1
• Normoglycaemia	17	• Standardized urinary catheter placement	1
• Intracavity lavage	11	• ASA score documented	1
• Oxygen delivery (FiO ₂ ≥80%)	8	• Double gloving for all scrubbed surgical team members	1
• Triclosan-coated sutures	7	• Pulse lavage of subcutaneous tissues for all open surgery	1
• Limited OR traffic	6	• Antibacterial dressings	1
• Changing of suction and electrocautery	5	• Antiseptic-impregnated incise drapes	1
• Laparoscopic surgery	4	• Florescent imaging instrument to assess perfusion in the anastomosis and to check for microleaks	1
• Closure strategy	3	• Subcutaneous heparin	1
• Sterile occlusive incisional wound dressing placed in OR	3	• Isolation technique	1
• Surgical hand hygiene with hydroalcoholic solutions	3	• Ensuring surgical masks are worn at all times	1
• Altemeier wound classification documented	3	• Sticker with dressing change instructions	1
• Glove change after each intra-operative digital rectal exam	2	• Do not leave OR in scrubs	1
• Sleeve placement and glove change after anastomosis established	2		
Postoperative	N		N
• Normoglycaemia	17	• Iodopovidone wound wash	1
• Normothermia	16	• Low volume negative pressure to aid in healing	1
• Removal of sterile dressing within 48 h	10	• Dressing changes using sterile technique	1
• Oxygen delivery (FiO ₂ ≥80%)	8	• Shower on postoperative day 2	1
• Washing of incisions with CHG	6	• Patients asked to ambulate evening of surgery with staff	1
• Hand hygiene	3	• Start of regular diet in evening of surgery	1
• SSI assessment and documentation	3	• Patients are maintained on a minimal narcotic regimen	1
• Protect primary closure incisions for 5 days with silver-impregnated or polyhexamethylene biguanide dressing	3	• Padded tegaderm dressing left in place for 5 days	1
• Urinary catheter removal within postoperative 2 days	3	• Update and assessment of surgical antibiotic prophylaxis	1
• Standardized wound care orders	2	• Wound care service consultation obtained for all ostomies and complex wounds	1
• Wound surveillance	2	• Standardized debriefing at the end of the case	1
• Patient education	2	• 'Enhanced' recovery after surgery interventions	1
• Standard handoffs between pre-op, OR, PACU and/or ICU	1	• Systolic blood pressure ≥90 mmHg	1
• Review of wound by attending surgeon post discharge	1	• Stoma bag applied before dressing if required	1
• Patient shower with CHG after dressing removal	1	• Discharge is upon laxation	1

ASA, American Society of Anaesthesiologists; CHG, chlorhexidine gluconate; CO₂, carbon dioxide; FiO₂, fraction of inspired oxygen; ICU, intensive care unit; MRSA, methicillin-resistant *Staphylococcus aureus*; OR, operating room; PACU, postanesthesia care unit; PATOS, (infection) present at time of surgery.

of other databases may have resulted in the omission of relevant studies. This limitation could affect the comprehensiveness of the review and potentially bias the findings.

Secondly, the included studies exhibited considerable heterogeneity in terms of population size, study design and geographical distribution. This variability can complicate comparison and synthesis of the results. Moreover, a significant proportion of the studies were conducted in the USA, which may limit the generalizability of the findings to other healthcare settings with different practices and resources.

This review provides valuable insights into the implementation and effectiveness of SSI prevention bundles in colorectal surgery, highlighting areas for future research and potential improvements in clinical practice.

Research on infection prevention care bundles has increased significantly in recent years, reflecting a growing recognition of their potential to improve clinical outcomes in colorectal surgery. Despite this progress, there remains a lack of consensus on the most effective interventions within the pre-operative, intra-operative and postoperative periods. The diversity of approaches underscores the need for standardized protocols to ensure consistency and effectiveness across different healthcare settings.

Implementing SSI prevention bundles is a critical strategy for enhancing the quality of care and improving surgical outcomes. These bundles are designed to reduce SSI rates systematically, thereby lowering associated healthcare costs. This scoping review has identified the most prevalent interventions utilized in each period, providing valuable insights that can inform the design and development of future care bundles tailored specifically for colorectal surgery.

This review demonstrates that while SSI prevention care bundles are generally effective in reducing SSI rates in colorectal surgery, their success is highly dependent on consistent and comprehensive compliance with the prescribed interventions. This finding highlights the importance of adherence to all components of the care bundles to achieve the desired outcomes.

Overall, this scoping review identified the most common interventions used in care bundles across the different periods in colorectal surgery. Further research is essential to optimize the implementation and compliance rates of these interventions. Such efforts will facilitate their wider dissemination and more effective use in clinical practice, ultimately contributing to the reduction of SSI rates in colorectal surgery.

Future research should also aim to determine the optimal composition of care bundles for SSI prevention. This will involve identifying which specific interventions, when combined, yield the greatest effectiveness while maintaining feasibility and ease of implementation in various clinical settings. Establishing such evidence-based standards will be crucial for advancing the efficacy of infection prevention strategies, and improving patient outcomes in colorectal surgery.

Conflict of interest statement

None declared.

Funding sources

This research did not receive any specific grant funding from agencies in the public, commercial or not-for-profit sectors.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.jhin.2024.10.010>.

References

- [1] Keenan JE, Speicher PJ, Thacker JKM, Walter M, Kuchibhatla M, Mantyh CR. The preventive surgical site infection bundle in colorectal surgery: an effective approach to surgical site infection reduction and health care cost savings. *JAMA Surg* 2014;149:1045–52.
- [2] Dixon LK, Biggs S, Messenger D, Shabbir J. Surgical site infection prevention bundle in elective colorectal surgery. *J Hosp Infect* 2022;122:162–7.
- [3] Mengistu DA, Alemu A, Abdulkadir AA, Mohammed Husen A, Ahmed F, Mohammed B, et al. Global incidence of surgical site infection among patients: systematic review and meta-analysis. *Inquiry* 2023;60:469580231162549.
- [4] Leaper DJ, Holy CE, Spencer M, Chitnis A, Hogan A, Wright GWJ, et al. Assessment of the risk and economic burden of surgical site infection following colorectal surgery using a US longitudinal database: is there a role for innovative antimicrobial wound closure technology to reduce the risk of infection? *Dis Colon Rectum* 2020;63:1628–38.
- [5] Xi Y, Xu P. Global colorectal cancer burden in 2020 and projections to 2040. *Transl Oncol* 2021;14:101174.
- [6] Falconer R, Ramsay G, Hudson J, Watson A; Highland Colorectal SSIG. Reducing surgical site infection rates in colorectal surgery – a quality improvement approach to implementing a comprehensive bundle. *Colorectal Dis* 2021;23:2999–3007.
- [7] Steiner HL, Strand EA. Surgical-site infection in gynecologic surgery: pathophysiology and prevention. *Am J Obstet Gynecol* 2017;217:121–8.
- [8] Kulkarni SP, Kothari O. Surgical site infection in obstetric and gynecological surgeries: a prospective observational study. *Cureus* 2023;15:e34855.
- [9] Lachiewicz MP, Moulton LJ, Jaiyeoba O. Pelvic surgical site infections in gynecologic surgery. *Infect Dis Obstet Gynecol* 2015;2015:614950.
- [10] Cunha TFS, Soares Melancia TA, Zagalo Fernandes Ribeiro CM, Almeida de Brito JA, Abreu Miguel SS, André Abreu Esteves Bogalhão do Casal D. Risk factors for surgical site infection in cervicofacial oncological surgery. *J Craniomaxillofac Surg* 2012;40:443–8.
- [11] European Centre for Disease Prevention and Control. Healthcare-associated infections: surgical site infections. Annual epidemiological report for 2017. Stockholm: ECDC; 2019. p. 18. Available at: https://www.ecdc.europa.eu/sites/default/files/documents/AER_for_2017-SSI.pdf.
- [12] European Centre for Disease Prevention and Control. Healthcare-associated infections: surgical site infections. Annual epidemiological report for 2018–2020. Stockholm: ECDC; 2023.
- [13] Tanner J, Padley W, Assadian O, Leaper D, Kiernan M, Edmiston C. Do surgical care bundles reduce the risk of surgical site infections in patients undergoing colorectal surgery? A systematic review and cohort meta-analysis of 8,515 patients. *Surgery* 2015;158:66–77.
- [14] Zywtot A, Lau CSM, Fletcher HS, Paul S. Bundles prevent surgical site infections after colorectal surgery: meta-analysis and systematic review. *J Gastrointest Surg* 2017;21:1915–30.
- [15] Wolfhagen N, Boldingh QJJ, Boermeester MA, de Jonge SW. Perioperative care bundles for the prevention of surgical-site infections: meta-analysis. *Br J Surg* 2022;109:933–42.
- [16] Aromataris E, Lockwood C, Porritt K, Pilla B, Jordan Z, editors. JBI manual for evidence synthesis. JBI; 2024. Available from: <https://synthesismanual.jbi.global>.

- [17] Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ* 2021;372:n71.
- [18] Tricco AC, Lillie E, Zarin W, O'Brien K, Colquhoun H, Kastner M, et al. A scoping review on the conduct and reporting of scoping reviews. *BMC Med Res Methodol* 2016;16:15.
- [19] Arroyo-Garcia N, Badia JM, Vazquez A, Pera M, Pares D, Limon E, et al. An interventional nationwide surveillance program lowers postoperative infection rates in elective colorectal surgery. A cohort study (2008–2019). *Int J Surg* 2022;102:106611.
- [20] Antonacci AC, Antonacci CL, Dechario SP, Husk G, Schilling ME, Cifu-Tursellino K, et al. Reducing surgical site infections after colectomy: bundle item compliance, process, and outlier identification. *Surg Endosc* 2022;36:6049–58.
- [21] Clayphan B, Dixon L, Biggs S, Jordan L, Pullyblank A, Holden K, et al. PreciSSlon: a collaborative initiative to reduce surgical site infections after elective colorectal surgery. *J Hosp Infect* 2022;130:131–7.
- [22] Schlick CJR, Huang RP, Brajcich BC, Halverson AL, Yang AD, Kreutzer L, et al. Unbundling bundles: evaluating the association of individual colorectal surgical site infection reduction bundle elements on infection rates in a statewide collaborative. *Dis Colon Rectum* 2022;65:1052–61.
- [23] Brajcich BC, Schlick CJR, Halverson AL, Huang RP, Yang AD, Love R, et al. Association between patient and hospital characteristics and adherence to a surgical site infection reduction bundle in a statewide surgical quality improvement collaborative. *J Am Coll Surg* 2022;234:783–92.
- [24] Jurt J, Hübner M, Clerc D, Curchod P, Abd El Aziz MA, Hahnloser D, et al. Challenges related to surgical site infection prevention – results after standardized bundle implementation. *J Clin Med* 2021;10:4524.
- [25] Faragher I, Tham N, Hong M, Guy S, Yeung J. Implementation of an organ space infection prevention bundle reduces the rate of organ space infection after elective colorectal surgery. *BMJ Open Qual* 2021;10:e001278.
- [26] Vicentini C, Scacchi A, Corradi A, Marengo N, Furmenti MF, Quattrocchio F, et al. Interrupted time series analysis of the impact of a bundle on surgical site infections after colon surgery. *Am J Infect Control* 2021;49:1024–30.
- [27] Zarain-Obrador L, Alonso-García M, Gil-Yonte P, Hijas-Gómez AI, Rodríguez-Villar D, Martínez-Amores B, et al. Effect of a surgical care bundle on the incidence of surgical site infection in colorectal surgery: a quasi-experimental intervention. *Rev Invest Clin* 2021;73:251–8.
- [28] Lohsiriwat V. High compliance with surgical site infection (SSI) prevention bundle reduces incisional SSI after colorectal surgery. *Ann Coloproctol* 2021;37:146–52.
- [29] D'Andrea AP, Khetan P, Miller R, Sylla P, Divino CM. Outcomes after bowel resection for inflammatory bowel disease in the era of surgical care bundles and enhanced recovery. *J Gastrointest Surg* 2020;24:123–31.
- [30] Martinez C, Omesiete P, Pandit V, Thompson E, Nocera M, Riall T, et al. A protocol-driven reduction in surgical site infections after colon surgery. *J Surg Res* 2020;246:100–5.
- [31] Park CM, Ha TS, Lee WY, Chung DR, Park YA, Choi JR, et al. Implementing a multidisciplinary care bundle to reduce colon surgical site infections. *Ann Surg Treat Res* 2020;99:285–93.
- [32] Hajirawala LN, Legare TB, Tiu SPT, DeKerlegand AM, Barton JS, Davis KG, et al. The impact of a colorectal care bundle for surgical site infections at an academic disproportionate share hospital with a level I trauma center. *Am Surg* 2020;86:848–55.
- [33] Dean HF, King E, Gane D, Hocking D, Rogers J, Pullyblank A. Introduction of a care bundle effectively and sustainably reduces patient-reported surgical site infection in patients undergoing colorectal surgery. *J Hosp Infect* 2020;105:156–61.
- [34] D'Souza K, Choi JI, Wootton J, Wallace T. Impact of sequential implementation of multimodal perioperative care pathways on colorectal surgical outcomes. *Can J Surg* 2019;62:25–32.
- [35] Guzman-Pruneda FA, Husain SG, Jones CD, Beal EW, Porter E, Grove M, et al. Compliance with preoperative care measures reduces surgical site infection after colorectal operation. *J Surg Oncol* 2019;119:497–502.
- [36] McGee MF, Kreutzer L, Quinn CM, Yang A, Shan Y, Halverson AL, et al. Leveraging a comprehensive program to implement a colorectal surgical site infection reduction bundle in a statewide quality improvement collaborative. *Ann Surg* 2019;270:701–11.
- [37] Albert H, Bataller W, Masroor N, Doll M, Cooper K, Spencer P, et al. Infection prevention and enhanced recovery after surgery: a partnership for implementation of an evidence-based bundle to reduce colorectal surgical site infections. *Am J Infect Control* 2019;47:718–9.
- [38] Hoang SC, Klipfel AA, Roth LA, Vrees M, Schechter S, Shah N. Colon and rectal surgery surgical site infection reduction bundle: to improve is to change. *Am J Surg* 2019;217:40–5.
- [39] Russell TA, Chung H, Riad C, Reardon S, Kazanjian K, Cherry R, et al. Sustaining improvement: implementation and spread of a surgical site infection bundle. *Am Surg* 2018;84:1665–9.
- [40] Vu JV, Collins SD, Seese E, Hendren S, Englesbe MJ, Campbell DA, et al. Evidence that a regional surgical collaborative can transform care: surgical site infection prevention practices for colectomy in Michigan. *J Am Coll Surg* 2018;226:91–9.
- [41] Lin DM, Carson KA, Lubomski LH, Wick EC, Pham JC. Statewide collaborative to reduce surgical site infections: results of the Hawaii surgical unit-based safety program. *Journal Am Coll Surg* 2018;227:189.
- [42] Weiser MR, Gonen M, Usiak S, Pottinger T, Samedy P, Pate D, et al. Effectiveness of a multidisciplinary patient care bundle for reducing surgical-site infections. *Br J Surg* 2018;105:1680–7.
- [43] Gachabayov M, Senagore AJ, Abbas SK, Yelika SB, You K, Bergamaschi R. Perioperative hyperglycemia: an unmet need within a surgical site infection bundle. *Tech Coloproctol* 2018;22:201–7.
- [44] Harris J. Success of a colorectal surgical site infection prevention bundle in a multihospital system. *AORN J* 2018;107:592–600.
- [45] Gorgun E, Rencuzogullari A, Ozben V, Stocchi L, Fraser T, Benlice C, et al. An effective bundled approach reduces surgical site infections in a high-outlier colorectal unit. *Dis Colon Rectum* 2018;61:89–98.
- [46] Jaffe TA, Meka AP, Semaan DZ, Okoro U, Hwang C, Papin J, et al. Optimizing value of colon surgery in Michigan. *Ann Surg* 2017;265:1178–82.
- [47] Kilan R, Moran D, Eid I, Okeahialam C, Quinn C, Binsaddiq W, et al. Improving antibiotic prophylaxis in gastrointestinal surgery patients: a quality improvement project. *Ann Med Surg* 2017;20:6–12.
- [48] Ohman KA, Wan LP, Guthrie T, Johnston B, Leinicke JA, Glasgow SC, et al. Combination of oral antibiotics and mechanical bowel preparation reduces surgical site infection in colorectal Surgery. *J Am Coll Surg* 2017;225:465–71.
- [49] Bert F, Giacomelli S, Amprino V, Pieve G, Ceresetti D, Testa M, et al. The 'bundle' approach to reduce the surgical site infection rate. *J Eval Clin Pract* 2017;23:642–7.
- [50] Hewitt DB, Tannouri SS, Burkhart RA, Altmark R, Goldstein SD, Isenberg GA, et al. Reducing colorectal surgical site infections: a novel, resident-driven, quality initiative. *Am J Surg* 2017;213:36–42.
- [51] Kwaan MR, Weight CJ, Carda SJ, Mills-Hokanson A, Wood E, Rivard-Hunt C, et al. Abdominal closure protocol in colorectal, gynecologic oncology, and urology procedures: a randomized quality improvement trial. *Am J Surg* 2016;211:1077–83.
- [52] Benlice C, Gorgun E. Using NSQIP data for quality improvement: the Cleveland Clinic SSI experience. *Semin Colon Rectal Surg* 2016;27:74–82.
- [53] Rumberger LK, Vittetoe D, Cathey L, Bennett H, Heidel RE, Daley BJ. Improving outcomes in elective colorectal surgery: a single-institution retrospective review. *Am Surg* 2016;82:325–30.

- [54] Tanner J, Kiernan M, Hilliam R, Davey S, Collins E, Wood T, et al. Effectiveness of a care bundle to reduce surgical site infections in patients having open colorectal surgery. *Ann R Coll Surg* 2016;98:270–4.
- [55] Keenan JE, Speicher PJ, Nussbaum DP, Adam MA, Miller TE, Mantyh CR, et al. Improving outcomes in colorectal surgery by sequential implementation of multiple standardized care programs. *J Am Coll Surg* 2015;221:404–414.e1.
- [56] Ghuman A, Chan T, Karimuddin AA, Brown CJ, Raval MJ, Phang PT. Surgical site infection rates following implementation of a colorectal closure bundle in elective colorectal surgeries. *Dis Colon Rectum* 2015;58:1078–82.
- [57] Pérez-Blanco V, García-Olmo D, Maseda-Garrido E, Nájera-Santos MC, García-Caballero J. Evaluation of a preventive surgical site infection bundle in colorectal surgery. *Cir Esp* 2015;93:222–8.
- [58] Waits SA, Fritze D, Banerjee M, Zhang WY, Kubus J, Englesbe MJ, et al. Developing an argument for bundled interventions to reduce surgical site infection in colorectal surgery. *Surgery* 2014;155:602–6.
- [59] Cima R, Dankbar E, Lovely J, Pendlimari R, Aronhalt K, Nehring S, et al. Colorectal surgery surgical site infection reduction program: a national surgical quality improvement program-driven multidisciplinary single-institution experience. *J Am Coll Surg* 2013;216:23–33.
- [60] Crolla R, van der Laan L, Veen EJ, Hendriks Y, van Schendel C, Kluytmans J. Reduction of surgical site infections after implementation of a bundle of care. *PLoS One* 2012;7:e44599.
- [61] Lutfiyya W, Parsons D, Breen J. A colorectal ‘care bundle’ to reduce surgical site infections in colorectal surgeries: a single-center experience. *Perm J* 2012;16:10–6.
- [62] Bull A, Wilson J, Worth LJ, Stuart RL, Gillespie E, Waxman B, et al. A bundle of care to reduce colorectal surgical infections: an Australian experience. *J Hosp Infect* 2011;78:297–301.
- [63] World Health Organization. Global guidelines on the prevention of surgical site infection. 2nd ed. Geneva: WHO; 2018. Available at: <https://apps.who.int/iris/bitstream/handle/10665/277399/9789241550475-eng.pdf?ua=1>.
- [64] National Institute for Health and Care Excellence. Surgical site infections: prevention and treatment. London: NICE; 2019.
- [65] Mangram AJ, Horan TC, Pearson ML, Silver LC, Jarvis WR. Guideline for prevention of surgical site infection, 1999. Centers for Disease Control and Prevention (CDC) Hospital Infection Control Practices Advisory Committee. *Am J Infect Control* 1999;27:97–132. quiz 3–4; discussion 96.
- [66] Gilhooly D, Green SA, McCann C, Black N, Moonesinghe SR. Barriers and facilitators to the successful development, implementation and evaluation of care bundles in acute care in hospital: a scoping review. *Implement Sci* 2019;14:47.
- [67] Resar R, Griffin FA, Haraden C, Nolan TW. Using care bundles to improve health care quality. IHI Innovation Series white paper. Cambridge, MA: Institute for Healthcare Improvement; 2012.
- [68] Pop-Vicas AE, Abad C, Baubie K, Osman F, Heise C, Safdar N. Colorectal bundles for surgical site infection prevention: a systematic review and meta-analysis. *Infect Control Hosp Epidemiol* 2020;41:805–12.
- [69] Wolfhagen N, Boldingh QJJ, Boermeester MA, de Jonge SW. Perioperative care bundles for the prevention of surgical-site infections: meta-analysis. *Br J Surg* 2022;109:933–42.
- [70] Berrios-Torres SI, Umscheid CA, Bratzler DW, Leas B, Stone EC, Kelz RR, et al. Centers for Disease Control and Prevention guideline for the prevention of surgical site infection, 2017. *JAMA Surg* 2017;152:784–91.