

Reliability and Validity of the Global Physical Activity Questionnaire for Portuguese Adults

Perceptual and Motor Skills
2024, Vol. 131(5) 1548–1570
© The Author(s) 2024



Article reuse guidelines:

sagepub.com/journals-permissions

DOI: 10.1177/00315125241266341

journals.sagepub.com/home/pms



Mariana Ribeiro¹ , Elisabete Fernandes², Mariana Borges¹,
Madalena Pires¹, Xavier Melo³, Fausto J. Pinto¹, Ana Abreu^{1,2,4},
and Rita Pinto^{1,2} 

Abstract

The Global Physical Activity Questionnaire (GPAQ) has been used often to assess physical activity (PA) patterns. However, the European Portuguese version of this instrument has not been validated. We aimed to validate the self-administered GPAQ, version 2, (GPAQv2) for Portuguese adults. We included 32 participants in a pilot study of a Portuguese adaptation of the test and 108 participants in an assessment of their PA patterns and sedentary behavior (SB) through the GPAQv2. For its validation, we compared the GPAQv2 to the International PA Questionnaire-Long Form (IPAQ-LF) (concurrent validity) and the *ActiGraph wGT3X-BT* accelerometer (criterion validity). We evaluated PA and SB at baseline and after seven consecutive days. Test-retest reliability with the Kappa test (k) and the Intraclass Correlation Coefficient (ICC) ranged from strong to almost perfect (k : 0.864–0.976) and from moderate to excellent (ICC: 0.56–0.994), respectively. Concurrent validity, assessed by Spearman's Correlation Coefficient,

¹Centro Cardiovascular da Universidade de Lisboa (CCUL@RISE), CAML, Faculdade de Medicina, Universidade de Lisboa, Lisboa, Portugal

²Instituto de Medicina Preventiva e Saúde Pública, Faculdade de Medicina, Universidade de Lisboa, Lisboa, Portugal

³Egas Moniz Interdisciplinary Research Center (CiiEM), Egas Moniz School of Health and Science, Campus Universitário, Almada, Portugal

⁴Instituto de Saúde Ambiental, Faculdade de Medicina, Universidade de Lisboa, Lisboa, Portugal

Corresponding Author:

Mariana Ribeiro, Exercise and Cardiovascular Rehabilitation Laboratory, CCUL, Faculdade de Medicina, Universidade de Lisboa, Av. Prof. Egas Moniz, Lisboa 1649-028, Portugal.

Email: marianafbr3@gmail.com

was moderate to substantial (ρ : 0.471–0.680), and there was fair to substantial criterion validity (ρ : 0.226–0.672). Bland-Altman plots showed that the GPAQv2 overestimated vigorous and moderate to vigorous PA and underestimated moderate PA. The largest difference values were related to SB, since the GPAQv2 underestimated sitting time. In sum, we found the GPAQv2 to have acceptable validity and reliability for assessing PA and SB patterns, and we recommend its use for Portuguese adults.

Keywords

physical activity, questionnaires, accelerometer, validity, comparison of methods

Introduction

Regular physical activity (PA) is a fundamental component of a salutogenic lifestyle, due to its multiple proven benefits in promoting health (Bull et al., 2009; WHO, 2020b) and preventing such chronic diseases as cardiovascular disease (Kyu et al., 2016; Tucker et al., 2022), type 2 diabetes (Fletcher et al., 2018), cancer (D'Ascenzi et al., 2021; Matthews et al., 2020) and other health outcomes (Angulo et al., 2020; Dipietro et al., 2019), including mental health (Barnes, 2015; Sawan et al., 2023). The World Health Organization (WHO) guidelines recommend that adults (18–64 years old) practice at least 150–300 minutes of moderate-intensity aerobic PA (MPA) or at least 75–150 minutes of vigorous-intensity aerobic PA (VPA) or an equivalent combination of moderate and vigorous-intensity PA (MVPA) each week to maintain or improve overall health and reduce the risk of or even prevent chronic disease (WHO, 2020b). However, over the last few decades there has been an exponential increase in physical inactivity levels in contemporary society (Nikitaro et al., 2021), leading to an increasingly sedentary lifestyle (Stamatakis et al., 2019). Furthermore, physical inactivity is currently considered the fourth risk factor for global mortality (WHO, 2010).

Monitoring population PA levels is only possible if there are reliable and valid PA assessment instruments (Armstrong & Bull, 2006). Subjective self-report methods, namely questionnaires, are among the most often used instruments in epidemiological studies (Sember et al., 2020), with the advantage of suitability for large samples of people and with low associated costs (Strath et al., 2013). Nevertheless, self-administered PA questionnaires can be cognitively difficult for respondents to complete (Janz, 2006), may induce social desirability bias (Strath et al., 2013), may use unfamiliar terms for gauging PA intensities (e.g.; moderate and vigorous), and may over or under-estimate patterns of PA (Sallis, 2010). To overcome these problems, accelerometers have been broadly used as an objective method of assessing the criterion validity of PA questionnaires (Sember et al., 2020; Skender et al., 2016).

Several PA questionnaires have been validated and frequently used (Strath et al., 2013) in different countries for public health monitoring and epidemiology research, including the International Physical Activity Questionnaire (IPAQ), with its long

and short forms (IPAQ-LF; IPAQ-SF) (Craig et al., 2003). While the IPAQ-LF was created to assess PA domains, it has been criticized as too long and complex; the shorter IPAQ-SF focuses on PA generally, rather than specifying its various sub-domains (Armstrong & Bull, 2006; Craig et al., 2003). The gap between the long and short forms of the IPAQ prompted the development of another reliable, valid and standardized questionnaire, taking into account cultural context, socioeconomic and/or literacy levels of respondents, and country-specific types of PA (Armstrong & Bull, 2006; Bull et al., 2009; Sember et al., 2020). The WHO prepared the Global Physical Activity Questionnaire (GPAQ) as part of their STEPwise approach to surveilling risk factors for chronic diseases. The IPAQ improved upon the long and short versions of the IPAQ, since the GPAQ evaluates PA in multiple domains (work, moving between places and leisure activities) and includes an assessment of sedentary behavior (SB) (Armstrong & Bull, 2006; WHO, 2005, 2022). Though the GPAQ, version 2, (GPAQv2) has been applied in over 100 countries (WHO, 2020a) and widely used (Mengesha et al., 2019; Whiting et al., 2021; WHO, 2011), there has been no translation and validation of it for the European Portuguese population. In this study, we aimed to validate and analyze the psychometric properties of the GPAQv2 for the Portuguese adult population, by comparing it to both an objective (accelerometry) and subjective (IPAQ-LF) method.

Method

Study Design

This was a cross-sectional, observational, and prospective study, consisting of two phases: a qualitative phase in which we translated and adapted the GPAQv2, and a pre-test and quantitative phase in which we tested the translated/adapted instrument's reliability and validity in a sample of Portuguese adults. In this second stage, we evaluated the PA level of all participants through two administrations of the GPAQv2 at two time points (M1 and M2), separated by seven consecutive days. We also compared the GPAQv2 Portuguese version with the respondents' completion of the IPAQ-LF and with ActiGraph *wGT3X-BT* accelerometer values. These two different correlates served as measures of concurrent and criterion validity, respectively.

Ethical Considerations

This study was authorized by the WHO, and the research protocol was approved, on 22 April 2022, by our local Ethics Committee (N.º 61/22). All participants were informed about the purpose of the study and provided written informed consent for the pilot test and qualitative and quantitative phases in person.

Sample Size

We conducted a power analysis (*G*Power 3.1.9.7*) to estimate a required sample size based on a previous study (Ács et al., 2020), in which investigators adapted and validated the GPAQv2 (by self-administration) for an apparently healthy Hungarian population and compared it with accelerometry and the IPAQ-LF. We assumed a Pearson's linear correlation coefficient of 0.48 for our findings and assumed a statistical power of 80% and a statistical significance level of 5%. The minimum necessary number of individuals for the quantitative phase was 108 individuals, ensuring equal distribution between sexes.

Participants

We first recruited a sample of 118 adults from the local Faculty of Medicine between May 2022 and September 2022. Inclusion criteria were: (a) age between 18 and 60 years old, (b) both females and males, (c) Portuguese nationality and resident in mainland Portugal, (d) availability to participate in the two assessment time points, (e) able to read and understand Portuguese, and (f) willingness to sign informed consent. Exclusion criteria were: (a) any physical or cognitive deficits that would interfere with engaging in activities of daily living; (b) being pregnant; and (c) being an athlete (defined as individuals who were simultaneously training in sports to improve their performance, actively participating in sports competitions, and registered in a local, regional or national sport federation with sport training and competition as their major activity or focus of personal interest to which they devoted several hours on all of most days, exceeding the time they allocated to other types of professional or league activities (Araújo & Scharhag, 2016). While all recruited participants met these inclusion/exclusion criteria, our final sample contained 108 participants, due to some missing data as explained below (Supplemental Material 1).

Procedures

Qualitative Phase. We conducted the instrument translation phase of this research by following WHO recommendations (WHO, 2005). Two qualified professionals translated the original questionnaire from English to European Portuguese, and a PA expert performed a back-translation into English. Subsequently, two PA expert analyzed all versions created and any discrepancies between the translations were discussed and a consensus was reached unanimously. The new version of the GPAQv2 received a final revision from a European Portuguese language teacher.

We performed a pilot test on this finalized new version of the GPAQv2, by self-administration, in a sample of 32 participants, homogeneously distributed by sex and age groups (18–60 years old). These respondents reported no difficulties understanding the questions, with average ratings of 4.75 in the 5-points on a Likert scale.

Quantitative Phase. We explained the study protocol to participants and obtained their written informed consent, and we then asked them to complete both the GPAQv2 and

IPAQ-LF digitally (via a link provided) and to wear an accelerometer for seven consecutive days. We collected their sociodemographic and anthropometric data through the provided link. At the end of the evaluation week, all participants were once again asked to complete both questionnaires and return the accelerometers to the research team. Participants received short message service reminders on the days they were scheduled to complete the questionnaires. A report of the results obtained was sent to each participant as a strategy for promoting PA and health literacy.

Physical Activity Outcome Measures

Global Physical Activity Questionnaire, Version 2 (GPAQv2). The GPAQv2 consists of 16 questions that assess adults' PA levels during a typical week. The questionnaire assesses three PA domains: (a) work, (b) travel to and from places, and (c) recreational activities and sedentary behavior (SB). The duration and frequency of PA and SB were evaluated in all domains in minutes/day. We used the GPAQ Analysis Guide (WHO, 2012) for scoring responses. We calculated the total MPA, VPA, MVPA and SB in minutes/week, and we converted the SB into hours/day, as this scale for this variable is easier to perceive and interpret for most people. Previous studies have also validated the GPAQv2 in other countries (Herrman et al., 2013; Keating et al., 2019).

International Physical Activity Questionnaire (IPAQ-LF). We used the self-administered IPAQ-LF Portuguese version (Campaniço, 2016) as a subjective method of testing the concurrent validity of the GPAQv2. This questionnaire contains 31 questions and was designed to assess PA in a usual week in the following domains: work, commuting/transportation between locations, household chores, general maintenance and family care, leisure activities and SB. Consistent with the organization of the GPAQv2, MPA, VPA, MVPA and SB were calculated in minutes/week, and SB in hours/day. The data was cleaned and processed according to the guidelines outlined in the IPAQ-LF manual (International Physical Activity Questionnaire group, 2004). The IPAQ is one of the most used questionnaires for assessing PA and several studies have already examined its validity and reliability. (Cerin et al., 2012; Cleland et al., 2018; Ryan et al., 2018; Van Holle et al., 2015).

ActiGraph wGT3X-BT. We measured PA objectively with triaxial *ActiGraph wGT3X-BT* (AG; *ActiGraph, Pensacola, FL, USA*) accelerometers. All participants were instructed to wear the accelerometer around their right hip (via an elastic belt) in the extension of the mid-axillary line to the iliac crest. Participants were instructed to wear the accelerometer throughout the day, except during water-related activities and during their sleep. The accelerometers were set to a frequency of 100 Hz, storing all information in 10 second epochs. At least three valid days, including one weekend day, with a minimum of 600 minutes of utilization, were required for a participant's accelerometer data to be included in subsequent data analysis. Accelerometer activation,

downloading, and data processing were carried out using *ActiLife Software version 6.13.4 (ActiGraph, Pensacola, FL, USA)*. Calculation of average time spent by participants in different types of PA intensity was carried out. We used the cut-off points by Troiano (Troiano et al., 2008) to distinguish PA intensity: sedentary (0–99 counts/min); light (100–2019 counts/min); moderate (2020–5998 counts/min) and vigorous (5999 counts/min). All variables were converted to time (minutes) per valid day.

Statistical Analysis

Our exploratory data analysis included the calculation of absolute frequencies and percentages for categorical variables; and we used means or *Ms* (and standard deviations or *SDs*) and medians (and interquartile ranges or *IQRs* as descriptors for numerical variables. We tested the normality of data distributions of continuous quantitative variables with the Kolmogorov-Smirnov test. Differences in the mean values of sociodemographic data between females and males were tested with the Student-t test for two independent samples or the Mann-Whitney U test for non-parametric data, as appropriate. We used the chi-square test for homogeneity for qualitative variables, or the Fisher's Exact test or the Fisher-Freeman-Halton test when the chi-square test could not be applied. We tested the reliability of categorical and continuous variables with the Cohen's Kappa coefficient (*k*) and Intraclass Correlation Coefficients test (ICC) - Model: Mixed two-factor; Type: Absolute agreement, respectively. The reliability ratings for *k* values were defined as: none (0–0.20), minimal (0.21–0.39), weak (0.40–0.59), moderate (0.60–0.79), strong (0.80–0.90) and almost perfect (>0.90) (McHugh, 2012). The ICC was categorized as: poor (<0.50), moderate (0.50–0.75), good (0.75–0.90) and excellent (>0.90) (Koo & Li, 2016). A reliability value of ≥ 0.70 was considered acceptable (Nunnally & Bernstein, 1994). Concurrent validity between questionnaires (GPAQv2 vs. IPAQ-LF) and criterion validity between GPAQv2-accelerometer were determined for all participants using the Spearman Correlation Coefficient (*rho*) with the following criteria: 0–0.2 = poor; 0.21–0.40 = fair; 0.41–0.60 = moderate; 0.61–0.80 substantial and 0.81–1.0 = near perfect (Bull et al., 2009). Agreement between the GPAQv2 and the accelerometer was assessed using Bland-Altman graphs, with the respective limits of agreement (Mean $\pm 1.96 \times SD$) at 95% (Bland & Altman, 1986). We performed all statistical analyses with the Statistical Package for Social Science (SPSS), version 28. We set statistical significance at 5% ($p < .05$).

Results

Participant Characteristics

Out of the initially recruited 118 eligible participants, 10 were excluded from data analyses for not using the accelerometer properly and/or failing to complete the questionnaires within the stipulated deadline. Table 1 shows the sociodemographic

characteristics of the remaining 108 participants whose average age was 39 years ($SD = 11$). In terms of health status, 80.6% ($n = 87$) of participants reported having no diagnosed chronic disease. Only one participant reported a diagnosed cardiovascular disease, and none mentioned having diabetes mellitus, cancer, chronic kidney disease and chronic obstructive pulmonary disease. Regarding PA levels (based on accelerometry data), 90.7% ($n = 98$) of the participants were physically active with an average of 49 minutes ($SD = 22$) MVPA/day. Their mean SB was 11.0 hours/day ($SD = 1.5$), and their mean number of valid accelerometer days was 6.9 ($SD = 0.4$). The PA levels of the participants evaluated using the three instruments, mentioned above, can be observed in [Table 2](#).

Reliability

The reliability study using the test-retest technique seven days apart were conducted on the categorical variables of the GPAQv2, and the results are presented in [Table 3](#). The

Table 1. Participant Characteristics.

<i>n</i>	Total	Male	Female	<i>p</i> -Value
	108	55	53	
	Mean (SD)	Mean (SD)	Mean (SD)	
Age (years)	39 (12)	39 (11)	39 (12)	.825
Weight (kg)	71.5 (12.3)	80.4 (8.9)	62.9 (8.5)	<.001
Height (meters)	1.70 (0.09)	1.76 (0.07)	1.63 (0.06)	<.001
BMI (kg/m ²)	24.7 (2.9)	25.8 (2.3)	23.6 (3.0)	<.001
Marital status <i>n</i> (%)				
Single	44 (40.70)	21 (39.60)	23 (41.80)	
Married	45 (41.70)	22 (41.50)	23 (41.80)	.988
Divorced	6 (5.60)	3 (5.70)	3 (5.50)	
Common law marriage	13 (12.00)	7 (13.20)	6 (10.90)	
Education level <i>n</i> (%)				
≤Middle school	7 (6.50)	3 (5.70)	4 (7.30)	
Middle school	44 (40.70)	20 (37.70)	24 (43.60)	.638
Higher education	57 (52.80)	30 (56.60)	27 (49.10)	
Risk factors <i>n</i> (%)				
Smoker	9 (8.30)	3 (5.70)	6 (10.90)	.489
Ex-smoker	9 (8.30)	7 (13.20)	2 (3.60)	.090
Dyslipidemia	12 (11.1)	6 (11.30)	6 (10.90)	.999
Family history of CVD	3 (2.80)	2 (1.90)	1 (1.80)	.614
HTN	7 (6.50)	3 (5.70)	4 (7.30)	.999

Note. Abbreviations: BMI, body mass index; CVD, cardiovascular disease; Fr, frequency; HTN, hypertension; *n*, sample size; SD, standard deviation.

value of k ranged from 0.864 to 0.976, indicating strong to almost perfect reliability. The percentage of agreement between the two evaluation moments was very high (94.4%–99.1%). For the continuous variables analysis, we used ICC for the frequency of PA practice (days/week and min/day) (Table 4). These values were statistically significant for all questions ($p < .001$) and ranged from moderate to excellent (ICC: 0.562–0.994). We found higher correlations regarding time in VPA, especially in the domains of work and leisure. On the other hand, the lowest ICC value corresponded to the duration of MPA in the leisure activities domain.

Concurrent and Criterion Validity

The criterion validity of the GPAQv2 was assessed against the accelerometer, while concurrent validity was tested against the IPAQ-LF (Supplemental Material 2). We found statistically significant differences for all PA levels and SB when comparing the results obtained from the three assessment instruments. The correlations between GPAQv2 and the accelerometer were positive and ranged from fair to substantial. The weakest association was verified for the SB, whereas the highest was for the MVPA. Comparing the GPAQv2 and IPAQ-LF, we found moderate to substantially positive correlations, where the highest association corresponded to MPA and the lowest to the SB.

Bland-Altman Analysis

Bland-Altman plots were used to measure the agreement between PA activity intensities assessed by GPAQv2 and the accelerometer records (Figures 1A–D). We observed significant bias between the two assessment instruments for all PA levels intensity and SB. Several outliers were identified, 6 for MPA, 8 VPA, 5 MVPA and 5 SB, with these data points falling outside the limits of agreement. For MPA, the mean bias between the two instruments was 16.05 minutes/week, with the range of variation from -327.91 to 295.80 . This suggests that the GPAQv2 underestimated PA for this level of intensity. For VPA, the mean bias was 109.67 minutes/week, with a range of variation from -204.14 to 423.48 , suggesting a tendency towards overestimation of GPAQv2 compared to the accelerometer. As for MVPA, the mean bias was 93.62 min/w, with a range of variation from -347.18 to 534.42 . The greatest mean difference was observed for SB which was 172.67 min/w (-468.74 ; 123.41) or 2.88 h/d (-7.81 ; 2.06). These results confirm the GPAQv2's tendency to underestimate this level of activity.

Discussion

In this study, we examined the reliability and validity of the self-administered GPAQv2 for assessing PA in Portuguese adults by comparing it with an accelerometer (objective method) and the self-administered IPAQ-LF (subjective method). The GPAQv2 showed acceptable and statistically significant reliability on test-retest

Table 2. Pattern of Physical Activity of the Sample According to Accelerometry, GPAQv2 and IPAQ-LF.

PA intensity	Accelerometer			GPAQv2			IPAQ-LF		
	Mean (SD)	Median (IQR)	Mean (SD)	Median (IQR)	Mean (SD)	Median (IQR)	Mean (SD)	Median (IQR)	
PA intensity									
MPA (min/w)	310.7 (133.3)	290.3 (171.4)	294.6 (209.0)	240.0 (270.0)	554.5 (691.7)	335.0 (601.0)			
VPA (min/w)	28.2 (54.6)	9.3 (31.5)	137.8 (172.9)	95.0 (180.0)	219.7 (400.5)	120.0 (278.0)			
MVPA (min/w)	338.8 (156.4)	322.2 (195.3)	432.5 (284.4)	375.0 (251.0)	774.2 (893.2)	515.0 (784.0)			
SB (h/d)	11.0 (1.5)	11.1 (2.1)	8.2 (2.5)	8.0 (3.4)	8.1 (3.0)	7.6 (3.6)			

Note. Abbreviations: GPAQv2, global physical activity questionnaire, version 2; h/d, hours per day; IPAQ-LF, international physical activity questionnaire, long form; IQR, interquartile range; Min/w, minutes per week; MPA, moderate physical activity; MVPA, moderate to vigorous physical activity; PA, physical activity; SB, sedentary behavior; SD, standard deviation; VPA, vigorous physical activity.

Table 3. Test-Retest Results for the Kappa Reliability Value.

<i>n</i> = 108						
PA Domain		Question	Moment	k	% Agreement	<i>p</i> -Value
Work	VPA	Q1	Test (M1)	0.947	99.074	<.001
			Retest (M2)			
	MPA	Q4	Test (M1)	0.976	99.074	<.001
			Retest (M2)			
Transport	MPA	Q7	Test (M1)	0.864	94.444	<.001
			Retest (M2)			
Leisure	VPA	Q10	Test (M1)	0.961	98.148	<.001
			Retest (M2)			
	MPA	Q13	Test (M1)	0.944	97.222	<.001
			Retest (M2)			

Note. Abbreviations: MPA, moderate physical activity; k, Kappa value; Q, question; %, percentage; PA, physical activity; VPA, vigorous physical activity.

correlations one week apart, with correlation values ranging from strong to almost perfect for *k* values, and moderate to excellent for the ICC. These values are consistent with those from other studies in different countries (Bull et al., 2009; Keating et al., 2019). For instance, Ács et al. (2020) reported an ICC: 0.899–0.987 in the adult Hungarian population, while Bull et al. (2009) found *k* values ranging from 0.67 to 0.73 across nine countries, with higher values in Shanghai. Rivière et al. (2018) reported poor to good reliability (ICC = 0.37–0.94; *k* = 0.50–0.62) in a French version of the GPAQ and Lee et al. (2020) found *k* values ranging from 0.30–0.67 and *rho* between 0.47–0.70, in Korea. Reliability results across studies may have oscillated due to several factors, including the wide age range of our target population (e.g., 15–79 years old) (Keating et al., 2019), cultural and socioeconomic differences between developed and developing countries (Bull et al., 2009; Keating et al., 2019), and variations in the participants' education and health levels (Alkahtani, 2016; Bull et al., 2009; Mumu et al., 2017). In our study, 52.80% of the participants were highly educated. Similar findings were reported by Lee et al. (2011), who found that participants with higher education were less likely to overestimate their PA levels on the IPAQ-SF. We found the reliability of self-reported VPA to be highest for the work and leisure domains, likely because of the routine nature of work activities facilitated recall when completing the GPAQv2 (Adıgüzel et al., 2021). VPA in leisure activities, was also more reliably reported, as this intensity level is typically achieved through easier to recall structured recreational exercises (Chu et al., 2015; Meh et al., 2021). In contrast, activities such as moving between places or practicing MPA in daily life may be more challenging to accurately report (Chu et al., 2015).

Our concurrent validity findings showed moderate to substantial correlations between GPAQ-IPAQ PA intensity scores, closer to Rivière et al. (2018) results (*rho*: 0.410–

Table 4. Test-Retest Reliability Analysis with Intraclass Correlation Coefficients (ICCs).

n = 108

PA Domain		Question	Moment	ICC	<i>p</i> -Value
Work	VPA	Q2	Score -test (M1)	0.991	<.001
			Score – retest (M2)		
	MPA	Q3	Score - test (M1)	0.989	<.001
			Score – retest (M2)		
		Q5	Score - test (M1)	0.981	<.001
			Score – retest (M2)		
Q6	Score - test (M1)	0.978	<.001		
	Score – retest (M2)				
Transport	MPA	Q8	Score - test (M1)	0.971	<.001
			Score – retest (M2)		
		Q9	Score - test (M1)	0.938	<.001
Leisure	VPA	Q11	Score - test (M1)	0.994	<.001
			Score – retest (M2)		
		Q12	Score - test (M1)	0.989	<.001
	Score – retest (M2)				
	MPA	Q14	Score - test (M1)	0.965	<.001
			Score – retest (M2)		
Q15		Score - test (M1)	0.562	<.001	
SB		Q16	Score - test (M1)	0.973	<.001
			Score – retest (M2)		

Note. Abbreviations: ICC, intraclass correlation coefficients; MPA, moderate physical activity; Q, question; SB, sedentary behavior; VPA, vigorous physical activity.

0.860) than Herrman et al. (2013) (ρ : 0.260–0.630). These differences in results may be due to variations in the IPAQ form used (Rivière et al., 2018). The short form (IPAQ-SF), used in some studies, lacks the ability to differentiate PA by domains, whereas the IPAQ-LF and GPAQv2 assess PA by domains, which may facilitate participants' recall (Armstrong & Bull, 2006; Meh et al., 2021). However, the IPAQ-LF has been criticized for being too long (Armstrong & Bull, 2006), potentially leading to PA overestimation. This highlights the importance of using the same questionnaire when comparing PA data across populations (Rivière et al., 2018). SB showed the strongest association between questionnaires, consistent with Bull et al. (2009) findings ($\rho = 0.650$, $p < .01$) and Adıgüzel et al. (2021) findings ($\rho = 0.940$, $p < .001$). A possible explanation for these similarities is that SB is similarly approached in both questionnaires (Adıgüzel et al., 2021). However, the single SB question in subjective PA assessments (GPAQv2 and IPAQ-SF) can lead to underestimation with these instruments (Meh et al., 2021),




Figure 1. Bland-Altman Plots of the Criterion Validity of the Global Physical Activity Questionnaire, Version 2.

Note. (A) – Agreement GPAQv2 versus accelerometer for moderate physical activity (MPA); (B) agreement GPAQv2 versus accelerometer for vigorous physical activity (VPA); (C) agreement GPAQv2 versus Accelerometer for Moderate to vigorous physical activity (MVPA) and (D) agreement GPAQv2 versus accelerometer for sedentary behavior (SB). GPAQv2 – global physical activity questionnaire, version 2; h/d, hours per day; IPAQ-LF, international physical activity questionnaire, long form; min/w, minutes per week.




Figure 1. Continued.

suggesting that the IPAQ-LF, with its two SB questions, distinguishing between weekdays and weekends may be favored for SB (Craig et al., 2003).

Our criterion validity findings showed that accelerometer-assessed sedentary time was significantly higher than self-reported SB in the GPAQv2. This is consistent with previous findings (Ács et al., 2020; Cleland et al., 2014; Lee et al., 2020; Meh et al., 2021). We

found a significant correlation ($\rho = 0.226, p = .019$) between these measures, falling within the range identified in a recent systematic review ($\rho: 0.007\text{--}0.47$) (Keating et al., 2019), but our values were higher than that those reported by Cleland et al. (2014) ($\rho = 0.187, p = .135$) (2014), Lee et al. (2020) ($\rho = 0.18, p < .01$) (2020) and Ács et al. (2020) ($\rho = -0.007, p = .936$). However, Meh et al. (2021) obtained a higher correlation ($\rho = 0.400, p \leq 0.01$), using a different brand of accelerometers.

In our Bland-Altman analysis, the agreement between the GPAQv2 and accelerometer data was consistent with findings from other studies (Ács et al., 2020; Keating et al., 2019). Ács et al. (2020) reported an overestimation for the VPA (212.75 minutes/week) and MVPA (104.93 minutes/week) and underestimated SB time measured by an accelerometer (6336.79 minutes/week), while Meh et al. (2021) overestimated MVPA ($M = 64, SD = 143$ minutes) and underestimated SB time by an average of 151 minutes/day ($SD = 172$ or about 2.5 hours/day), a value similar to our data with a Portuguese population ($M = 172.67, SD = 151.06$ minutes/day or 2.88 hours/day). Discrepancies may be due to differences in measurement methods, as the GPAQv2 only assesses PA lasting at least 10 consecutive minutes, while accelerometers measure all movement (Rivière et al., 2018). Study design factors, such as the particular week of accelerometer use, could also contribute to variability (Keating et al., 2019). Finally, individual differences in PA literacy and subjective interpretation of bodily sensations during PA could affect accuracy between measures, particularly for MVPA (Canning et al., 2014; Meh et al., 2021). The mean difference in reported PA obtained and represented in the Bland-Altman graphs confirmed an PA overestimation tendency of the subjective self-report methods (especially for VPA) and SB underestimation by these methods (Keating et al., 2019), compared with objective methods (Colley, 2018). Our study has several strengths. The GPAQv2 demonstrated acceptable reliability with correlation values ranging from strong to almost perfect for k values, and moderate to excellent for the ICC. Furthermore, the self-administered GPAQv2 allows for ease of use and widespread application without the need for trained interviewers. Our use of a sample with a high proportion of highly educated participants provided insights into how education levels can influence the accuracy of PA self-reports. Additionally, we adhered to the WHO's standardized recommendations for validating the GPAQv2, adapting the example images to portray PA types (Bull et al., 2009) in the Portuguese cultural context.

Limitations and Directions for Future Research

This study is not without limitations. We opted for the self-administered version of the GPAQv2, noting that similar results have been obtained by these two methods in at least one prior study (Chu et al., 2015). Still, the original questionnaire was designed for use by a trained interviewer. Our participant sample was a convenience sample, and most participants were physically active, which may not be representative of the Portuguese population; latest reports concluded that 45% of adults in Portugal are physically inactive (OECD & WHO, 2023). The choice of our threshold upper age limit of 60 years of age was justified by a previous study (Wanner et al., 2017) that warned of a need for

cautiously interpreting PA for people over 60 years old, due to low correlations of self-reported and actual PA found for this sub-group. However, our choice limits generalization of these data to older individuals. Another limitation presented was our lack of control over the environment when the questionnaire was completed, since environmental factors can influence the ability to respond and concentrate. Similarly, we did not consider the average completion time (Vilelas, 2020), which may be useful for future time management in applying the GPAQv2. As for PA assessment methodology the accelerometer is not effective in evaluating certain activities such as cycling, swimming, lifting weights, or movements performed mostly with the upper limbs (Warren et al., 2010), including some domestic tasks (Hoos et al., 2012). Finally, the criterion of “10 consecutive minutes” present in the GPAQv2 may have been a barrier in self-reported PA assessment, as the accelerometer always evaluated continuously rather than within this time interval. Troiano et al. (2020) considered removing this rule and one recent study accounted for every minute of PA (WHO, 2020b). Future investigators might update the GPAQv2, since the WHO mentioned that it would be subject to revision for possible adjustments and such adjustments have now been recommended (WHO, 2020b) regarding removing the rule of counting PA in intervals of 10 consecutive minutes. Additionally, we recommend including reports of muscle strengthening activities in the GPAQv2, as these activities are not present in any question (Meh et al., 2021), despite multiple health benefits associated with this practice. It would be interesting to validate the GPAQv2 in more countries and different cultural contexts, and future researchers might apply hetero-administration, through an interview. In this sense, our study validated another PA assessment instrument, which could positively contribute to the monitoring of public health in Portugal (or in portuguese-speaking countries) where physical inactivity has a high prevalence of 45% (OECD & WHO, 2023). Its application could enable the characterization of the PA levels across different domains within populations, consequently leading to the development of health promotion policies and PA programs.

Conclusion

In this study, we provide valuable evidence on the psychometric properties of the GPAQv2 in the Portuguese cultural context. Our results demonstrate acceptable reliability, moderate to substantial concurrent validity with the IPAQ-LF, and reasonable to substantial criterion validity with accelerometry. However, self-reported PA levels were higher for VPA and MVPA and lower for MPA and SB compared to accelerometry. Our validation and test-retest reliability data enables the use of GPAQv2 among Portuguese adults to study PA within the WHO’s STEPwise approach for the surveillance of chronic disease risk factors.

Acknowledgments

We would like to express our gratitude to the WHO for authorizing the translation process of the GPAQv2, and to all participants who contributed for this study. Special thanks go to Maria Isabel

Barbosa, José Manuel Fiuza and Sérgio Carvalho for their support during the translation and cultural adaptation process of the GPAQv2.

Author Contributions

MR participated in the concept and design of the study, contributed to data collection and data statistical analysis, and drafting of the manuscript. EF contributed to data analysis and interpretation of results. MLP and MB participated in the design of the study and data collection. AA, FJP and XM participated in the design of the study and drafting of the manuscript. RP participated in the concept and design of the study, supervision and contributed to data collection. All authors contributed to the manuscript writing. All authors have read and approved the final version of the manuscript and agree with the order of presentation of the authors.

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

ORCID iDs

Mariana Ribeiro  <https://orcid.org/0000-0002-6811-5352>

Rita Pinto  <https://orcid.org/0000-0003-0925-4245>

Supplemental Material

Supplemental material for this article is available online.

References

- Ács, P., Betlehem, J., Oláh, A., Bergier, B., Morvay-Sey, K., Makai, A., & Prémusz, V. (2020). Cross-cultural adaptation and validation of the global physical activity questionnaire among healthy Hungarian adults. *BMC Public Health*, 20(S1), 1056. <https://doi.org/10.1186/s12889-020-08477-z>
- Adıgüzel, İ., Raika Durusoy Onmuş, İ., Mandıracıoğlu, A., & Aslı Öcek, Z. (2021). Adaptation of the global physical activity questionnaire (GPAQ) into Turkish: A validation and reliability study. *Turkish Journal of Physical Medicine and Rehabilitation*, 67(2), 175–186. <https://doi.org/10.5606/tftrd.2021.1675>
- Alkahtani, S. A. (2016). Convergent validity: Agreement between accelerometry and the global physical activity questionnaire in college-age Saudi men. *BMC Research Notes*, 9(1), 436. <https://doi.org/10.1186/s13104-016-2242-9>

- Angulo, J., El Assar, M., Álvarez-Bustos, A., & Rodríguez-Mañas, L. (2020). Physical activity and exercise: Strategies to manage frailty. *Redox Biology*, 35(101513), 101513. <https://doi.org/10.1016/j.redox.2020.101513>
- Araújo, C. G. S., & Scharhag, J. (2016). Athlete: A working definition for medical and health sciences research. *Scandinavian Journal of Medicine & Science in Sports*, 26(1), 4–7. <https://doi.org/10.1111/sms.12632>
- Armstrong, T., & Bull, F. (2006). Development of the World health organization global physical activity questionnaire (GPAQ). *Journal of Public Health*, 14(2), 66–70. <https://doi.org/10.1007/s10389-006-0024-x>
- Barnes, J. (2015). Exercise, cognitive function, and aging. *Advances in Physiology Education*, 39(2), 55–62. <https://doi.org/10.1152/advan.00101.2014>
- Bland, J. M., & Altman, D. G. (1986). Statistical methods for assessing agreement between two methods of clinical measurement. *Lancet*, 1(8476), 307–310. <https://www.sciencedirect.com/science/article/abs/pii/S0140673686908378?via%3Dihub>
- Bull, F., Maslin, T. S., & Armstrong, T. (2009). Global physical activity questionnaire (GPAQ): Nine country reliability and validity study. *Journal of Physical Activity and Health*, 6(6), 790–804. <https://doi.org/10.1123/jpah.6.6.790>
- Campaniço, H. (2016). *Validade simultânea do questionário internacional de actividade física através da medição objectiva da actividade física por actigrafia proporcional*. [Dissertação elaborada com vista à obtenção do Grau de Mestre em Exercício e Saúde, Faculdade de Motricidade Humana da Universidade de Lisboa]. https://www.repository.utl.pt/bitstream/10400.5/11866/1/DISSERTA%C3%87%C3%83O_2016_Helena_Campani%C3%A7o.pdf
- Canning, K. L., Brown, R. E., Jamnik, V. K., Salmon, A., Arden, C. I., & Kuk, J. L. (2014). Individuals underestimate moderate and vigorous intensity physical activity. *PLoS One*, 9(5), Article e97927. <https://doi.org/10.1371/journal.pone.0097927>
- Cerin, E., Barnett, A., Cheung, M. C., Sit, C. H. P., Macfarlane, D. J., & Chan, W. M. (2012). Reliability and validity of the IPAQ-L in a sample of Hong Kong urban older adults: Does neighborhood of residence matter? *Journal of Aging and Physical Activity*, 20(4), 402–420. <https://doi.org/10.1123/japa.20.4.402>
- Chu, A. H. Y., Ng, S. H. X., Koh, D., & Müller-Riemenschneider, F. (2015). Reliability and validity of the self- and interviewer-administered versions of the global physical activity questionnaire (GPAQ). *PLoS One*, 10(9), Article e0136944. <https://doi.org/10.1371/journal.pone.0136944>
- Cleland, C., Ferguson, S., & Ellis, G. (2018). Validity of the International Physical Activity Questionnaire (IPAQ) for assessing moderate-to-vigorous physical activity and sedentary behaviour of older adults in the United Kingdom. *BMC Medical Research Methodology*, 18(1), 176. <https://doi.org/10.1186/s12874-018-0642-3>
- Cleland, C. L., Hunter, R. F., Kee, F., Cupples, M. E., Sallis, J. F., & Tully, M. A. (2014). Validity of the Global Physical Activity Questionnaire (GPAQ) in assessing levels and change in moderate-vigorous physical activity and sedentary behaviour. *BMC Public Health*, 14(1), 1255. <https://doi.org/10.1186/1471-2458-14-1255>
- Colley, R. C. (2018). Comparison of self-reported and accelerometer-measured physical activity in Canadian adults. *Health Reports*, 29(12), 3–15. 30566204.

- Craig, C. L., Marshall, A. L., Sjoström, M., Bauman, A. E., Booth, M. L., Ainsworth, B. E., Pratt, M., Ekelund, U., Yngve, A., Sallis, J. F., & Oja, P. (2003). International physical activity questionnaire: 12-country reliability and validity. *Medicine & Science in Sports & Exercise*, 35(8), 1381–1395. <https://doi.org/10.1249/01.MSS.0000078924.61453.FB>
- D'Ascenzi, F., Anselmi, F., Fiorentini, C., Mannucci, R., Bonifazi, M., & Mondillo, S. (2021). The benefits of exercise in cancer patients and the criteria for exercise prescription in cardiology. *European Journal of Preventive Cardiology*, 28(7), 725–735. <https://doi.org/10.1177/2047487319874900>
- Dipietro, L., Campbell, W. W., Buchner, D. M., Erickson, K. I., Powell, K. E., Bloodgood, B., Hughes, T., Day, K. R., Piercy, K. L., Vaux-Bjerke, A., & Olson, R. D. (2019). Physical activity, injurious falls, and physical function in aging: An umbrella review. *Medicine & Science in Sports & Exercise*, 51(6), 1303–1313. <https://doi.org/10.1249/MSS.0000000000001942>
- Fletcher, G. F., Landolfo, C., Niebauer, J., Ozemek, C., Arena, R., & Lavie, C. J. (2018). Promoting physical activity and exercise. *Journal of the American College of Cardiology*, 72(14), 1622–1639. <https://doi.org/10.1016/j.jacc.2018.08.2141>
- Herrmann, S. D., Heumann, K. J., Der Ananian, C. A., & Ainsworth, B. E. (2013). Validity and reliability of the global physical activity questionnaire (GPAQ). *Measurement in Physical Education and Exercise Science*, 17(3), 221–235. <https://doi.org/10.1080/1091367X.2013.805139>
- Hoos, T., Espinoza, N., Marshall, S., & Arredondo, E. M. (2012). Validity of the global physical activity questionnaire (GPAQ) in adult latinas. *Journal of Physical Activity and Health*, 9(5), 698–705. <https://doi.org/10.1123/jpah.9.5.698>
- International Physical Activity Questionnaire Group. (2004). *Guidelines for data processing and analysis of the international physical activity questionnaire (IPAQ) – short and long forms*. International Physical Activity Questionnaire Group. https://www.physio-pedia.com/images/c/c7/Quidelines_for_interpreting_the_IPAQ.pdf
- Janz, K. F. (2006). Physical activity in epidemiology: Moving from questionnaire to objective measurement. *British Journal of Sports Medicine*, 40(3), 191–192. <https://doi.org/10.1136/bjism.2005.023036>
- Keating, X. D., Zhou, K., Liu, X., Hodges, M., Liu, J., Guan, J., Phelps, A., & Castro-Piñero, J. (2019). Reliability and concurrent validity of global physical activity questionnaire (GPAQ): A systematic review. *International Journal of Environmental Research and Public Health*, 16(21), 4128. <https://doi.org/10.3390/ijerph16214128>
- Koo, T. K., & Li, M. Y. (2016). A guideline of selecting and reporting Intraclass correlation coefficients for reliability research. *Journal of Chiropractic Medicine*, 15(2), 155–163. <https://doi.org/10.1016/j.jcm.2016.02.012>
- Kyu, H. H., Bachman, V. F., Alexander, L. T., Mumford, J. E., Afshin, A., Estep, K., Veerman, J. L., Delwiche, K., Iannarone, M. L., Moyer, M. L., Cercy, K., Vos, T., Murray, C. J. L., & Forouzanfar, M. H. (2016). Physical activity and risk of breast cancer, colon cancer, diabetes, ischemic heart disease, and ischemic stroke events: Systematic review and dose-response meta-analysis for the global burden of disease study 2013. *BMJ*, 354, i3857. <https://doi.org/10.1136/bmj.i3857>

- Lee, J., Lee, C., Min, J., Kang, D.-W., Kim, J.-Y., Yang, H. I., Park, J., Lee, M.-K., Lee, M., Park, I., Jae, S. Y., Jekal, Y., Jee, S. H., & Jeon, J. Y. (2020). Development of the Korean global physical activity questionnaire: Reliability and validity study. *Global Health Promotion, 27*(3), 44–55. <https://doi.org/10.1177/1757975919854301>
- Lee, P. H., Yu, Y., McDowell, I., Leung, G. M., Lam, T., & Stewart, S. M. (2011). Performance of the international physical activity questionnaire (short form) in subgroups of the Hong Kong Chinese population. *International Journal of Behavioral Nutrition and Physical Activity, 8*(1), 81. <https://doi.org/10.1186/1479-5868-8-81>
- Matthews, C. E., Moore, S. C., Arem, H., Cook, M. B., Trabert, B., Håkansson, N., Larsson, S. C., Wolk, A., Gapstur, S. M., Lynch, B. M., Milne, R. L., Freedman, N. D., Huang, W.-Y., Berrington de Gonzalez, A., Kitahara, C. M., Linet, M. S., Shiroma, E. J., Sandin, S., Patel, A. V., & Lee, I.-M. (2020). Amount and intensity of leisure-time physical activity and lower cancer risk. *Journal of Clinical Oncology, 38*(7), 686–697. <https://doi.org/10.1200/JCO.19.02407>
- McHugh, M. L. (2012). Interrater reliability: The kappa statistic. *Biochemia Medica, 22*(3), 276–282. <https://doi.org/10.11613/BM.2012.031>
- Meh, K., Jurak, G., Sorić, M., Rocha, P., & Sember, V. (2021). Validity and reliability of IPAQ-SF and GPAQ for assessing sedentary behaviour in adults in the European union: A systematic review and meta-analysis. *International Journal of Environmental Research and Public Health, 18*(9), 4602. <https://doi.org/10.3390/ijerph18094602>
- Meh, K., Sember, V., Đurić, S., Vähä-Ypyä, H., Rocha, P., & Jurak, G. (2021). Reliability and validity of slovenian versions of IPAQ-SF, GPAQ, and EHIS-PAQ for assessing physical activity and sedentarism of adults. *International Journal of Environmental Research and Public Health, 19*(1), 430. <https://doi.org/10.3390/ijerph19010430>
- Mengesha, M. M., Roba, H. S., Ayele, B. H., & Beyene, A. S. (2019). Level of physical activity among urban adults and the socio-demographic correlates: A population-based cross-sectional study using the global physical activity questionnaire. *BMC Public Health, 19*(1), 1160. <https://doi.org/10.1186/s12889-019-7465-y>
- Mumu, S. J., Ali, L., Barnett, A., & Merom, D. (2017). Validity of the global physical activity questionnaire (GPAQ) in Bangladesh. *BMC Public Health, 17*(1), 650. <https://doi.org/10.1186/s12889-017-4666-0>
- Nikitara, K., Odani, S., Demenagas, N., Rachiotis, G., Symvoulakis, E., & Vardavas, C. (2021). Prevalence and correlates of physical inactivity in adults across 28 European countries. *The European Journal of Public Health, 31*(4), 840–845. <https://doi.org/10.1093/eurpub/ckab067>
- Nunnally, J., & Bernstein, I. (1994). The assessment of reliability. Standards of reliability. In *Psychometric theory (3a)*. McGraw Hill.
- OECD, & World Health Organization. (2023). *Step up! Tackling the burden of insufficient physical activity in Europe*. WHO. <https://www.oecd-ilibrary.org/content/publication/500a9601-en>
- Rivière, F., Widad, F. Z., Speyer, E., Erpelding, M.-L., Escalon, H., & Vuillemin, A. (2018). Reliability and validity of the French version of the global physical activity questionnaire.

- Journal of Sport and Health Science*, 7(3), 339–345. <https://doi.org/10.1016/j.jshs.2016.08.004>
- Ryan, D. J., Wullems, J. A., Stebbings, G. K., Morse, C. I., Stewart, C. E., & Onambele-Pearson, G. L. (2018). Reliability and validity of the international physical activity questionnaire compared to calibrated accelerometer cut-off points in the quantification of sedentary behaviour and physical activity in older adults. *PLoS One*, 13(4), Article e0195712. <https://doi.org/10.1371/journal.pone.0195712>
- Sallis, J. F. (2010). Measuring physical activity: Practical approaches for program evaluation in native American communities. *Journal of Public Health Management and Practice*, 16(5), 404–410. <https://doi.org/10.1097/PHH.0b013e3181d52804>
- Sawan, S. A., Nunes, E. A., Lim, C., McKendry, J., & Phillips, S. M. (2023). The health benefits of resistance exercise: Beyond hypertrophy and big weights. *Exercise, Sport, and Movement*, 1(1), 1–5. <https://doi.org/10.1249/ESM.0000000000000001>
- Sember, V., Meh, K., Sorić, M., Starc, G., Rocha, P., & Jurak, G. (2020). Validity and reliability of international physical activity questionnaires for adults across EU countries: Systematic review and meta analysis. *International Journal of Environmental Research and Public Health*, 17(19), 7161. <https://doi.org/10.3390/ijerph17197161>
- Skender, S., Ose, J., Chang-Claude, J., Paskow, M., Brühmann, B., Siegel, E. M., Steindorf, K., & Ulrich, C. M. (2016). Accelerometry and physical activity questionnaires—a systematic review. *BMC Public Health*, 16(1), 515. <https://doi.org/10.1186/s12889-016-3172-0>
- Stamatakis, E., Gale, J., Bauman, A., Ekelund, U., Hamer, M., & Ding, D. (2019). Sitting time, physical activity, and risk of mortality in adults. *Journal of the American College of Cardiology*, 73(16), 2062–2072. <https://doi.org/10.1016/j.jacc.2019.02.031>
- Strath, S. J., Kaminsky, L. A., Ainsworth, B. E., Ekelund, U., Freedson, P. S., Gary, R. A., Richardson, C. R., Smith, D. T., & Swartz, A. M. (2013). Guide to the assessment of physical activity: Clinical and research applications: A scientific statement from the American heart association. *Circulation*, 128(20), 2259–2279. <https://doi.org/10.1161/01.cir.0000435708.67487>
- Troiano, R. P., Berrigan, D., Dodd, K. W., Mâsse, L. C., Tilert, T., & McDowell, M. (2008). Physical activity in the United States measured by accelerometer. *Medicine & Science in Sports & Exercise*, 40(1), 181–188. <https://doi.org/10.1249/mss.0b013e31815a51b3>
- Troiano, R. P., Stamatakis, E., & Bull, F. C. (2020). How can global physical activity surveillance adapt to evolving physical activity guidelines? Needs, challenges and future directions. *British Journal of Sports Medicine*, 54(24), 1468–1473. <https://doi.org/10.1136/bjsports-2020-102621>
- Tucker, W. J., Fegers-Wustrow, I., Halle, M., Haykowsky, M. J., Chung, E. H., & Kovacic, J. C. (2022). Exercise for primary and secondary prevention of cardiovascular disease. *Journal of the American College of Cardiology*, 80(11), 1091–1106. <https://doi.org/10.1016/j.jacc.2022.07.004>
- Van Holle, V., De Bourdeaudhuij, I., Deforche, B., Van Cauwenberg, J., & Van Dyck, D. (2015). Assessment of physical activity in older Belgian adults: Validity and reliability of an adapted interview version of the long international physical activity questionnaire (IPAQ-L). *BMC Public Health*, 15(433), 1–14. <https://doi.org/10.1186/1471-2458-15-1>

- Vilelas, J. (2020). *Investigação—O Processo de Construção do Conhecimento (3ª)*. Edições Sílabo, Lda.
- Wanner, M., Hartmann, C., Pestoni, G., Martin, B. W., Siegrist, M., & Martin-Diener, E. (2017). Validation of the global physical activity questionnaire for self-administration in a European context. *BMJ Open Sport & Exercise Medicine*, 3(1), Article e000206. <https://doi.org/10.1136/bmjsem-2016-000206>
- Warren, J. M., Ekelund, U., Besson, H., Mezzani, A., Geladas, N., & Vanhees, L. (2010). Assessment of physical activity – a review of methodologies with reference to epidemiological research: A report of the exercise physiology section of the European association of cardiovascular prevention and rehabilitation. *European Journal of Cardiovascular Prevention & Rehabilitation*, 17(2), 127–139. <https://doi.org/10.1097/HJR.0b013e32832ed875>
- Whiting, S., Mendes, R., Abu-Omar, K., Gelius, P., Crispo, A., McColl, K., Simmonds, P., Fedkina, N., Andreasyan, D., Gahraman, H., Migal, T., Sturua, L., Obreja, G., Abdurakhmanova, Z., Saparkulovna, I. N., Erguder, T., Ekinci, B., Keskinkilic, B., Shukurov, S., & Breda, J. (2021). Physical inactivity in nine European and Central Asian countries: An analysis of national population-based survey results. *European Journal of Public Health*, 31(4), 846–853. <https://doi.org/10.1093/eurpub/ckab028>
- World Health Organization. (2020a). *Noncommunicable disease surveillance, monitoring and reporting—physical activity surveillance*. WHO. <https://www.who.int/teams/noncommunicable-diseases/surveillance/systems-tools/physical-activity-surveillance>
- World Health Organization. (2005). *WHO STEPS surveillance manual: The WHO stepwise approach to chronic disease risk factor surveillance*. World Health Organization. <https://apps.who.int/iris/handle/10665/43376>
- World Health Organization. (2010). *Global recommendations on physical activity for health* (Vol. 58). WHO.
- World Health Organization. (2011). *Review of physical activity surveillance data sources in European Union Member States*. WHO Regional Office for Europe. <https://apps.who.int/iris/handle/10665/108041>
- World Health Organization. (2012). *Global physical activity questionnaire (GPAQ). Analysis guide*. WHO. https://cdn.who.int/media/docs/default-source/ncds/ncd-surveillance/gpaq-analysis-guide.pdf?sfvrsn=1e83d571_2
- World Health Organization. (2020b). *Who guidelines on physical activity and sedentary behaviour*. World Health Organization. <https://www.who.int/publications/i/item/9789240015128>
- World Health Organization. (2022). *Global status report on physical activity 2022*. World Health Organization. <https://apps.who.int/iris/handle/10665/363607>

Author Biographies

Mariana Ribeiro is a registered nurse, with bachelor's degree in nursing from Nursing School of Lisbon and invited assistant professor. She has a master's degree in cardiovascular rehabilitation from the Faculty of Medicine of the University of Lisbon (FMUL) and has advanced training in long-term and palliative care. Her research

interests are physical activity, exercise, cardiovascular disease, cardio-oncology and health promotion.

Elisabete Fernandes is an invited assistant professor and researcher in biostatistics at the FMUL. She has a PhD in statistics at the Faculty of Sciences of the University of Lisbon. Her main research interests include biostatistics, clinical trials, epidemiological studies, and public health research.

Mariana Borges is an exercise physiologist who has been working in the field of cardiac rehabilitation since 2020. She holds a Bachelor's degree in Sports Science from the Faculty of Human Kinetics at the University of Lisbon and a Master's degree in Exercise and Health from the same institution. Currently, she is pursuing a Ph.D. at FMUL. She is also a researcher at Cardiovascular Exercise and Rehabilitation Laboratory from the Cardiovascular Center of the University of Lisbon (CCUL) and her main areas of interest are cardiovascular disease, cardiac rehabilitation, long-term adherence, exercise physiology, motivational interviewing, physical activity and sedentary behavior.

Madalena Pires is a certified exercise physiologist, researcher at the Exercise and Cardiovascular Rehabilitation Laboratory of CCUL and a PhD student at FMUL. She holds a Sports Science bachelor's degree and an Exercise and Health master's degree from the Faculty of Human Kinetics of the University of Lisbon. She specializes in exercise assessment and prescription for people with diagnosed cardiovascular disease. Her field of research targets physical activity and sedentary behavior, maintenance cardiovascular rehabilitation and the particularities of women with cardiovascular disease.

Xavier Melo holds a Ph.D. in Physical Activity and Health from the Faculty of Human Kinetics at the University of Lisbon. Currently is Associate Professor at Egas Moniz School of Health & Science. His research focuses on pediatric, chronic disease, and intellectual disability populations, with a dual emphasis on 1) exploring central and peripheral cardiovascular adaptations in both acute and chronic exercise response models, and 2) investigating associations between multiple cardiovascular risk factors and surrogate markers of cardiovascular disease. He exhibits significant expertise in multiple areas, including Vascular Imaging, Arterial Stiffness, Endothelial and Autonomic Function, Cardiopulmonary Exercise Testing, Energy Expenditure, Objective Measures of Body Composition, and Physical Activity.

Fausto J. Pinto is a senior cardiologist and full Professor of Cardiology at FMUL. He was the FMUL dean from 2015 to 2022. Currently he is the coordinator of the Cardiovascular Centre of University of Lisbon. He is the Director both of Cardiology department at Santa Maria Hospital, and the Department of Heart and Vessels of the Lisbon North Hospital Centre. During his career, devoted to cardiology clinical practice, teaching and research. His work has been awarded by multiple scientific and clinical societies, both national and international.

Ana Abreu is a senior cardiologist, Professor at FMUL, director of the Instituto de Medicina Preventiva and Instituto de Saúde Ambiental and elect-president of the European Association of Preventive Cardiology (EAPC) from the European Society of Cardiology (ESC). She possesses extensive knowledge in the cardiac rehabilitation field. Currently, she is the director of the cardiac rehabilitation program of CHULN/FMUL/CRECUL, first national center accredited in 2020 by the ESC. Her main focus, in clinical, research and advocacy are now: cardiovascular risk and prevention, Environmental and Planetary Health.

Rita Pinto is a researcher at the CCUL, an Exercise Physiologist at the Cardiovascular Rehabilitation Center of the University of Lisbon, and an invited assistant professor at FMUL. She is the coordinator of the Exercise and Cardiovascular Rehabilitation Laboratory at CCUL and a member of the scientific council of the RISE Laboratory – Health Research Network. She completed a degree in Sports Science and a Master’s degree in Exercise and Health. She received her PhD in Physical Activity and Health from Faculty of Human Kinetics at the University of Lisbon. In research, she focuses on cardiovascular disease, cardiac rehabilitation, exercise physiology, stress testing, assessment and prescription of physical exercise, physical activity, and sedentary behavior.